

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19816 A1

(51) International Patent Classification⁷: C07D 401/12,
413/12, 417/12, 205/04, 403/06, A61K 31/4535, A61P
9/10, 25/28, 19/10, 11/06, C07D 211/66, 207/16, 309/14,
335/02, 401/04

(21) International Application Number: PCT/US00/23584

(22) International Filing Date: 28 August 2000 (28.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/153,738 13 September 1999 (13.09.1999) US
60/222,900 3 August 2000 (03.08.2000) US

(71) Applicant (for all designated States except US):
**BOEHRINGER INGELHEIM PHARMACEUTI-
CALS, INC.** [US/US]; 900 Ridgebury Road, Ridgefield,
CT 06877 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **EMMANUEL,
Michel, J.** [HT/US]; 22B Myrtle Avenue, Danbury,
CT 06810 (US). **FRYE, Leah, L.** [US/US]; 53 Manor
Road, Patterson, NY 12563 (US). **HICKEY, Eugene, R.**
[US/US]; 5 Woodbury Drive, Danbury, CT 06811 (US).
LIU, Weimin [CN/US]; 56 Wakelee Avenue Ext. #12,
Shelton, CT 06484 (US). **MORWICK, Tina, M.** [US/US];
114 Taylor Road, New Milford, CT 06776 (US). **SPERO,**

Denice, M. [US/US]; 18 Limekiln Road, West Redding,
CT 06896 (US). **SUN, Sanxing** [CN/US]; 38 Padanaram
Avenue, 21B, Danbury, CT 06811 (US). **THOMSON,
David, S.** [GB/US]; 166 Minuteman Road, Ridgefield, CT
06877 (US). **WARD, Yancey, D.** [US/US]; 11 Narraganset
Trail, Sandy Hook, CT 06482 (US). **YOUNG, Erick, R.,
R.** [US/US]; 10232 Avalon Valley Drive, Danbury, CT
06810 (US).

(74) Agents: **RAYMOND, Robert** et al.; Boehringer Ingel-
heim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box
368, Ridgefield, CT 06877 (US).

(81) Designated States (national): AE, AU, BG, BR, BY, CA,
CN, CZ, EE, HR, HU, ID, IL, IN, JP, KR, KZ, LT, LV, MX,
NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, UZ, VN,
YU, ZA.


(84) Designated States (regional): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

Published:


- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.


(54) Title: NOVEL SPIROHETEROCYCLIC COMPOUNDS USEFUL AS REVERSIBLE INHIBITORS OF CYSTEINE PRO-
TEASES




(I)



(II)



(Ia)



(Ib)

(57) Abstract: Disclosed are novel cathepsin S, K, F, L and B reversible inhibitory compounds of formulas (I), (II), (Ia) and (Ib) further defined herein. The compounds are useful for treating autoimmune diseases. Also disclosed are processes for making such novel compounds.

WO 01/19816 A1

Novel Spiroheterocyclic Compounds Useful As Reversible Inhibitors of Cysteine Proteases

RELATED APPLICATION DATA

5

This application claims benefit to US Provisional Application serial nos. 60/153,738 filed 9/13/99 and 60/222,900 filed August 3, 2000.

TECHNICAL FIELD OF THE INVENTION

10

This invention relates to peptidyl spiroheterocyclic, amidino and guanidino compounds. The compounds are reversible inhibitors of the cysteine protease cathepsin S, K, F, L and B and are therefore useful in the treatment of autoimmune and other related diseases. The invention also relates to processes for preparing such compounds and pharmaceutical compositions comprising them.

15

BACKGROUND OF THE INVENTION

20

Cathepsin S and cathepsin K are members of the papain family, within the papain superfamily of cysteine proteases. The papain family is the largest group of cysteine proteases and includes proteases such as cathepsins B, H, K, L, O and S. (A.J. Barrett et al., 1996, Perspectives in Drug Discovery and Design, 6, 1). The cysteine proteases have important roles in human biology and diseases including atherosclerosis, emphysema, osteoporosis, chronic inflammation and immune disorders (H.A. Chapman et al., 1997, Ann. Rev. Physiol., 59, 63). Cathepsin S plays a key role in regulating antigen presentation and immunity (H.A. Chapman, 1998, Current Opinion in Immunology, 10, 93; R. J. Riese et al., 1998, J. Clin. Invest., 101, 2351; R.J. Riese et al., 1996, Immunity, 4, 357). Cathepsin S deficient mice have impaired invariant chain degradation resulting in decreased antigen presentation and germinal center formation, and diminished susceptibility to collagen-induced arthritis indicating the therapeutic potential for a

30

cathepsin S inhibitor (G. Shi et al., 1999, *Immunity*, 10, 197; T.Y. Nakagawa et al, 1999, *Immunity*, 10, 207)

The specificity of the immune response relies on processing of foreign protein and presentation of antigenic peptide at the cell surface. Antigenic peptide is presented bound to MHC Class II, a heterodimeric glycoprotein expressed in certain antigen presenting cells of hematopoietic lineage, such as B cells, macrophages and dendritic cells. Presentation of antigen to effector cells, such as T-cells, is a fundamental step in recognition of non-self and thus initiation of the immune response.

Recently MHC Class II heterodimers were shown to associate intracellularly with a third molecule designated invariant chain. Invariant chain facilitates Class II transport to the endosomal compartment and stabilizes the Class II protein prior to loading with antigen. Invariant chain interacts directly with Class II dimers in the antigen-binding groove and therefore must be proteolyzed and removed or antigen cannot be loaded or presented. Current research suggests that invariant chain is selectively proteolyzed by cathepsin S, which is compartmentalized with MHC Class II complexes within the cell. Cathepsin S degrades invariant chain to a small peptide, termed CLIP, which occupies the antigen – binding groove. CLIP is released from MHC Class II by the interaction of MHC Class II with HLA-DM, a MHC-like molecule thus freeing MHC Class II to associate with antigenic peptides. MHC Class II-antigen complexes are then transported to the cell surface for presentation to T-cells, and initiation of the immune response.

Cathepsin S, through proteolytic degradation of invariant chain to CLIP, provides a fundamental step in generation of an immune response. It follows that inhibition of antigen presentation via prevention of invariant chain degradation by cathepsin S could provide a mechanism for immuno-regulation. Control of antigen-specific immune responses has long been desirable as a useful and safe therapy for autoimmune diseases. Such diseases include Crohn's disease and arthritis, as well as other T-cell-mediated immune responses (C. Janeway and P. Travers, 1996, *Immunobiology, The Immune System in Health and Disease*, Chapter 12). Furthermore, cathepsin S, which has broad

pH specificity, has been implicated in a variety of other diseases involving extracellular proteolysis, such as Alzheimer's disease (U. Muller-Ladner et al., 1996, Perspectives in Drug Discovery and Design, 6, 87) and atherosclerosis (G.K. Sukhova et al., 1998, J. Clin. Invest., 102, 576).

5

A cathepsin S inhibitor has been found to block the rise in IgE titers and eosinophil infiltration in the lung in a mouse model of pulmonary hypersensitivity, suggesting that cathepsin S may be involved in asthma (R.J. Riese et al., J. Clin. Investigation, 1998, 101, 2351).

10

Another cysteine protease, cathepsin F has been found in macrophages and is also involved in antigen processing. It has been postulated that cathepsin F in stimulated lung macrophages and possibly other antigen presenting cells could play a role in airway inflammation (G.-P. Shi et al., J. Exp. Med., 2000, 191, 1177).

15

Cathepsin K, another cysteine protease has been found to be highly expressed in osteoclasts and to degrade bone collagen and other bone matrix proteins. Inhibitors of cathepsin K have been shown to inhibit bone resorption in mice. Therefore, cathepsin K may play a role in osteoclastic bone resorption and cathepsin K inhibitors may be useful in the treatment of diseases involving bone resorption such as osteoporosis (F. Lazner et al., Human Molecular Genetics, 1999, 8, 1839).

20

Cysteine proteases are characterized by having a cysteine residue at the active site which serves as a nucleophile. The active site also contains a histidine residue. The imidazole ring on the histidine serves as a base to generate a thiolate anion on the active site cysteine, increasing its nucleophilicity. When a substrate is recognized by the protease, the amide bond to be cleaved is directed to the active site, where the thiolate attacks the carbonyl carbon forming an acyl-enzyme intermediate and cleaving the amide bond, liberating an amine. Subsequently, water cleaves the acyl-enzyme species regenerating the enzyme and liberating the other cleavage product of the substrate, a carboxylic acid.

25

30

Inhibitors of cysteine proteases contain a functionality that can react reversibly or irreversibly with the active site cysteine. Examples of reactive functionalities that have been described (D. Rasnick, 1996, Perspectives in Drug Discovery and Design, 6, 47) on cysteine protease inhibitors include peptidyl diazomethanes, epoxides,

- 5 monofluoroalkanes and acyloxymethanes, which irreversibly alkylate the cysteine thiol. Other irreversible inhibitors include Michael acceptors such as peptidyl vinyl esters and other carboxylic acid derivatives (S. Liu et al., J. Med Chem., 1992, 35, 1067) and vinyl sulfones (J.T. Palmer et al., 1995, J. Med Chem., 38, 3193).
- 10 Reactive functionalities that form reversible complexes with the active site cysteine include peptidyl aldehydes (R.P. Hanzlik et al., 1991, Biochim. Biophys. Acta., 1073, 33), which are non-selective, inhibiting both cysteine and serine proteases as well as other nucleophiles. Peptidyl nitriles (R.P. Hanzlik et al., 1990, Biochim. Biophys. Acta., 1035, 62) are less reactive than aldehydes and therefore more selective for the more
- 15 nucleophilic cysteine proteases. Various reactive ketones have also been reported to be reversible inhibitors of cysteine proteases (D. Rasnick, 1996, *ibid*). In addition to reacting with the nucleophilic cysteine of the active site, reactive ketones may react with water, forming a hemiketal which may act as a transition state inhibitor.
- 20 Examples of cathepsin S inhibitors have been reported. J.L. Klaus et al. (WO 9640737) described reversible inhibitors of cysteine proteases including cathepsin S, containing an ethylene diamine. In US Patent No. 5,776,718 to Palmer et al. there is disclosed in it's broadest generic aspect a protease inhibitor comprising a targeting group linked through a two carbon atom chain to an electron withdrawing group (EWG). The compounds of the
- 25 present application are structurally distinct and thus excluded from the 5,776,718 patent with particular embodiments possessing unexpectedly greater activity than the closest compounds of the prior art. Other examples of cathepsin S inhibitors have been reported by E.T. Altmann et al, (WO 9924460, 1999) which describes dipeptide nitriles asserted to have activity as inhibitors of Cathepsins B, K, L and S. The WO publication does not
- 30 disclose any compounds possessing an imino or guanidino moiety and fails to provide

any description, methods or examples for particular spiroheterocyclic moieties at the P2 position.

Additional peptidyl nitriles have been reported as protease inhibitors. For example, both
5 nitriles and ketoheterocycles are described by B.A. Rowe et al. (US 5,714,471) as
protease inhibitors useful in the treatment of neurodegenerative diseases. Peptidyl nitriles
are reported by B. Malcolm et al. (WO 9222570) as inhibitors of picornavirus protease.
B.J. Gour-Salin (Can. J. Chem., 1991, 69, 1288) and T.C. Liang (Arch. Biochim.
Biophys., 1987, 252, 626) described peptidyl nitriles as inhibitors of papain

10 A reversible inhibitor presents a more attractive therapy than irreversible inhibitors. Even
compounds with high specificity for a particular protease can bind non-target enzymes.
An irreversible compound could therefore permanently inactivate a non-target enzyme,
increasing the likelihood of toxicity. Furthermore, any toxic effects resulting from
15 inactivation of the target enzyme would be mitigated by reversible inhibitors, and could
be easily remedied by modified or lower dosing. Finally, covalent modification of an
enzyme by an irreversible inhibitor could potentially generate an antibody response by
acting as a hapten.

20 In light of the above, there is a clear need for compounds which reversibly and selectively
inhibit cysteine proteases cathepsin S, K, F, L and B for indications in which these
proteases exacerbate disease.

25 BRIEF DESCRIPTION OF THE INVENTION

It is therefore an object of this invention to provide novel compounds according to the
formulas (I), (II), (Ia) and (Ib) as described herein which reversibly inhibit the cysteine
30 proteases cathepsin S, K, F, L and B. It is a further object of the invention to provide
methods for treating diseases and pathological conditions exacerbated by these cysteine
proteases such as, but not limited, to rheumatoid arthritis, multiple sclerosis, asthma and


osteoporosis. It is yet a further object of the invention to provide novel processes for preparation of the above-mentioned novel compounds.

5 DETAILED DESCRIPTION OF THE INVENTION

A proposed mechanism of action of the cysteine protease inhibitors of this invention is that the inhibitors contain a functionality that can react (reversibly or irreversibly) with the active site cysteine. The reactive functionality is attached to a peptide or peptide mimic that can be recognized and accommodated by the region of the protease surrounding the active site. The nature of both the reactive functionality and the remaining portion of the inhibitor determine the degree of selectivity and potency toward a particular protease.

15 Given the similarity of the active sites in cysteine proteases, it may be anticipated that a given class of inhibitors might have activity against more than one cysteine protease. It may also be expected that due to structural differences between individual cysteine proteases, different compounds of the invention may have different inhibitory potencies against different cysteine proteases. Thus some of the compounds of the invention may
20 also be expected to be most effective in treating diseases mediated by cysteine proteases that they inhibit most potently. The activity of particular compounds disclosed herein against cysteine proteases cathepsin S, K, F, L and B may be determined by the screens described in the section entitled "Assessment of Biological Properties."

25 In one broad generic aspect, the invention provides novel compounds of the formula (I):



wherein:

Het is piperidinyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, hexahydropyrimidinyl, hexahydropyridazinyl, piperazinyl, 1,4,5,6-
5 tetrahydropyrimidin-2-ylamine, dihydro-oxazolyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide or imidazolidinyl-2,4-dione, each being optionally substituted with one or more R₅;

10 Y is O or S;

R₁ is C1-5 alkyl, C1-5 alkoxy, aryloxy, C3-7 cycloalkyl, phenyl, benzyl, naphthyl, tetrahydronaphthyl, C1-5alkylsulfonylC1-5alkyl, C3-7cycloalkylsulfonylC1-5alkyl,
15 arylsulfonylC1-5alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl, quinoxalinyl, or amino; wherein R₁ is optionally substituted by one or
20 more R_a;

R_a is C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl,
25 pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkoxy, C1-5alkanoyl, C1-5alkanoyloxy, aryloxy, benzyloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl,
30 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl,

pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio wherein the

5 sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, C1-5
10 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoxyloxy, arylcarbamoxyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, pyrrolidinyl, piperidinyl,
15 morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_a may be further optionally
20 substituted by one or more R_b ;

R_b is C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino;

25 R_2 is hydrogen or C1-3 alkyl;

R_3 is hydrogen, C1-5 alkyl, C2-5alkylene, C3-7 cycloalkyl, arylC1-3alkyl or aryl wherein R_3 is optionally substituted by one or more R_c ;

30 R_c is C1-5 alkyl, C3-7 cycloalkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl,

thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl,
pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl,
quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5
alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy,
5 aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or
di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl,
thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl,
thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl,
indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl,
10 isoquinolinyl, quinazolinyl or quinoxalinyl,
or R_c is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom
may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may
be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl,
15 morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl,
oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl,
pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl,
quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy,
20 arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5
alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be
independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl,
piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl,
pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl,
25 pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl,
benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, halogen,
hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further
optionally substituted by one or more R_d;

R₄ is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylalkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino;

5

R₄ is hydrogen or C1-3 alkyl;

R₅ is C1-5 alkyl chain optionally interrupted by one or two O or S, phenyl, naphthyl,
10 arylC1-3alkyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, C1-5
alkanoyl, aroyl, C1-5 alkoxy carbonyl, aryloxy carbonyl, benzyloxy carbonyl, carbamoyl
wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl,
aryl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl, pyrrolyl,
thiazolyl, imidazolyl, pyridinyl, benzimidazolyl or quinolinyl,
15 or R₅ is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be
oxidised to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidised to a
sulfoxide or sulfone, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5
alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be
independently mono or disubstituted by alkyl, aryl, pyrrolidinyl, piperidinyl,
20 morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl,
thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyridinyl carbonyl, pyrimidinyl,
pyrazinyl, indolyl, benzofuranyl, benzo thienyl, benzimidazolyl, benzthiazolyl, quinolinyl,
isoquinolinyl, quinazolinyl, quinoxalinyl or arylsulfonyl, or R₅ is halogen, hydroxy, oxo,
carboxy, cyano, nitro, amidino or guanidino, R₅ may be further optionally substituted by
25 one or more R₆;

R₆ is C1-5 alkyl, C3-6 cycloalkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl,
thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl
pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl,
30 indolyl, benzofuranyl, benzo thienyl, benzimidazolyl, benzthiazolyl, quinolinyl,
isoquinolinyl, quinazolinyl, benzoxazolyl, quinoxalinyl, C1-5 alkoxy, aryloxy,

aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, furanyl, thienyl, pyrrolyl or pyridinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, benzyloxy, arylC1-3alkoxycarbonyl, amidino or guanidino;

5

X is O or S and

pharmaceutically acceptable derivatives thereof.

10

In another embodiment of the invention, there are provided novel compounds of the formula (I) as described immediately above, and wherein:

15 Het is piperidinyl, pyrrolidinyl, tetrahydropyranyl or tetrahydrothiopyranyl each ring being substituted with one or more R₅;

Y is O;

20 R₁ is C1-3 alkyl, C1-3alkoxy, C3-7 cycloalkyl, phenyl, benzyl, naphthyl, tetrahydronaphthyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl, pyridinyl, isoxazolyl, pyrazinyl, indolyl, quinolinyl, benzofuranyl, benzimidazolyl, benzoisoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

25 R_a is C1-3 alkyl, phenyl, naphthyl, piperidinyl, indolinyl, morpholinyl, piperazinyl, furanyl, thienyl, benzimidazolyl, C1-3 alkoxy, C1-3 alkanoyl, phenoxy, naphthyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl or pyridinyl,
30 or R_a is C1-5 alkanoylamino, benzoylamino, C1-3 alkylsulfonyl, phenylsulfonyl, ureido wherein either nitrogen atom may be independently substituted by alkyl,

phenyl, piperidinyl, morpholinyl, furanyl, thienyl or pyridinyl, C1-3
alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, C1-5 alkylsulfonylamino,
phenylsulfonylamino, C1-5 alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-5
5 alkyl, phenyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl or pyridinyl,
halogen, hydroxy, oxo, carboxy, nitro or cyano, R_a may be further optionally
substituted by one or more R_b;

R_b is halogen, hydroxy, benzyloxy, oxo or cyano;

R₂ is hydrogen;

R₃ is C1-5 alkyl or C2-5 alkylene, C4-6 cycloalkyl or benzyl wherein R₃ is optionally
substituted by one or more R_c;

R_c is C1-4 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, C1-4 alkoxy, phenoxy,
benzoyl, benzyloxy, indolinyl, imidazolyl, C1-3alkylthio, C1-3alkylsulfonyl,
halogen, hydroxy, oxo, carboxy, nitro or cyano, R_c may be further optionally
substituted by one or more R_d;

R_d is methyl, phenyl, benzyl, benzyloxy, C1-3alkoxy, halogen, hydroxy,
nitro or cyano;

R₄ is hydrogen;

R₅ is C1-4alkyl chain optionally interrupted by one O or S atom, phenyl, phenylC1-
2alkyl, furanyl, pyrimidinyl, thienyl, C1-3 alkanoyl, benzoyl, C1-4 alkoxy carbonyl,
carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by
C1-5 alkyl, phenyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl or pyridinyl,
C1-3 alkylthio, phenylthio, C1-5 alkylaminosulfonyl, phenylaminosulfonyl, C1-
5alkylamino wherein the nitrogen atom may be independently mono- or disubstituted by

naphthylsulfonyl or pyridinylcarbonyl, halogen, hydroxy, carboxy, oxo or cyano, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthylmethyl, piperidinyl, morpholinyl, piperazinyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, benzimidazolyl, quinolinyl, isoquinolinyl, C1-4 alkoxy, benzoyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, piperidinyl, morpholinyl, furanyl, thienyl or pyridinyl, halogen, hydroxy, oxo or cyano; and

X is O.

In yet another embodiment of the invention, there are provided novel compounds of the formula (I) as described immediately above, and wherein:

R₁ is methyl, ethyl, phenyl, piperidinyl, morpholinyl, piperazinyl, pyridinyl, pyrazinyl, furanyl, thienyl, benzyl, benzofuranyl, cyclohexyl, quinolinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is C1-3 alkyl, phenyl, piperidinyl, thienyl, C1-3 alkoxy, phenoxy, C1-3 alkanoyl, C1-3 alkoxycarbonyl, benzyloxy, C1-3 alkanoylamino, thiophenyl, benzimidazolyl, C1-3 alkylthio or chloro, R_a may be further optionally substituted by one or more R_b;

R_b is bromo, chloro, fluoro, iodo, hydroxy, oxo or cyano;

R₃ is methyl, ethyl, n-propyl, n-butyl, isobutyl, propene, butene, isobutene, C3-7 cycloalkyl or benzyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, methoxy, ethoxy, methylthio, ethylthio, cyclohexyl, phenyl, naphthyl, imidazolyl, indolinyl, cyclohexyl, bromo, chloro, fluoro, iodo, hydroxy, oxo, carboxy, nitro, benzoyl, benzyloxy, N-benzylimidazolyl or cyano, R_c may be further optionally substituted
5 by one or more R_d;

R_d is methyl, methoxy, ethoxy, chloro, fluoro, nitro or hydroxy;

R₅ is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, phenyl, methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, isobutyloxycarbonyl, tert-butoxycarbonyl and pyrimidinyl, R₅ may be
10 further optionally substituted by one or more R_e;

R_e is methyl, ethyl, n-propyl, isopropyl, phenyl, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, tert-butoxycarbonyl, bromo, chloro, fluoro, iodo, hydroxy, oxo or
15 cyano.

In yet still another embodiment of the invention, there are provided novel compounds of
20 the formula (I) as described immediately above, and wherein:

Het is piperidinyl or pyrrolidinyl;

R₁ is N-acetylamino, phenyl, chlorophenyl, methoxyphenyl, m-phenoxyphenyl, morpholinyl, pyrazinyl, pyridinyl, furanyl, chlorothieryl, thienyl or thienylmethyl;
25

R₃ is n-butyl, isobutyl, 2,2-dimethylpropyl, cyclohexylmethyl, p-methoxybenzyl or 2-naphthylmethyl; and

30 wherein the configuration at the stereocenter defined by R₂ and R₃ when they are different and the carbon they are attached to is defined as L; and

R₅ is methyl, propyl, isopropyl, ethoxycarbonyl, benzyloxycarbonyl, benzyl, phenethyl, N,N-dimethylaminoacetyl or pyrimidinyl.

5

In yet a further embodiment of the invention, there are provided novel compounds of the formula (I) as described immediately above, and wherein:

Het is piperidin-4-yl or pyrrolidinyl;

10

R₁ is morpholinyl or N-acetylamino-phenyl;


R₃ is 2,2-dimethylpropyl or cyclohexylmethyl; and

15

R₅ is methyl, propyl, isopropyl, ethoxycarbonyl, benzyloxycarbonyl, benzyl, phenethyl, N,N-dimethylaminoacetyl or pyrimidinyl.

20

In a second broad generic aspect of the invention, there are provided novel compounds of the formula (II):



25

wherein:

Het is azepanyl, piperidiny, pyrrolidiny, azetidiny, oxepanyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, azocanyl, oxocanyl, 1,3-diazocanyl, 1,4-diazocanyl, 1,5-diazocanyl, 1,3-dioxocanyl, 1,4-dioxocanyl, 1,5-dioxocanyl, 1,3-oxazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,3-diazepanyl, 1,4-diazepanyl, 1,3-dioxepanyl, 1,4-dioxepanyl, 1,3-oxazepanyl, 1,4-oxazepanyl, 1,2-thiazocanyl-1,1-dioxide, 1,2,8-thiadiazocanyl-1,1-dioxide, 1,2-thiazepanyl-1,1-dioxide, 1,2,7-thiadiazepanyl-1,1-dioxide, tetrahydrothiophenyl, hexahydropyrimidiny, hexahydropyridaziny, piperaziny, 1,4,5,6-tetrahydropyrimidiny, pyrazolidiny, dihydro-oxazolyl, dihydrothiazolyl, dihydroimidazolyl, isoxazoliny, oxazolidiny, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidiny-1,1-dioxide, imidazolidiny-2,4-dione, imidazolidiny, morpholiny, dioxanyl, tetrahydropyridiny, thiomorpholiny, thiazolidiny, dihydropyranyl, dithianyl, decahydro-quinoliny, decahydro-isoquinoliny, 1,2,3,4-tetrahydro-quinoliny, indoliny, octahydro-quinoliziny, dihydro-indoliziny, octahydro-indoliziny, octahydro-indolyl, decahydroquinazoliny, decahydroquinoxaliny, 1,2,3,4-tetrahydroquinazoliny or 1,2,3,4-tetrahydroquinoxaliny;

A C6-C10 bridged bicyclo wherein one or more carbon atoms are optionally replaced by a heteroatom chosen from N, O and S;

each being optionally substituted with one or more R₅;

Y is C(O), C(S) or S(O)₂;

25

R₁ is a bond, hydrogen, C1-10 alkyl, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, C3-8 cycloalkyloxy, aryl, benzyl, tetrahydronaphthyl, indenyl, indanyl, C1-10alkylsulfonylC1-10alkyl, C3-8cycloalkylsulfonylC1-10alkyl, arylsulfonylC1-10alkyl, heterocyclyl selected from pyrrolidiny, piperidiny, morpholiny, thiomorpholiny, piperaziny, indoliny, pyranly, tetrahydropyranyl, tetrahydrothiopyranyl, thiopyranyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridiny,

30

pyrimidinyl, pyrazinyl, pyridazinyl, tetrazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzisoxazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl, tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocycloxy wherein the
5 heterocyclyl moiety is selected from those herein described in this paragraph, hydroxy or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond, C1-10 alkyl, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,
10 indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10 alkoxy, C1-10alkanoyl, C1-10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, aryloxycarbonyl,
15 aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
20 or R_a is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl,
25 oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein
30

the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b; with the proviso that R₁ and R_a simultaneously cannot be a bond;

R_b is a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more carbon atoms are optionally replaced by O, N, S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2 oxo groups, -NH₂, or one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl;

or R_b is C3-6 cycloalkyl, aryl, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, mono-C1-5alkylamino, di-C1-5alkylamino, carboxamide, amidino or guanidino;

R₂ is hydrogen or C1-3 alkyl;

R₃ is a bond, hydrogen, C1-10 alkyl, C2-10alkylene, C3-8 cycloalkyl, arylC1-5alkyl or aryl wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-10 alkyl, C3-8 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl,

bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, decahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranyl, tetrahydropyranyl, tetrahydrothiopyranyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, dihydrobenzofuranyl, octahydrobenzofuranyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, tetrahydroquinolinyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10 alkoxy, aryloxy, C1-10 alkanoyl, aroyl, C1-10 alkoxycarbonyl, aryloxycarbonyl, C1-10 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_c is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_c is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,

benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

5

R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-5alkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino;

10 R₂ and R₃ together with the carbon they are attached optionally form a nonaromatic 5-7 membered cycloalkyl or heterocyclic ring;

R₄ is hydrogen, hydroxy or C1-3 alkyl;

15 R₅ is a bond, hydrogen, carbonyl, C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, aryl, benzyl, tetrahydronaphthyl, indenyl, indanyl, C3-7cycloalkylsulfonylC1-5alkyl, arylsulfonylC1-5alkyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl,

20 thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, thiopyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridiziny, tetrazolyl, triazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl,

25 tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, benzyloxy, C1-10alkoxycarbonyl, arylC1-5alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl,

30 morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,

benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,

or R₅ is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to

5 a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, indoliny, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridiny, pyrimidinyl, pyraziny, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,
 10 or R₅ is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, indoliny, furanyl, thienyl, pyrrolyl, oxazolyl,
 15 thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridiny, pyrimidinyl, pyraziny, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,

or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R₅ may be further optionally substituted by one or more R_e;

20

R_e is C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-10 alkoxy, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, indanyl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl,
 25 indoliny, thiopyranyl, tetrahydrothiopyranyl, pyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridiny, pyrimidinyl, pyraziny, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinoliny, isoquinoliny, quinazoliny, quinoxaliny, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, arylC1-3alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen

30

atom may be independently mono or di-substituted by C1-10 alkyl, aryl,
pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl,
furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl,
pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,
5 benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or
quinoxalinyl,

or R_c is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom
may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom
10 may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl,
morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl,
oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl,
pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl,
quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

15 or R_c is C1-10 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-10
alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino,
arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein
the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl,
aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,
20 indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl,
tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,
benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or
quinoxalinyl,

or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or
25 guanidino, R_c may be further optionally substituted by one or more R_f;

R_f is C1-5 alkyl, C3-6 cycloalkyl, tolylsulfonyl, C1-5 alkoxy, aryl,
aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro,
carboxamide, amidino or guanidino;

30 X is O or S and

pharmaceutically acceptable derivatives thereof.

- 5 In another embodiment of the invention, there are provided novel compounds of the formula (II) as described immediately above, and wherein:

Het is piperidinyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, azetidiny, azepanyl, oxepanyl, tetrahydrofuranyl, oxetanyl, hexahydropyrimidinyl,
10 hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, octahydro-indoliziny, octahydro-quinoliziny, decahydro-quinoliny, 1,2,3,4-tetrahydro-quinoliny, dihydro-oxazolyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide, imidazolidinyl, pyrazolidinyl or a bridged bicyclo chosen from aza-bicyclo[3.2.1]octane, aza-bicyclo[2.2.1]heptane, aza-bicyclo[2.2.2]octane, aza-bicyclo[3.2.2]nonane, aza-bicyclo[2.1.1]hexane, aza-bicyclo[3.1.1]heptane, aza-bicyclo[3.3.2]decane and 2-oxa or 2-thia-5-aza-bicyclo[2.2.1]heptane;
15 each ring being substituted with one or more R₅;

Y is C(O) or S(O)₂;

20

R₁ is a bond, hydrogen, C1-7 alkyl, C1-7 alkoxy, C3-7 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, tetrahydronaphthyl, C1-7alkylsulfonylC1-7alkyl, C3-7cycloalkylsulfonylC1-7alkyl, arylsulfonylC1-7alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indoliny, pyranyl, thiopyranyl, furanyl,
25 thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, isoxazolyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinoliny, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoisoxazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

30

R_a is a bond C1-7 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl,

thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl,
indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl,
quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-7 alkoxy, C1-7alkanoyl,
C1-7alkanoyloxy, aryloxy, benzyloxy, C1-7 alkoxycarbonyl, aryloxycarbonyl,
5 aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or
di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl,
thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl,
triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl,
benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl,
10 quinazolinyl or quinoxalinyl,
or R_a is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom
may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may
be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl,
15 morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl,
imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl,
isoquinolinyl, quinazolinyl or quinoxalinyl,
or R_a is C1-7 alkoxycarbonylamino, aryloxycarbonylamino, C1-7
20 alkylcarbamoyloxy, arylcarbamoyloxy, C1-7 alkylsulfonylamino,
arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein
the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl,
aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl,
thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl,
25 pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl,
benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or
guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

5 R₂ is hydrogen or methyl or ethyl;

R₃ is a bond, hydrogen, C1-5 alkyl, C2-5alkylene, C3-7 cycloalkyl, arylC1-3alkyl or aryl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is C1-5 alkyl, C3-7 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranal, tetrahydropyranal, thienyl, pyrrolyl, oxazolyl, 15 thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5 alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or 20 di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, 25 or R_c is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, 30 oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl,

pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_c is C1-5 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-5 alkyl-carbamoyloxy, aryl-carbamoyloxy, C1-5 alkyl-sulfonylamino, aryl-sulfonylamino, C1-5 alkyl-aminosulfonyl, aryl-aminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-4 alkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, halogen, hydroxy, oxo or cyano;

R₄ is hydrogen or methyl;

R₅ is a bond, hydrogen, carbonyl, C1-8 alkyl, C1-8alkoxyC1-8alkyl, C1-8alkylaminoC1-8alkyl, C1-8alkylthioC1-8alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-8 alkoxy, aryloxy, C3-7 cycloalkyl, aryl, benzyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, thiopyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, tetrazolyl, triazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-7alkanoyl, aroyl, C1-7alkanoyloxy, benzyloxy, C1-7 alkoxy-carbonyl, arylC1-4alkoxy-carbonyl,

aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-7 alkoxycarbonylamino, aryloxycarbonylamino, C1-7 alkylcarbamoyloxy, arylcarbamoyloxy, C1-7 alkylsulfonylamino, arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-7 alkyl, C1-7alkoxyC1-7alkyl, C1-7alkylaminoC1-7alkyl, C1-7alkylthioC1-7alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-7 alkoxy, C3-7 cycloalkyl, aryl, tetrahydronaphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thiopyranyl, tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkanoyl, aroyl, C1-5alkanoyloxy, aryloxy, benzyloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl,

aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_e is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_e is C1-5 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-5 alkyl-carbamoyloxy, aryl-carbamoyloxy, C1-5 alkyl-sulfonylamino, aryl-sulfonylamino, C1-5 alkyl-aminosulfonyl, aryl-aminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_e may be further optionally substituted by one or more R_f ,

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, C1-3 alkoxy, cyclopropyl, cyclohexyl, phenyl, naphthyl, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

and

X is O.

In yet another embodiment of the invention, there are provided novel compounds of the formula (II) as described immediately above, and wherein:

5 wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, octahydro-indoliziny, octahydro-quinoliziny or aza-bicyclo[3.2.1]octanyl, each ring being optionally substituted with one
10 or more R₅;

R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, C1-3alkylsulfonylC1-3alkyl, C3-6cycloalkylsulfonylC1-3alkyl, arylsulfonylC1-3alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl,
15 piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, isoxazolyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinoliny, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

20 R_a is a bond, C1-3 alkyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl
25 wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,
or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom
30 may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be

independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
or R_a is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino,
5 arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

10 R_b is C1-3 alkyl, C3-6 cycloalkyl, aryl, C1-3 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

15 R₂ is hydrogen or methyl;

R₃ is a bond, hydrogen, C1-5 alkyl, C2-5alkylene, C4-6 cycloalkyl or arylC1-2alkyl wherein R₃ is optionally substituted by one or more R_c;

20 R_c is C1-4 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranal, tetrahydropyranal, thienyl, pyrrolyl,
25 oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-4 alkoxy, phenoxy, naphthyloxy, C1-3 alkanoyl, benzoyl, C1-3 alkoxycarbonyl, phenoxycarbonyl, C1-3 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be
30 independently mono or di-substituted by C1-5 alkyl or aryl,

or R_c is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or aryl,

5 or R_c is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl,

10 or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-3 alkyl, C3-6 cycloalkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, phenylC1-3alkoxy, benzoyl, halogen, hydroxy, oxo or cyano;

15 R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-6 alkyl, C1-6alkoxyC1-6alkyl, C1-6alkylaminoC1-6alkyl, C1-6alkylthioC1-6alkyl wherein the sulfur atom may be oxidized to a sulfoxide or
20 sulfone, C1-6 alkoxy, , phenoxy, naphthyloxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl,
25 isoquinolinyl and benzoxazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-4alkanoyloxy, benzyloxy, C1-4 alkoxycarbonyl, arylC1-2alkoxycarbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl,
30 morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R₅ is C1-4 alkanoylamino, aroylamino, C1-4 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, 5 piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-4 alkoxycarbonylamino, phenoxycarbonylamino, C1-4 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-4 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be 10 independently mono or di-substituted by C1-4 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R_c;

15 R_c is C1-4 alkyl, C1-4 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl, 20 benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-4 alkanoyl, aroyl, C1-4 alkanoyloxy, phenoxy, naphthyloxy, benzyloxy, C1-4 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, 25 furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, or benzthiazolyl, or R_c is C1-4 alkanoylamino, benzoylamino, C1-4 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl, 30 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl,

thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-4 alkoxy-carbonylamino, phenoxy-carbonylamino, C1-4

alkyl-carbamoyloxy, phenyl-carbamoyloxy, C1-4 alkyl-sulfonylamino,

phenyl-sulfonylamino, C1-4 alkyl-aminosulfonyl, phenyl-aminosulfonyl, amino

wherein the nitrogen atom may be independently mono or di-substituted by C1-3

alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl,

piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl,

pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_e may be further optionally substituted by one or more R_f;

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, methoxy, cyclopropyl, phenyl,

phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy or

carboxamide.

In yet still another embodiment of the invention, there are provided novel compounds of the formula (II) as described immediately above, and wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl,

oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R₅;

R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl,

naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl,

thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl,

indolyl, quinoliny, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl,

benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

5 R_a is a bond, C1-3 alkyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxy carbonyl, aryloxy carbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkoxy carbonylamino, aryloxy carbonylamino, C1-3 alkyl carbamoyloxy, aryl carbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

20 R_b is methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, methoxy, ethoxy, n-propoxy, i-propoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

25 R₂ is hydrogen;

 R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

30 R_c is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl,

bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, C1-3 alkoxy, phenoxy, naphthyloxy, C1-2 alkanoyl, benzoyl, C1-2 alkoxycarbonyl, phenoxycarbonyl, C1-2alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or aryl, or R_c is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or aryl, or R_c is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, arylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, or R_c is halogen, hydroxy, oxo, carboxy or cyano, R_c may be further optionally substituted by one or more R_d;

R_d is methyl, cyclopropyl, cyclohexyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, fluoro, chloro, oxo or cyano;

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-3alkanoyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be

independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-3 alkoxy-carbonylamino, phenoxy-carbonylamino, C1-3 alkyl-carbamoyloxy, phenyl-carbamoyloxy, C1-3 alkyl-sulfonylamino, phenyl-sulfonylamino, C1-3 alkyl-aminosulfonyl, phenyl-aminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R₅ may be further optionally substituted by one or more R₆;

R₆ is C1-3 alkyl, C1-3 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-3 alkanoyl, aroyl, C1-3 alkanoyloxy, phenoxy, benzyloxy, C1-3 alkoxy-carbonyl, phenoxy-carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₆ is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-3 alkoxy-carbonylamino, phenoxy-carbonylamino, C1-3
alkyl-carbamoyloxy, phenyl-carbamoyloxy, C1-3 alkyl-sulfonylamino,
phenyl-sulfonylamino, C1-3 alkyl-aminosulfonyl, phenyl-aminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-3
5 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,
oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or
benzthiazolyl,

or R_e is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R_e may be further
optionally substituted by one or more R_f;

10 and

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro,
chloro, bromo, hydroxy, oxo, carboxy or carboxamide.

15 In yet still another embodiment of the invention, there are provided novel compounds of
the formula (II) as described immediately above, and wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl or tetrahydropyranyl each ring being
20 substituted with one or more R₅;

Y is C(O);

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, ethoxy, cyclopropyl, cyclopentyl,
25 cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl,
thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl
or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond, methyl, ethyl, cyclopropyl, phenyl, pyrrolidinyl, piperidinyl,
30 morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, acetyl,
acetoxyl, phenoxy, benzyloxy, methoxycarbonyl, phenoxy-carbonyl, benzyloxy,

carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_a is acetylamino, benzoylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R_a is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

or R_a is fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_a may be further optionally substituted by one or more R_b;

R_b is methyl, cyclopropyl, phenyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, i-propyl, C5-6 cycloalkyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, thienyl, oxazolyl, thiazolyl, indolyl, benzofuranyl, benzothieryl, benzthiazolyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or aryl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or aryl,

or R_c is methoxycarbonylamino, phenoxy carbonylamino, methylcarbamoxyloxy, phenylcarbamoxyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
5 or R_c is fluoro, chloro or oxo, R_c may be further optionally substituted by one or more R_d;

R_d is methyl, cyclopropyl, phenyl, methoxy, fluoro, chloro or oxo;

10 R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-4alkoxyC1-4alkyl, C1-4alkylaminoC1-4alkyl, C1-4alkylthioC1-4alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-4 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl,
15 benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-2alkanoyl, benzoyl, naphthoyl, C1-2alkanoyloxy, benzyloxy, C1-2 alkoxy carbonyl, benzyloxy carbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl,
20 morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,
25 oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-2 alkoxy carbonylamino, phenoxy carbonylamino, C1-2 alkylcarbamoxyloxy, phenylcarbamoxyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl,
30 morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R₅ is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C1-2 alkoxy, C3-6 cycloalkyl, phenyl, naphthyl, indanyl,
5 pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl,
thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl,
benzthiazolyl, benzoxazolyl, C1-2 alkanoyl, aroyl, C1-2alkanoyloxy, phenoxy,
benzyloxy, C1-2 alkoxy carbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl
wherein the nitrogen atom may be independently mono or di-substituted by C1-2
10 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl,
thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R_e is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen
15 atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl,
piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or
pyrimidinyl,

or R_e is C1-2 alkoxy carbonylamino, phenoxy carbonylamino, C1-2
alkyl carbamoyloxy, phenyl carbamoyloxy, C1-2 alkylsulfonylamino,
20 phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-2
alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,
oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R_e is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R_e may be
25 further optionally substituted by one or more R_f;

and

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro,
chloro, hydroxy, oxo, carboxy or carboxamide.

30 In yet a further embodiment of the invention, there are provided novel compounds of the
formula (II) as described immediately above, and wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl, azepan-3-yl, azepan-4-yl or tetrahydropyran-4-yl, each ring being optionally substituted with one or more R₅;

5 R₁ is a bond, methyl, ethyl, i-propyl, methoxy, cyclopropyl, cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

10 R_a is methyl, phenyl, thienyl, methoxy, acetyl, acetoxy, phenoxy, benzyloxy, methoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
or R_a is acetylamino, methylthio, phenylthio wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
15 independently substituted by methyl or phenyl,
or R_a is methoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
or R_a is fluoro, chloro, hydroxy, oxo, carboxy, cyano or carboxamide;

20

R₃ is a bond, methyl, ethyl, n-propyl, propenyl, butenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

25

R_c is methyl, ethyl, n-propyl, i-propyl, cyclohexyl, cyclopentyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be
30 oxidized to a sulfoxide or sulfone, fluoro, chloro or oxo;

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-2alkoxyC1-2alkyl, C1-2alkylaminoC1-2alkyl, C1-2alkylthioC1-2alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-2 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_c;

R_c is methyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidinyl, morpholinyl, indolyl, thienyl, pyridinyl, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R_c is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino,

phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f;

and

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo.

10

In yet still a further embodiment of the invention, there are provided novel compounds of the formula (II) as described immediately above, and wherein:

15

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl or tetrahydropyran-4-yl, each ring being substituted with one or more R₅;

20

R₁ is i-propyl, benzyloxy, cyclohexyl, phenyl, 4-(acetylamino)-phenyl, 4-(methanesulfonylamino)-phenyl, 4-methoxyphenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-chlorophenyl, naphthyl, thienylmethyl, piperidinyl, morpholinyl, pyrrolidinyl, piperazinyl, furanyl, thienyl, 5-chlorothieryl, pyridin-4-yl, pyrazinyl, methylamino, ethylamino, dimethylamino or diethylamino;

25

R₃ is ethyl, n-propyl, propenyl, butenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

30

R_c is methyl, cyclohexyl, cyclopentyl, indanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro or chloro;

R₅ is a bond, carbonyl, methyl, ethyl, n-propyl, n-butyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, piperidinyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy, R₅ may be
5 further optionally substituted by one or more R_e;

R_e is methyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, thienyl, 5-methylthienyl, methoxy, phenoxy, benzyloxy, piperidinyl, pyridinyl, indolyl, 1-(tolyl-sulfonyl)-indolyl, carbamoyl wherein the nitrogen atom may be
10 independently mono or di-substituted by methyl, phenyl or benzyl, or R_e is hydroxy, fluoro, chloro, oxo, dimethylamino or trifluoromethyl;

15 In yet another embodiment of the invention, there are provided novel compounds of the formula (II) as described immediately above, and wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl or azetidin-3-yl, each ring being substituted with one or more R₅;

20 R₁ is phenyl, 4-(acetylamino)-phenyl, 4-(methanesulfonylamino)-phenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-fluorophenyl, thienylmethyl, morpholinyl, pyrrolidinyl, piperidinyl, piperazinyl, 5-chlorothienyl, pyridin-4-yl or pyrazinyl;

25 R₃ is n-butyl, i-butyl, 2,2-dimethylpropyl, cyclohexylmethyl, propenyl, i-butenyl, 4-methoxybenzyl, 4-chlorobenzyl, 3,4-dichlorobenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylbenzyl, 3-methylbenzyl or naphth-2-ylmethyl; wherein the configuration at the stereocenter defined by R₂ and R₃ when they are different and the carbon they are attached to is defined as L;

30 and

R₅ is a bond, methyl, ethyl, n-propyl, n-butyl, n-pentyl, 2-pentyl, 3-pentyl, phenethyl, phenpropyl, 2,2-dimethylpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, phenyl, benzyl, naphthylmethyl, indanylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, 5-methylthienylmethyl, piperidinyl, piperidinylcarbonyl, pyridinylcarbonyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, methylcarbamoyl, phenylcarbamoyl, benzylcarbamoyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, methylcyclohexyl, methylbenzyl, methoxybenzyl, phenoxybenzyl, benzyloxybenzyl, N-[(4-methylphenyl)-sulfonyl]-indolylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, N,N-dimethylaminoacetyl, trifluoromethylbenzyl, fluoro, oxo or carboxy.

Another embodiment of the invention provides for the following compounds of the formulas (I) and (II) above which have demonstrated potent inhibition of Cathepsin S in a cell based assay at concentrations of 15 uM or less.

Morpholine-4-carboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

4-Acetylamino-N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

4-Acetylamino-N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid *t*-butyl ester;

4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester;

- Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 5 Morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride;
- Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;
- 10 Morpholine-4-carboxylic acid [1-(4-cyano-1-phenethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- Morpholine-4-carboxylic acid {1-[3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;
- 15 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid benzyl ester;
- 20 Morpholine-4-carboxylic acid [1-(4-cyano-1-isopropyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- 25 Morpholine-4-carboxylic acid [1-(1-phenethyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- Morpholine-4-carboxylic acid [1-(1-n-propyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- 30 Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- Morpholine-4-carboxylic acid [1-(4-cyano-tetrahydro-thiopyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 35 Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-dimethylamino-acetyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;
- 40 4-Cyano-4-{3-cyclohexyl-2-[(4-acetylamino)-phenyl-1-carbonyl]-amino}-propionylamino}-piperidine-1-carboxylic acid ethyl ester;
- Morpholine-4-carboxylic acid [1-(3-cyano-1-pyrimidin-2-yl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 45

- 4-Cyano-4-{4,4-dimethyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid benzyl ester;
- 5 4-Acetylamino-*N*-[1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;
- 4-Acetylamino-*N*-[1-(4-cyano-1-isopropyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;
- 10 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 4-Cyano-4-{3-cyclohexyl-2-[(4-acetylamino)-phenyl-1-carbonyl]-amino}-propionylamino}-piperidine-1-carboxylic acid benzyl ester;
- 15 *N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;
- Morpholine-4-carboxylic acid [1-(1-carbamimidoyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide p-toluenesulfonate;
- 20 4-Acetylamino-*N*-[1-(4-cyano-1-phenethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;
- 4-(Acetylamino-methyl)-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;
- 25 4-Cyano-4-{4,4-dimethyl-2-[(morpholine-4-carbonyl)-amino]-pentanoylamino}-piperidine-1-carboxylic acid ethyl ester;
- 30 Morpholine-4-carboxylic acid [1-(1-acetyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- Morpholine-4-carboxylic acid [1-(1-benzoyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- 35 3-Cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-pyrrolidine-1-carboxylic acid benzyl ester;
- N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-isonicotinamide;
- 40 Pyrazine-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- Morpholine-4-carboxylic acid [1-(1-acetyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 45

Morpholine-4-carboxylic acid [1-(1-benzoyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

5 Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-chloro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

5-Chloro-thiophene-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

10 4-Chloro-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-phenylcarbamoyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

15 Morpholine-4-carboxylic acid [1-(1-benzylcarbamoyl 4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

20 *N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-methanesulfonylamino-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-3-phenoxy-benzamide;

25 *N*-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl-ethyl]-isonicotinamide;

Pyrazine-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

30 *N*-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-3-cyclohexyl-2-(2-thiophen-2-yl-acetylamino)-propionamide;

35 5-Chloro-thiophene-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(cyclohexyl-methyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

40 *N*-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-3-phenoxy-benzamide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

45

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-chloro-benzamide;

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

5

Pyrazine-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-methyl-ethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

10

N-[1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide;

15

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-benzyloxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

20

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-methanesulfonylamino-benzamide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-benzyloxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

25

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3,5-difluoro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

30

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2,6-difluoro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

35

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-2-fluoro-benzamide;

40

4-Chloro-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-2-fluoro-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-methoxy-benzamide;

45

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-fluoro-benzamide;

5 *N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-phenoxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

10 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-methyl-piperidine-4-yl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

15 Morpholine-4-carboxylic acid [1-(3-cyano-1-ethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

20 Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-methyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

25 Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-phenoxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-methyl-pent-2-enyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

30 Morpholine-4-carboxylic acid {1-[3-cyano-1-(4-fluoro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

35 Morpholine-4-carboxylic acid {1-[3-cyano-1-(2,4,6-trimethyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1*H*-indol-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

40 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopropyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

45 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-pyridin-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Pyrrolidine-1-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

5 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-1-oxy-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Isoxazole-5-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

10 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

15 1*H*-Imidazole-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

20 Morpholine-4-carboxylic acid {1-[3-cyano-1-(5,5-dimethyl-3-oxo-cyclohex-1-enyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-isopropyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

25 Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

30 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;

35 Morpholine-4-carboxylic acid [1-(3-cyano-1-phenethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopropylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

40 Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-azetidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

45

4-Cyano-4-{3-cyclohexyl-2-[(4-methyl-piperazine-1-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester;

5 Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

10 Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(*cis*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

15 *N*-[4-Cyano-tetrahydro-pyran-4-ylcarbamoyl]-2-cyclohexyl-ethyl]-isonicotinamide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

20 Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

25 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(*cis*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;

30 Morpholine-4-carboxylic acid [1-(3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;

35 Morpholine-4-carboxylic acid (1-{3-cyano-1-[1-(toluene-4-sulfonyl)-1*H*-indol-3-ylmethyl]-pyrrolidin-3-ylcarbamoyl}-2-cyclohexyl-ethyl)-amide;

40 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-(4-iodo-phenyl)-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-*p*-tolyl-ethyl]-amide;

45 Morpholine-4-carboxylic acid [(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-cyclohexyl-methyl]-amide;

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-naphthalen-2-yl-ethyl]-amide;

- 5 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-(4-chloro-phenyl)-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

10

Morpholine-4-carboxylic acid {1-[3-cyano-1-(indan-2-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

15

Morpholine-4-carboxylic acid {1-[3-cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

1-Benzyl-3-cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-pyrrolidine-2-carboxylic acid methyl ester;

20

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-isobutyramide;

[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-carbamic acid benzyl ester;

25

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-but-3-enyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-methoxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

30

Morpholine-4-carboxylic acid {1-[3-cyano-1-(naphthalen-2-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

35

Cyclohexanecarboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

40

Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-piperidine-4-carbonyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[4-cyano-1-(pyridine-4-carbonyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

45

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-2-hydroxymethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

5 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

4-Chloro-*N*-[1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

10 Pyrazine-2-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

4,4-dimethyl-2-(2-thiophen-2-yl-acetylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide;

15 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

20 *N*-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(morpholine-4-carbothioyl)-amino]-propionamide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-cyclohexyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

25 Morpholine-4-carboxylic acid {1-[4-cyano-1-(tetrahydro-pyran-4-yl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;.

Morpholine-4-carboxylic acid [2-(4-chloro-phenyl)-1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-ethyl]-amide;

30 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-(3,4-dichloro-phenyl)-ethyl]-amide;

35 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-naphthalen-2-yl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-amide;

40 Morpholine-4-carboxylic acid [1-(4-cyano-1,2-dimethyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Naphthalene-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

45

and the pharmaceutically acceptable derivatives thereof.

The following are preferred compounds of the formulas (I) and (II) of the invention:

- 5 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 10 4-Acetylamino-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;
- 4-Acetylamino-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;
- 15 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 20 Morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride;
- Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;
- 25 Morpholine-4-carboxylic acid [1-(4-cyano-1-phenethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 30 Morpholine-4-carboxylic acid {1-[3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;
- Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;
- 35 Morpholine-4-carboxylic acid [1-(4-cyano-1-isopropyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- Morpholine-4-carboxylic acid [1-(1-phenethyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- 40 Morpholine-4-carboxylic acid [1-(1-*n*-propyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;
- 45 Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

4-Acetylamino-*N*-[1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

5 4-Acetylamino-*N*-[1-(4-cyano-1-isopropyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

10 *N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

4-(Acetylamino-methyl)-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

15 *N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-isonicotinamide;

Pyrazine-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

20 5-Chloro-thiophene-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

25 4-Chloro-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-3-phenoxy-benzamide;

30 *N*-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl-ethyl]-isonicotinamide;

Pyrazine-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

35 Morpholine-4-carboxylic acid {1-[3-cyano-1-(cyclohexyl-methyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

40 Morpholine-4-carboxylic acid [1-(3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

45 Pyrazine-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-methyl-ethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

5 *N*-[1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide;

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-methanesulfonylamino-benzamide;

10 *N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-fluoro-benzamide;

15 *N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

20 Morpholine-4-carboxylic acid [1-(3-cyano-1-ethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

25 Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-methyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

30 Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-methyl-pent-2-enyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1*H*-indol-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

35 Pyrrolidine-1-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

40 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

45 Morpholine-4-carboxylic acid [1-(3-cyano-1-isopropyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

- 5 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;

- 10 Morpholine-4-carboxylic acid [1-(3-cyano-1-phenethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

- 15 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopropylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

- 20 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-azetidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

- 25 Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

- 30 Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

- 35 Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

- 40 Morpholine-4-carboxylic acid [1-(3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

- 45 Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-naphthalen-2-yl-ethyl]-amide;

5 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-(4-chloro-phenyl)-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

10 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-but-3-enyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

15 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

20 4-Chloro-*N*-[1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

Pyrazine-2-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

25 4,4-dimethyl-2-(2-thiophen-2-yl-acetylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide;

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

30 Morpholine-4-carboxylic acid [1-(4-cyano-1-cyclohexyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

35 Morpholine-4-carboxylic acid [2-(4-chloro-phenyl)-1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-(3,4-dichloro-phenyl)-ethyl]-amide;

40 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-naphthalen-2-yl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-amide;

45

Morpholine-4-carboxylic acid [1-(4-cyano-1,2-dimethyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

5 and the pharmaceutically acceptable derivatives thereof.

The activity of particular compounds disclosed herein against cathepsin K may be
10 determined without undue experimentation by one of ordinary skill in the art in view of the art, the guidance provided throughout this specification and by the screens described in the section entitled "Assessment of Biological Properties."

The following subgeneric aspect of the compounds of the formula (II) have Cathepsin K
15 activity:

The compound according to the third embodiment above of formula (II)
and wherein:

20 Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R_5 ;

R_1 is a bond, C1-4 alkyl, C1-4 alkoxy, cyclopropyl, cyclohexyl, phenoxy, naphthyloxy,
25 phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinoliny, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R_1 is optionally substituted by one or more R_a ;

30 R_a is methyl, ethyl, propyl, i-propyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, ethoxy, acetyl, acetoxyl, phenoxy, naphthyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, phenoxycarbonyl, naphthyloxycarbonyl,

benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ethylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, naphthylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, naphthylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, naphthylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is methyl, ethyl, cyclopropyl, cyclohexyl, phenyl, methoxy, ethoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₂ is hydrogen or methyl;

R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, naphthyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, furanyl,

- tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
- 5 or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,
- 10 or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
- or R_c is chloro, fluoro, hydroxy, oxo, carboxy or cyano;
- 15 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl or tetrahydrothiophenyl;
- 20 R₄ is hydrogen;
- R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, naphthyloxy, cyclopropyl, cyclopentyl, cyclohexyl,
- 25 phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-
- 30 substituted by methyl, ethyl or phenyl,

or R₅ is acetylamino, benzoylamino, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino,
5 methylcarbamoyloxy, phenylcarbamoyloxy, phenylsulfonylamino, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

10

R_e is methyl ethyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidinyl, morpholinyl, indolyl, thienyl, pyridinyl, methoxy, ethoxy, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be
15 independently mono or di-substituted by methyl, ethyl or phenyl,

or R_e is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

20

or R_e is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

25

or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f;

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or
30 oxo.

30

Another embodiment of the compounds of the formula (II) having Cathepsin K activity are those described immediately above and wherein:

R₁ is a bond, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, benzyloxy, cyclopropyl, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is methyl, cyclopropyl, phenyl, halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

or R_c is acetylamino, benzoylamino, methylthio, methoxycarbonylamino, methylcarbamoyloxy, methylsulfonylamino, methylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl,

or R_c is fluoro or oxo;

R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, pyrrolidinyl or piperidinyl;

R₅ is methyl, ethyl, n-propyl, n-butyl, n-pentyl, 2-pentyl, 3-pentyl, phenethyl, phenpropyl, 2,2-dimethylpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methylbenzyl, 3-methylbenzyl, 4-methylbenzyl, 2,6-dimethylbenzyl, 2,5-dimethylbenzyl, 2,4-dimethylbenzyl, 2,3-dimethylbenzyl, 3,4-dimethylbenzyl, 3,5-dimethylbenzyl, 2,4,6-trimethylbenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-phenoxybenzyl, 3-phenoxybenzyl, 4-phenoxybenzyl, 2-benzyloxybenzyl, 3-benzyloxybenzyl, 4-benzyloxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 2,6-difluorobenzyl, 2,5-difluorobenzyl, 2,4-difluorobenzyl, 2,3-difluorobenzyl, 3,4-difluorobenzyl, 3,5-difluorobenzyl, 2,4,6-trifluorobenzyl, 2-trifluoromethylbenzyl, 3-trifluoromethylbenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, indanylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, 5-methylthienylmethyl, piperidinyl, piperidinylcarbonyl, pyridinylcarbonyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, methylcarbamoyl, phenylcarbamoyl, benzylcarbamoyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy.

Yet another embodiment of the compounds of the formula (II) having Cathepsin K activity are those described immediately above and wherein:

R₁ is methoxy, benzyloxy, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is methyl, phenyl, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl or benzyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, acetilamino, methylthio, methylsulfonylamino or fluoro;

- 5 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl or tetrahydrofuranlyl;

- R₅ is methyl, ethyl, n-propyl, n-butyl, phenethyl, phenpropyl, t-butyl, i-propyl, i-butyl, 10 cyclopropyl, cyclohexyl, cyclopropylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl 4-fluorobenzyl, 3,5-difluorobenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, phenylcarbamoyl, phenylsulfonylamino or fluoro.

15

Yet still another embodiment of the compounds of the formula (II) having Cathepsin K activity are those described immediately above and wherein:

Het is pyrrolidinyl, piperidinyl or tetrahydropyranyl;

20

R₁ is benzyloxy, phenoxy, naphthyloxy, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyridinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or phenylamino;

- 25 R₃ is n-propyl, i-butyl, propenyl, i-butenyl or 2,2-dimethylpropyl;

R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, or cycloheptyl;

R₅ is methyl, ethyl, n-propyl, phenethyl, t-butyl, i-propyl, i-butyl, cyclohexyl, cyclohexylmethyl, benzyl, 4-fluorobenzyl, naphthylmethyl, acetyl, benzoyl or benzyloxycarbonyl.

5 Representative compounds possessing CAT K activity are the following:

[1-(1-Benzyl-4-cyano-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;

10 [1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-cyclohexyl]-carbamic acid *t*-butyl ester;

[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;

15 [1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-cyclohexyl]-carbamic acid benzyl ester;

Naphthalene-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

20 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-amide;

25 Naphthalene-2-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;


30 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

[1-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;

35 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

40 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-but-3-enyl]-amide.

In a third broad generic aspect of the invention, there are provided novel compounds of the formulas (Ia) and (Ib):



wherein:

- 10 Het is
 azepanyl, piperidinyl, pyrrolidinyl, azetidiny, oxepanyl, tetrahydropyranyl,
 tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, azocanyl, oxocanyl, 1,3-diazocanyl,
 1,4-diazocanyl, 1,5-diazocanyl, 1,3-dioxocanyl, 1,4-dioxocanyl, 1,5-dioxocanyl, 1,3-
 oxazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,3-diazepanyl, 1,4-diazepanyl, 1,3-
 15 dioxepanyl, 1,4-dioxepanyl, 1,3-oxazepanyl, 1,4-oxazepanyl, 1,2-thiazocanyl-1,1-
 dioxide, 1,2,8-thiadiazocanyl-1,1-dioxide, 1,2-thiazepanyl-1,1-dioxide, 1,2,7-
 thiadiazepanyl-1,1-dioxide, tetrahydrothiophenyl, hexahydropyrimidinyl,
 hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, pyrazolidinyl, dihydro-
 oxazolyl, dihydrothiazolyl, dihydroimidazolyl, isoxazolinyl, oxazolidinyl, 1,2-
 20 thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide,
 imidazolidinyl-2,4-dione, imidazolidinyl, morpholinyl, dioxanyl, tetrahydropyridinyl,
 thiomorpholinyl, thiazolidinyl, dihydropyranyl, dithianyl, decahydro-quinolinyl,
 decahydro-isoquinolinyl, 1,2,3,4-tetrahydro-quinolinyl, indolinyl, octahydro-quinoliziny,
 dihydro-indoliziny, octahydro-indoliziny, octahydro-indolyl, decahydroquinazolinyl,
 25 decahydroquinoxaliny, 1,2,3,4-tetrahydroquinazolinyl or 1,2,3,4-tetrahydroquinoxaliny;

A C6-C10 bridged bicyclo wherein one or more carbon atoms are optionally replaced by a heteroatom chosen from N, O and S;

each being optionally substituted with one or more R₅;

- 5 R₁ is a bond, hydrogen, C1-10 alkyl, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, C3-8 cycloalkyloxy, aryl, benzyl, tetrahydronaphthyl, indenyl, indanyl, C1-10alkylsulfonylC1-10alkyl, C3-8cycloalkylsulfonylC1-10alkyl, arylsulfonylC1-10alkyl, heterocyclyl selected from azepanyl, azocanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, tetrahydrothiopyranyl, thiopyranyl, furanyl, tetrahydrofuranlyl, thienyl, pyrrolyl, oxazolyl, 10 isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, tetrazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzisoxazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl, tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy 15 wherein the heterocyclyl moiety is selected from those herein described in this paragraph, hydroxy or amino; wherein R₁ is optionally substituted by one or more R_a;

- R_a is a bond, C1-10 alkyl, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, 20 indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10 alkoxy, C1-10alkanoyl, C1-10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or 25 di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, 30 isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

with the proviso that R₁ and R_a simultaneously cannot be a bond;

R_b is a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more carbon atoms are optionally replaced by O, N, S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2 oxo groups, -NH₂, or one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl;

or R_b is C3-6 cycloalkyl, aryl, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, mono-C1-5alkylamino, di-C1-5alkylamino, carboxamide, amidino or guanidino;

5 R₂ is hydrogen or C1-3 alkyl;

R₃ is a bond, hydrogen, C1-10 alkyl, C2-10alkylene, C3-8 cycloalkyl, arylC1-5alkyl or aryl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is C1-10 alkyl, C3-8 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, decahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranal, tetrahydropyranal, tetrahydrothiopyranal, 15 thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, dihydrobenzofuranyl, octahydrobenzofuranyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, tetrahydroquinolinyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10 alkoxy, aryloxy, C1-10 alkanoyl, 20 aroyl, C1-10 alkoxy carbonyl, aryloxy carbonyl, C1-10 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, 25 indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_c is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom 30 may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl,

oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
 or R_c is C1-10 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
 or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-5alkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino;

R₂ and R₃ together with the carbon they are attached optionally form a nonaromatic 5-7 membered cycloalkyl or heterocyclic ring;

each R₄ is independently hydrogen, hydroxy or C1-3 alkyl;

R₅ is a bond, hydrogen, carbonyl, C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, aryl, benzyl, tetrahydronaphthyl, indenyl, indanyl, C3-7cycloalkylsulfonylC1-5alkyl, arylsulfonylC1-5alkyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranal, tetrahydropyranal, thiopyranal, tetrahydrothiopyranal, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridiziny, tetrazolyl, triazolyl,

pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl, tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocycloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, benzyloxy, C1-10alkoxycarbonyl, arylC1-5alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R₅ may be further optionally substituted by one or more R₆;

R₆ is C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be oxidized to a sulfoxide or

sulfone, C1-10 alkoxy, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, thiopyranyl, tetrahydrothiopyranyl, pyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, arylC1-3alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_e is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_e is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,

benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_e may be further optionally substituted by one or more R_f ;

5

R_f is C1-5 alkyl, C3-6 cycloalkyl, tolylsulfonyl, C1-5 alkoxy, aryl, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

10

R_6 is

hydrogen, hydroxy, nitrile or

a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH,

15 S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2 oxo groups, -NH₂, one or more C₁₋₄ alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl,
20 quinazolinyl, benzoxazolyl or quinoxalinyl;

wherein R_1 and R_6 in the formulas (Ia) or (Ib) optionally form a 4 to 8 membered mono- or 7-12 membered polycyclo heteroring system, each aromatic or nonaromatic, wherein each heteroring is optionally substituted by one or more R_7 ;

25

each R_7 and R_8 are independently:

C1-5 alkyl chain optionally interrupted by one or two N, O or S(O)_m and optionally
30 substituted by 1-2 oxo, amino, hydroxy, halogen, C1-4alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl,

thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl or quinoxaliny,

5 aryl, aryloxy, aroyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, C1-5 alkanoyl, C1-5 alkoxy carbonyl, aryloxy carbonyl, benzyloxy carbonyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, C3-6 cycloalkyl and benzyloxy

10 each of the aforementioned are optionally halogenated, halogen, hydroxy, oxo, carboxy, nitrile, nitro or $\text{NH}_2\text{C}(\text{O})-$;

m is 0, 1 or 2;

15 X is $=\text{O}$, $=\text{S}$ or $=\text{N}-\text{R}_6$ wherein R_6 is as defined above, and

pharmaceutically acceptable derivatives thereof.

20 In another embodiment of the invention, there are provided novel compounds of the formula (Ia) and formula (Ib) as described immediately above, and wherein:

Het is piperidinyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, azetidiny,

azepanyl, oxepanyl, tetrahydrofuranyl, oxetanyl, hexahydropyrimidinyl,

25 hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, octahydro-indoliziny,

octahydro-quinoliziny, decahydro-quinoliny, 1,2,3,4-tetrahydro-quinoliny, dihydro-oxazolyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide, imidazolidinyl, pyrazolidinyl or a bridged bicyclo chosen from aza-bicyclo[3.2.1]octane, aza-bicyclo[2.2.1]heptane, aza-bicyclo[2.2.2]octane, aza-bicyclo[3.2.2]nonane, aza-bicyclo[2.1.1]hexane, aza-bicyclo[3.1.1]heptane, aza-bicyclo[3.3.2]decane and 2-oxa or 2-thia-5-aza-bicyclo[2.2.1]heptane;

30 each ring being substituted with one or more R_5 ;

R₁ is a bond, hydrogen, C1-7 alkyl, C1-7 alkoxy, C3-7 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, tetrahydronaphthyl, C1-7alkylsulfonylC1-7alkyl, C3-7cycloalkylsulfonylC1-7alkyl, arylsulfonylC1-7alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranal, thiopyranal, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, isoxazolyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoisoxazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond C1-7 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, C1-7 alkoxy, C1-7alkanoyl, C1-7alkanoyloxy, aryloxy, benzyloxy, C1-7 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalyl,

or R_a is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalyl,

or R_a is C1-7 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-7 alkyl-carbamoyloxy, aryl-carbamoyloxy, C1-7 alkylsulfonylamino, arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

R₂ is hydrogen or methyl or ethyl;

R₃ is a bond, hydrogen, C1-5 alkyl, C2-5alkylene, C3-7 cycloalkyl, arylC1-3alkyl or aryl wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-5 alkyl, C3-7 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranyl, tetrahydropyranyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5 alkanoyl, aroyl, C1-5 alkoxy-carbonyl, aryloxy-carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl,

thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyll or quinoxalinyll,

5 or R_c is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, 10 oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyll or quinoxalinyll,

or R_c is C1-5 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, 15 arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, 20 benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyll or quinoxalinyll,

or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

25 R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-4 alkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, halogen, hydroxy, oxo or cyano;

R_d is hydrogen or methyl;

R₅ is a bond, hydrogen, carbonyl, C1-8 alkyl, C1-8alkoxyC1-8alkyl, C1-8alkylaminoC1-8alkyl, C1-8alkylthioC1-8alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-8 alkoxy, aryloxy, C3-7 cycloalkyl, aryl, benzyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, thiopyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, tetrazolyl, triazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-7alkanoyl, aroyl, C1-7alkanoyloxy, benzyloxy, C1-7 alkoxycarbonyl, arylC1-4alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-7 alkoxycarbonylamino, aryloxycarbonylamino, C1-7 alkylcarbamoyloxy, arylcarbamoyloxy, C1-7 alkylsulfonylamino, arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R_e;

5 R_e is C1-7 alkyl, C1-7alkoxyC1-7alkyl, C1-7alkylaminoC1-7alkyl, C1-7alkylthioC1-7alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-7 alkoxy, C3-7 cycloalkyl, aryl, tetrahydronaphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thiopyranyl, tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl, 10 oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxaliny, C1-5 alkanoyl, aroyl, C1-5alkanoyloxy, aryloxy, benzyloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or 15 di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny,

20 or R_e is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, 25 imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny,

30 or R_e is C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl,

aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny,

5 or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_e may be further optionally substituted by one or more R_f;

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, C1-3 alkoxy, cyclopropyl, cyclohexyl, phenyl, naphthyl, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy, cyano,
10 nitro or carboxamide;

R_g is

hydrogen, hydroxy, nitrile or

a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially
15 or fully halogenated wherein one or more C atoms are optionally replaced by O, NH, S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2 oxo groups, -NH₂, one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
20 benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl or quinoxaliny;

R₁ and R₆ of the formula (Ia) or formula (Ib) form a monocyclic 5, 6 or 7 membered aromatic or nonaromatic heterocyclic ring optionally substituted by R₇;

25 or a bicyclic ring having one 5, 6 or 7 membered aromatic or nonaromatic heterocyclic ring fused to a second 5-7 membered aromatic or nonaromatic heterocyclic or carbocyclic ring wherein each ring is optionally independently substituted by one or more R₇;

R₇ and R₈ are independently C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, benzyloxy each of the aforementioned are optionally halogenated or R_x is halogen, hydroxy, oxo, carboxy, nitrile, nitro or NH₂C(O)-;

5 m is 0, 1 or 2 and

X is O or S.

10 In yet another embodiment of the invention, there are provided novel compounds of the formulas (Ia) and (Ib) as described immediately above, and wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, octahydro-indoliziny, octahydro-quinoliziny or aza-bicyclo[3.2.1]octanyl, each ring being optionally substituted with one
15 or more R₅;

R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, C1-3alkylsulfonylC1-3alkyl, C3-6cycloalkylsulfonylC1-3alkyl,
20 arylsulfonylC1-3alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, isoxazolyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinoliny, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

25

R_a is a bond, C1-3 alkyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy,
30 aryloxy, benzyloxy, C1-3 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3

alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is C1-3 alkyl, C3-6 cycloalkyl, aryl, C1-3 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

R₂ is hydrogen or methyl;

R₃ is a bond, hydrogen, C1-5 alkyl, C2-5alkylene, C4-6 cycloalkyl or arylC1-2alkyl wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-4 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranyl, tetrahydropyranyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl,

indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-4 alkoxy, phenoxy, naphthyloxy, C1-3 alkanoyl, benzoyl, C1-3 alkoxy carbonyl, phenoxycarbonyl, C1-3 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be

independently mono or di-substituted by C1-5 alkyl or aryl,

or R_c is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or aryl,

or R_c is C1-3 alkoxy carbonylamino, aryloxy carbonylamino, C1-3 alkyl carbamoyloxy, aryl carbamoyloxy, C1-3 alkyl sulfonylamino, aryl sulfonylamino, C1-3 alkyl aminosulfonyl, aryl aminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl,

or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-3 alkyl, C3-6 cycloalkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, phenyl C1-3 alkoxy, benzoyl, halogen, hydroxy, oxo or cyano;

R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-6 alkyl, C1-6 alkoxy C1-6 alkyl, C1-6 alkyl amino C1-6 alkyl, C1-6 alkyl thio C1-6 alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-6 alkoxy, , phenoxy, naphthyloxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl and benzoxazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3 alkanoyl, benzoyl, naphthoyl,

C1-4alkanoyloxy, benzyloxy, C1-4 alkoxy carbonyl, arylC1-2alkoxy carbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
5 or R₅ is C1-4 alkanoylamino, aroylamino, C1-4 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
10 benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl,
or R₅ is C1-4 alkoxy carbonylamino, phenoxy carbonylamino, C1-4 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-4 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-4 alkyl, aryl, pyrrolidinyl, piperidinyl,
15 morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,
or R₅ is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R_e;

20 R_e is C1-4 alkyl, C1-4 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl,
25 C1-4 alkanoyl, aroyl, C1-4alkanoyloxy, phenoxy, naphthyloxy, benzyloxy, C1-4 alkoxy carbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl,
30 benzimidazolyl, or benzthiazolyl,

or R_e is C1-4 alkanoylamino, benzoylamino, C1-4 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl,
5 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-4 alkoxy-carbonylamino, phenoxycarbonylamino, C1-4 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-4 alkylsulfonylamino,
10 phenylsulfonylamino, C1-4 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

15 or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_e may be further optionally substituted by one or more R_f,

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, methoxy, cyclopropyl, phenyl, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy or
20 carboxamide.

R₁ and R₆ of the formula (Ia) or Formula (Ib) optionally form a monocyclic 5 or 6 membered aromatic or nonaromatic heterocyclic ring optionally substituted by R₇;

25 or a bicyclic ring having one 5, 6 or 7 membered aromatic or nonaromatic heterocyclic ring fused to a second 5-6 membered aromatic or nonaromatic heterocyclic or carbocyclic ring wherein each ring is optionally independently substituted by one or more R₇;

30 R₇ and R₈ are independently C1-4 alkyl, C5-6 cycloalkyl, C1-4 alkoxy, halogen, hydroxy, oxo, carboxy, nitrile, nitro or NH₂C(O)-;

and

X is O.

- 5 In yet still another embodiment of the invention, there are provided novel compounds of the formulas (Ia) and (Ib) as described immediately above, and wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more

10 R₅;

R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond, C1-3 alkyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxy carbonyl, aryloxy carbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkoxy carbonylamino, aryloxy carbonylamino, C1-3 alkyl carbamoyloxy, aryl carbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein

the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

5

R_b is methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, methoxy, ethoxy, n-propoxy, i-propoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

10

R₂ is hydrogen;

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

15

R_c is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, C1-3 alkoxy, phenoxy, naphthyloxy, C1-2 alkanoyl, benzoyl, C1-2 alkoxycarbonyl, phenoxycarbonyl, C1-2alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or aryl, or R_c is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or aryl, or R_c is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, arylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2alkylaminosulfonyl, phenylaminosulfonyl, amino

20

25

30

wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl,

or R_c is halogen, hydroxy, oxo, carboxy or cyano, R_c may be further optionally substituted by one or more R_d;

5

R_d is methyl, cyclopropyl, cyclohexyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, fluoro, chloro, oxo or cyano;

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is
10 selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-3alkanoyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
15 or R₅ is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzofuranyl, benzothienyl,
20 benzimidazolyl or benzthiazolyl,
or R₅ is C1-3 alkoxycarbonylamino, phenoxycarbonylamino, C1-3 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl,
25 morpholinyl, thiomorpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

30

or R₅ is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C1-3 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl,
5 pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl,
thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl,
benzthiazolyl, benzoxazolyl, C1-3 alkanoyl, aroyl, C1-3alkanoyloxy, phenoxy,
benzyloxy, C1-3 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl
wherein the nitrogen atom may be independently mono or di-substituted by C1-3
10 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl,
thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,
or R_e is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen
15 atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl,
piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl,
pyrimidinyl, benzimidazolyl or benzthiazolyl,
or R_e is C1-3 alkoxycarbonylamino, phenoxycarbonylamino, C1-3
alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino,
20 phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-3
alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,
oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or
benzthiazolyl,
25 or R_e is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R_e may be further
optionally substituted by one or more R_f;
and

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, bromo,
hydroxy, oxo, carboxy or carboxamide;

30

R₁ and R₆ of the formula (Ia) or Formula (Ib) form a bicyclic ring having one 5 or 6 membered aromatic or nonaromatic heterocyclic ring fused to a second 5-6 membered heteroaryl, heterocycle or phenyl ring; wherein each ring is optionally independently substituted by one or two R₇.

5

In yet a further embodiment of the invention, there are provided novel compounds of the formulas (Ia) and (Ib) as described immediately above, and wherein:

10

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl or tetrahydropyranyl each ring being substituted with one or more R₅;

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

15

20

R_a is a bond, methyl, ethyl, cyclopropyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, acetyl, acetoxyl, phenoxy, benzyloxy, methoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

25

or R_a is acetylamino, benzoylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

30

or R_a is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

or R_a is fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_a may be further optionally substituted by one or more R_b;

5 R_b is methyl, cyclopropyl, phenyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is methyl, ethyl, n-propyl, i-propyl, C5-6 cycloalkyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, thienyl, oxazolyl, thiazolyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, 15 methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or aryl, or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be 20 oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or aryl, or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom 25 may be independently mono or di-substituted by methyl, ethyl or phenyl, or R_c is fluoro, chloro or oxo, R_c may be further optionally substituted by one or more R_d;

30 R_d is methyl, cyclopropyl, phenyl, methoxy, fluoro, chloro or oxo;

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-4alkoxyC1-4alkyl, C1-4alkylaminoC1-4alkyl, C1-4alkylthioC1-4alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-4 alkoxy,phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-2alkanoyl, benzoyl, naphthoyl, C1-2alkanoyloxy, benzyloxy, C1-2 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R₆;

R₆ is C1-3 alkyl, C1-2 alkoxy, C3-6 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-2 alkanoyl, aroyl, C1-2alkanoyloxy, phenoxy, benzyloxy, C1-2 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2

alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl,
thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
or R_e is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen
atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl,
piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or
pyrimidinyl,
or R_e is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2
alkylcarbamoxyloxy, phenylcarbamoxyloxy, C1-2 alkylsulfonylamino,
phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-2
alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,
oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
or R_e is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R_e may be
further optionally substituted by one or more R_f,

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro,
chloro, hydroxy, oxo, carboxy or carboxamide and

R₁ and R₆ of the formula (Ia) or Formula (Ib) form a bicyclic ring having one 5-6
membered aromatic or nonaromatic heterocyclic ring fused to a phenyl ring;
wherein each ring is optionally independently substituted by one or two R₇.

In yet still a further embodiment of the invention, there are provided novel compounds of
the formula (Ia) or formula (Ib) as described immediately above, and wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl, azepan-3-yl, azepan-4-yl or tetrahydropyran-4-yl, each ring being optionally substituted with one or more R₅;

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, cyclopropyl, cyclohexyl, phenoxy, phenyl,
5 benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is methyl, phenyl, thienyl, methoxy, acetyl, acetoxy, phenoxy, benzyloxy,
10 methoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
or R_a is acetylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,
15 or R_a is methoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
or R_a is fluoro, chloro, hydroxy, oxo, carboxy, cyano or carboxamide;

20 R₃ is a bond, methyl, ethyl, n-propyl, propenyl, butenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, i-propyl, cyclohexyl, cyclopentyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl,
25 bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro, chloro or oxo;

30

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-2alkoxyC1-2alkyl, C1-2alkylaminoC1-2alkyl, C1-2alkylthioC1-2alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-2 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is methyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidinyl, morpholinyl, indolyl, thienyl, pyridinyl, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_e is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R_e is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino,

phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further


5 optionally substituted by one or more R_f ,

and

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo;

10

R_1 and R_6 of the formula (Ia) or Formula (Ib) form the bicyclic ring



; wherein W is $-S(O)_n-$, $-O-C(O)-$ or $-N-C(O)-$, n is 0, 1 or 2 and wherein

15 each ring is optionally independently substituted by one or two R_7 .

In a further embodiment of the invention, there are provided novel compounds of the formulas (Ia) and (Ib) as described immediately above, and wherein:

20

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl or tetrahydropyran-4-yl, each ring being substituted with one or more R_5 ;

R_1 is i-propyl, benzyloxy, cyclohexyl, phenyl, 4-(acetlamino)-phenyl, 4-

25 (methanesulfonylamino)-phenyl, 4-methoxyphenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-

fluorophenyl, 2-fluorophenyl, 2-fluoro-4-chlorophenyl, naphthyl, thienylmethyl,

piperidinyl, morpholinyl, pyrrolidinyl, piperazinyl, furanyl, thienyl, 5-chlorothieryl,

pyridin-4-yl, pyrazinyl, methylamino, ethylamino, dimethylamino or diethylamino;

R₃ is ethyl, n-propyl, propenyl, butenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

5 R_c is methyl, cyclohexyl, cyclopentyl, indanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro or chloro;

10 R₅ is a bond, carbonyl, methyl, ethyl, n-propyl, n-butyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, piperidiny, tetrahydropyranyl, pyrimidiny, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy, R₅ may be further optionally substituted by one or more R_c;

15 R_e is methyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, thienyl, 5-methylthienyl, methoxy, phenoxy, benzyloxy, piperidiny, pyridiny, indolyl, 1-(tolyl-sulfonyl)-indolyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, phenyl or benzyl, or R_e is hydroxy, fluoro, chloro, oxo, dimethylamino or trifluoromethyl;

20

and

n is 2.

25 In another embodiment of the invention, there are provided novel compounds of the formulas (Ia) and (Ib) as described for the broadest generic aspect above and wherein:

R₁ and R₆ remain acyclic,

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R₅;

- 5 R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

10

R_a is a bond, C1-3 alkyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxycarbonyl, aryloxy carbonyl, aroyloxy, carbamoyl wherein the nitrogen atom
15 may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
20 independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkoxycarbonylamino, aryloxy carbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein
25 the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

30

R_b is methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, methoxy, ethoxy, n-propoxy, i-propoxy, phenoxy,

benzyloxy, fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₂ is hydrogen;

5

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl,
10 bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl,
bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-
tetrahydronaphthyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl,
imidazolyl, pyrimidinyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, C1-3
alkoxy, phenoxy, naphthyloxy, C1-2 alkanoyl, benzoyl, C1-2 alkoxycarbonyl,
15 phenoxycarbonyl, C1-2alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen
atom may be independently mono or di-substituted by C1-3 alkyl or aryl,
or R_c is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen
20 atom may be independently substituted by C1-3 alkyl or aryl,
or R_c is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2
alkylcarbamoyloxy, arylcarbamoyloxy, C1-2 alkylsulfonylamino,
phenylsulfonylamino, C1-2alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-3
25 alkyl or phenyl,
or R_c is halogen, hydroxy, oxo, carboxy or cyano, R_c may be further optionally
substituted by one or more R_d;

R_d is methyl, cyclopropyl, cyclohexyl, phenyl, benzyl, methoxy, phenoxy,
30 benzyloxy, benzoyl, fluoro, chloro, oxo or cyano;

R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-3alkanoyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-3 alkoxycarbonylamino, phenoxycarbonylamino, C1-3 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R₅ may be further optionally substituted by one or more R₆;

R₆ is C1-3 alkyl, C1-3 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl,

benzthiazolyl, benzoxazolyl, C1-3 alkanoyl, aroyl, C1-3alkanoyloxy, phenoxy, benzyloxy, C1-3 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl

wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl,

thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur

atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl,

piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-3 alkoxycarbonylamino, phenoxycarbonylamino, C1-3

alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino,

phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino

wherein the nitrogen atom may be independently mono or di-substituted by C1-3

alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,

oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or

benzthiazolyl,

or R_e is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R_e may be further

optionally substituted by one or more R_f;

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide;

R₆ is

hydroxy, nitrile or

a C1-5 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH, or S(O)₂ and wherein said chain is optionally independently substituted with 1-2 oxo groups,

-NH₂, one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl,

piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl,

isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl,

benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl,
benzoxazolyl or quinoxalinyl;
and

5 X is O.

In another embodiment of the invention, there are provided novel compounds of the
formula (Ia) and (Ib) as described immediately above, and wherein:

10

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, ethoxy, cyclopropyl, cyclopentyl,
cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl,
thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl
15 or amino; wherein R₁ is optionally substituted by one or more R_a;

20

R_a is a bond, methyl, ethyl, cyclopropyl, phenyl, pyrrolidinyl, piperidinyl,
morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, acetyl,
acetoxyl, phenoxy, benzyloxy, methoxycarbonyl, phenoxycarbonyl, benzoyloxy,
carbamoyl wherein the nitrogen atom may be independently mono or di-
substituted by methyl, ethyl or phenyl,

25

or R_a is acetylamino, benzoylamino, methylthio, phenylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen
atom may be independently substituted by methyl, ethyl or phenyl,

30

or R_a is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy,
phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino,
methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom
may be independently mono or di-substituted by methyl or phenyl,
or R_a is fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or
carboxamide, R_a may be further optionally substituted by one or more R_b;

R_b is methyl, cyclopropyl, phenyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl

5 wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, i-propyl, C5-6 cycloalkyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, thienyl, oxazolyl, thiazolyl, indolyl, benzofuranyl, 10 benzothienyl, benzthiazolyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or aryl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be 15 oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or aryl,

or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, 20 methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl, or R_c is fluoro, chloro or oxo, R_c may be further optionally substituted by one or more R_d;

25 R_d is methyl, cyclopropyl, phenyl, methoxy, fluoro, chloro or oxo;

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-4alkoxyC1-4alkyl, C1-4alkylaminoC1-4alkyl, C1-4alkylthioC1-4alkyl wherein the sulfur atom may be oxidized to a sulfoxide or 30 sulfone, C1-4 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl,

piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocycloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-2alkanoyl, benzoyl, naphthoyl, C1-2alkanoyloxy, benzyloxy, C1-2 alkoxycarbonyl, benzyloxycarbonyl, 5 phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized 10 to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-2 alkoxycarbonylamino, phenoxy carbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2 15 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

20 R_e is C1-3 alkyl, C1-2 alkoxy, C3-6 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-2 alkanoyl, aroyl, C1-2alkanoyloxy, phenoxy, 25 benzyloxy, C1-2 alkoxycarbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R_e is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur 30 atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen

atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R_e is C1-2 alkoxy-carbonylamino, phenoxy-carbonylamino, C1-2

alkyl-carbamoyloxy, phenyl-carbamoyloxy, C1-2 alkyl-sulfonylamino,

phenyl-sulfonylamino, C1-2 alkyl-aminosulfonyl, phenyl-aminosulfonyl, amino

wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R_e is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f,

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide and

R₆ is

nitrile or

a C1-5 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH, or

S(O)₂ and wherein said chain is optionally independently substituted with oxo, -NH₂, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyridinyl, pyrimidinyl or pyrazinyl.

In yet another embodiment of the invention, there are provided novel compounds of the formula (Ia) or formula (Ib) as described immediately above, and wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl, azepan-3-yl, azepan-4-yl or tetrahydropyran-4-yl, each ring being optionally substituted with one or more R₅;

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, cyclopropyl, cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

5

R_a is methyl, phenyl, thienyl, methoxy, acetyl, acetoxy, phenoxy, benzyloxy, methoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

10

or R_a is acetylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,

or R_a is methoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

15

or R_a is fluoro, chloro, hydroxy, oxo, carboxy, cyano or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, propenyl, butenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

20

R_c is methyl, ethyl, n-propyl, i-propyl, cyclohexyl, cyclopentyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-

tetrahydronaphthyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro, chloro or oxo;

25

and

wherein the configuration at the stereocenter defined by R₂ and R₃ when they are different and the carbon they are attached to is defined as L; and

30

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-2alkoxyC1-2alkyl, C1-2alkylaminoC1-2alkyl, C1-2alkylthioC1-2alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-2 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_c;

R_c is methyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidinyl, morpholinyl, indolyl, thienyl, pyridinyl, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R_c is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino,

phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further
5 optionally substituted by one or more R_f;

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo;

R₆ is

nitrile or

10 a C1-5 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH, or S(O)₂ and wherein said chain is optionally independently substituted with oxo, -NH₂, morpholinyl or piperazinyl.

15 In yet still another embodiment of the invention, there are provided novel compounds of the formulas (Ia) and (Ib) as described immediately above, and wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl or tetrahydropyran-4-yl, each ring being substituted with one or more R₅;

20

R₁ is i-propyl, benzyloxy, cyclohexyl, phenyl, 4-(acetylamino)-phenyl, 4-(methanesulfonylamino)-phenyl, 4-methoxyphenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-chlorophenyl, naphthyl, thienylmethyl, piperidinyl, morpholinyl, pyrrolidinyl, piperazinyl, furanyl, thienyl, 5-chlorothieryl,
25 pyridin-4-yl, pyrazinyl, methylamino, ethylamino, dimethylamino or diethylamino;

R₃ is ethyl, n-propyl, propenyl, butenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, cyclohexyl, cyclopentyl, indanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro or chloro;

5 R₅ is a bond, carbonyl, methyl, ethyl, n-propyl, n-butyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, piperidinyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy, R₅ may be further optionally substituted by one or more R_c;

10

R_c is methyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, thienyl, 5-methylthienyl, methoxy, phenoxy, benzyloxy, piperidinyl, pyridinyl, indolyl, 1-(tolyl-sulfonyl)-indolyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, phenyl or benzyl,

15

or R_c is hydroxy, fluoro, chloro, oxo, dimethylamino or trifluoromethyl;

and

R₆ is acetyl, C1-3alkylaminocarbonyl or C1-3alkoxycarbonyl.

20

In yet a further embodiment of the invention, there are provided novel compounds of the formulas (Ia) and (Ib) as described immediately above, and wherein:

25 Het is piperidin-4-yl or pyrrolidin-3-yl;

R₁ is morpholin-4-yl, p-fluorophenyl or p-methoxyphenyl;

R₃ is methyl, propyl, n-pentyl or cyclohexyl

30

and

R₆ is acetyl, ethylaminocarbonyl or ethoxycarbonyl.

The activity of particular compounds disclosed herein against cathepsin K may be determined without undue experimentation by one of ordinary skill in the art in view of the art, the guidance provided throughout this specification and by the screens described in the section entitled "Assessment of Biological Properties."

5

The following subgeneric aspect of the compounds of the formulas (Ia) and (Ib) is postulated to possess Cathepsin K activity:

The broadest embodiment of the formula (Ia) and (Ib) as described hereinabove and
10 wherein

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R₅;

15

R₁ is a bond, C1-4 alkyl, C1-4 alkoxy, cyclopropyl, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or
20 more R_a;

R_a is methyl, ethyl, propyl, i-propyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, ethoxy, acetyl, acetoxy, phenoxy, naphthyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, phenoxycarbonyl, naphthyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
25 or R_a is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ethylthio wherein the sulfur atom may be
30

oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
5 or R_a is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, naphthylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, naphthylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, naphthylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl,
10 phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

15 R_b is methyl, ethyl, cyclopropyl, cyclohexyl, phenyl, methoxy, ethoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₂ is hydrogen or methyl;

20 R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

25 R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, naphthyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or
30 di-substituted by methyl or phenyl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,

5 or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl, or R_c is chloro, fluoro, hydroxy, oxo, carboxy or cyano;

10 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl or tetrahydrothiophenyl;

15 R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, naphthyloxy, cyclopropyl, cyclopentyl, cyclohexyl, 20 phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is acetylamino, benzoylamino, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

30 or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, phenylsulfonylamino, phenylaminosulfonyl,

amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

5

R_e is methyl ethyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidiny, morpholinyl, indolyl, thienyl, pyridinyl, methoxy, ethoxy, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be

10

independently mono or di-substituted by methyl, ethyl or phenyl, or R_e is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

15

or R_e is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

20

or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f;

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo.

25

Preferred cathepsin K inhibitors are those as described immediately above and wherein:

R₁ is a bond, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, benzyloxy, cyclopropyl, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl,

30

benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino;
wherein R₁ is optionally substituted by one or more R_a;

5 R_a is methyl, cyclopropyl, phenyl, halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
15 or R_c is acetylamino, benzoylamino, methylthio, methoxycarbonylamino, methylcarbamoyloxy, methylsulfonylamino, methylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl,
or R_c is fluoro or oxo;

20 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, pyrrolidinyl or piperidinyl;

25 R₅ is methyl, ethyl, n-propyl, n-butyl, n-pentyl, 2-pentyl, 3-pentyl, phenethyl, phenpropyl, 2,2-dimethylpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methylbenzyl, 3-methylbenzyl, 4-methylbenzyl, 2,6-dimethylbenzyl, 2,5-dimethylbenzyl, 2,4-dimethylbenzyl, 2,3-dimethylbenzyl, 3,4-dimethylbenzyl, 3,5-dimethylbenzyl, 2,4,6-trimethylbenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-
30 phenoxybenzyl, 3-phenoxybenzyl, 4-phenoxybenzyl, 2-benzyloxybenzyl, 3-

benzyloxybenzyl, 4-benzyloxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 2,6-difluorobenzyl, 2,5-difluorobenzyl, 2,4-difluorobenzyl, 2,3-difluorobenzyl, 3,4-difluorobenzyl, 3,5-difluorobenzyl, 2,4,6-trifluorobenzyl, 2-trifluoromethylbenzyl, 3-trifluoromethylbenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, indanylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, 5-methylthienylmethyl, piperidinyl, piperidinylcarbonyl, pyridinylcarbonyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, methylcarbamoyl, phenylcarbamoyl, benzylcarbamoyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy.

10

Most preferred cathepsin K inhibitors are those as described immediately above and wherein:

R_1 is methoxy, benzyloxy, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R_1 is optionally substituted by one or more R_a ;

R_a is methyl, phenyl, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R_3 is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl or benzyl wherein R_3 is optionally substituted by one or more R_c ;

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, acetylamino, methylthio, methylsulfonylamino or fluoro;

R_2 and R_3 together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl or tetrahydrofuranlyl;

R₅ is methyl, ethyl, n-propyl, n-butyl, phenethyl, phenpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclohexyl, cyclopropylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 4-fluorobenzyl, 3,5-difluorobenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, phenylcarbamoyl, phenylsulfonylamino or fluoro.

Most preferred cathepsin K inhibitors are those as described immediately above and
10 wherein:

Het is pyrrolidinyl, piperidinyl or tetrahydropyranyl;

R₁ is benzyloxy, phenoxy, naphthyloxy, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyridinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or phenylamino;

R₃ is n-propyl, i-butyl, propenyl, i-butenyl or 2,2-dimethylpropyl;

20 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, or cycloheptyl;

R₅ is methyl, ethyl, n-propyl, phenethyl, t-butyl, i-propyl, i-butyl, cyclohexyl, cyclohexylmethyl, benzyl, 4-fluorobenzyl, naphthylmethyl, acetyl, benzoyl or
25 benzyloxycarbonyl.

The following are representative compounds of the invention which possess desirable inhibition activity of Cathepsin S in a cell based assay as described in Riese, R.J. et al.,
30 Immunity, 1996, 4, 357-366, incorporated herein by reference.

'2-[(Acetylimino-phenyl-methyl)-amino]-N-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide;

5 '({1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethylamino}-morpholin-4-yl-methylene)-carbamic acid ethyl ester;

'2-(N-Cyano-benzimidoyl-amino)-N-(4-cyano-1-methyl-piperidine-4-yl)-3-cyclohexyl-propionamide;

10 'N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-phenyl-methyl)-amino]-propionamide;

'N-[4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

'N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

20 N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

25 '2-(1,1-Dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

and the pharmaceutically acceptable derivatives thereof.

30

Another embodiment of the invention provides for the following compounds which have demonstrated potent inhibition of Cathepsin S in a cell based assay at concentrations of 50 nM or less.

35 {[1-(4-Cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylimino]-morpholin-4-yl-methyl}-carbamic acid ethyl ester;

N-(4-Cyano-methyl-piperidin-4-yl)-3-cyclohexyl-2-(3-oxo-3H-isoindol-1-ylamino)-propionamide;

40

4,4-Dimethyl-2-(3-oxo-3H-isoindol-1-ylamino)-pentanoic acid-(4-cyano-1-propyl-piperidin-4-yl)-amide;

5 N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(2-oxo-2H-benzo[e][1,3] oxazin-4-ylamino)-propionamide;

{[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-piperidin-1-yl-methyl}-carbamic acid ethyl ester;

10 '2-[(Acetylimino-phenyl-methyl)-amino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide;

'{[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylen}-carbamic acid ethyl ester;

15 '2-[(Acetylimino-phenyl-methyl)-amino]-N-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide;

20 '({[1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylene)-carbamic acid ethyl ester;

'N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-phenyl-methyl)-amino]-propionamide;

25 'N-[4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

'N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

30 N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

35 N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide and

'2-(1,1-Dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide.

40 Any compounds of this invention containing one or more asymmetric carbon atoms may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. All such isomeric forms of these compounds are expressly

included in the present invention. Each stereogenic carbon may be in the R or S configuration unless otherwise specified, or a combination of configurations.

Some of the compounds of formulas (I), (II), (Ia) and (Ib) can exist in more than one
5 tautomeric form. The invention includes all such tautomers.

It shall be understood by one of ordinary skill in the art that all compounds of the invention are those which are chemically stable.

10 The invention includes pharmaceutically acceptable derivatives of compounds of formula (I), (II), (Ia) and (Ib). A "pharmaceutically acceptable derivative" refers to any pharmaceutically acceptable acid, salt or ester of a compound of this invention, or any other compound which, upon administration to a patient, is capable of providing (directly or indirectly) a compound of this invention, a pharmacologically active metabolite or
15 pharmacologically active residue thereof.

In addition, the compounds of this invention include prodrugs of compounds of the formulas (I), (II), (Ia) and (Ib). Prodrugs include those compounds that, upon simple transformation, are modified to produce the compounds of the invention. Simple
20 chemical transformations include hydrolysis, oxidation and reduction which occur enzymatically, metabolically or otherwise. Specifically, when a prodrug of this invention is administered to a patient, the prodrug may be transformed into a compound of formula (I), (II), (Ia) and (Ib), thereby imparting the desired pharmacological effect.

25 In order that the invention herein described may be more fully understood, the following detailed description is set forth. As used herein, the following abbreviations are used:

BOC or t-BOC is tertiary-butoxycarbonyl;

t-Bu is tertiary-butyl;

30 DMF is dimethylformamide;

EtOAc is ethyl acetate;

THF is tetrahydrofuran;

Ar is argon;

EDC is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride and

HOBT is 1-hydroxybenzotriazole.

5

Also, as used herein, each of the following terms, used alone or in conjunction with other terms, are defined as follows (except where noted to the contrary):

10 The term "alkyl" refers to a saturated aliphatic radical containing from one to ten carbon atoms or a mono- or polyunsaturated aliphatic hydrocarbon radical containing from two to twelve carbon atoms. The mono- or polyunsaturated aliphatic hydrocarbon radical containing at least one double or triple bond, respectively. "Alkyl" refers to both branched and unbranched alkyl groups. Examples of "alkyl" include alkyl groups which
15 are straight chain alkyl groups containing from one to eight carbon atoms and branched alkyl groups containing from three to eight carbon atoms. Other examples include lower alkyl groups which are straight chain alkyl groups containing from one to six carbon atoms and branched alkyl groups containing from three to six carbon atoms. It should be understood that any combination term using an "alk" or "alkyl" prefix refers to analogs according to the above definition of "alkyl". For example, terms such as "alkoxy",
20 "alkythio" refer to alkyl groups linked to a second group via an oxygen or sulfur atom. "Alkanoyl" refers to an alkyl group linked to a carbonyl group (C=O). Each alkyl or alkyl analog described herein shall be understood to be optionally partially or fully halogenated.

25

The term "cycloalkyl" refers to the cyclic analog of an alkyl group, as defined above. Examples of cycloalkyl groups are saturated or unsaturated nonaromatic cycloalkyl groups containing from three to eight carbon atoms, and other examples include cycloalkyl groups having three to six carbon atoms. Each cycloalkyl described herein
30 shall be understood to be optionally partially or fully halogenated.

The term "aryl" refers to phenyl and naphthyl.

The term "halo" refers to a halogen radical selected from fluoro, chloro, bromo or iodo. Representative halo groups of the invention are fluoro, chloro and bromo.

- 5 The term "heteroaryl" refers to a stable 5-8 membered (but preferably, 5 or 6 membered) monocyclic or 7-12 membered polycyclic, preferably bicyclic aromatic heterocycle radical. Each heterocycle consists of carbon atoms and from 1 to 4 heteroatoms chosen from nitrogen, oxygen and sulfur. The heterocycle may be attached by any atom of the cycle, which results in the creation of a stable structure. Examples of "heteroaryl" include
- 10 radicals such as furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indoliziny, indolyl, isoindolyl, benzofuranyl, benzothienyl, indazolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, purinyl, quinoliziny, quinoliny, isoquinoliny, cinnoliny, phthalazinyl, quinazolinyl,
- 15 quinoxaliny, naphthyridiny, pteridiny, carbazolyl, acridiny, phenazinyl, phenothiazinyl and phenoxazinyl,

- The term "heterocycle" refers to a stable 4-8 membered (but preferably, 5 or 6 membered) monocyclic or 7-12 membered polycyclic, preferably bicyclic heterocycle
- 20 radical which may be either saturated or unsaturated, and is non-aromatic. Each heterocycle consists of carbon atoms and from 1 to 4 heteroatoms chosen from nitrogen, oxygen and sulfur. The heterocycle may be attached by any atom of the cycle, which results in the creation of a stable structure. Examples of "heterocycle" include radicals such as pyrroliny, pyrrolidinyl, pyrazoliny, pyrazolidiny, piperidinyl, morpholiny,
- 25 thiomorpholiny, pyranyl, thiopyranyl, piperazinyl, indoliny, azetidiny, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, hexahydropyrimidinyl, hexahydropyridazinyl, 1,4,5,6-tetrahydropyrimidin-2-ylamine, dihydro-oxazolyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidiny-1,1-dioxide and imidazolidiny-2,4-dione.

30

The terms "heterocycle", "heteroaryl" or "aryl", when associated with another moiety, unless otherwise specified shall have the same meaning as given above. For example, "aroyl" refers to phenyl or naphthyl linked to a carbonyl group (C=O).

5 Each aryl or heteroaryl unless otherwise specified includes its partially or fully hydrogenated derivative. For example, quinolinyl may include decahydroquinolinyl and tetrahydroquinolinyl, naphthyl may include its hydrogenated derivatives such as tetrahydronaphthyl. Other partially or fully hydrogenated derivatives of the aryl and heteroaryl compounds described herein will be apparent to one of ordinary skill in the art.

10

The term heterocycle as it pertains to "Het" shall to be understood to mean a stable non-aromatic spiroheterocycle, 4-8 membered (but preferably, 5 or 6 membered) monocyclic, 7-12 membered polycyclic, preferably bicyclic heterocycle radical which may be either saturated or unsaturated or a C6-C10 bridged bicyclo wherein one or more carbon atoms are optionally replaced by a heteroatom. Each heterocycle consists of carbon atoms and from 1 to 4 heteroatoms chosen from nitrogen, oxygen and sulfur. The heterocycle may be attached by any atom of the cycle, which results in the creation of a stable structure. Examples of "Het" include the following heterocycles: azepanyl, piperidinyl, pyrrolidinyl, azetidiny, oxepanyl, tetrahydropyranyl, tetrahydrothiopyranyl, 20 tetrahydrofuranyl, oxetanyl, azocanyl, oxocanyl, 1,3-diazocanyl, 1,4-diazocanyl, 1,5-diazocanyl, 1,3-dioxocanyl, 1,4-dioxocanyl, 1,5-dioxocanyl, 1,3-oxazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,3-diazepanyl, 1,4-diazepanyl, 1,3-dioxepanyl, 1,4-dioxepanyl, 1,3-oxazepanyl, 1,4-oxazepanyl, 1,2-thiazocanyl-1,1-dioxide, 1,2,8-thiadiazocanyl-1,1-dioxide, 1,2-thiazepanyl-1,1-dioxide, 1,2,7-thiadiazepanyl-1,1-dioxide, 25 tetrahydrothiophenyl, hexahydropyrimidinyl, hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, pyrazolidinyl, dihydro-oxazolyl, dihydrothiazolyl, dihydroimidazolyl, isoxazoliny, oxazolidinyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide, imidazolidinyl-2,4-dione, imidazolidinyl, morpholinyl, dioxanyl, tetrahydropyridinyl, thiomorpholinyl, thiazolidinyl, dihydropyranyl, dithianyl, decahydro-quinolinyl, decahydro-isoquinolinyl, 30 1,2,3,4-tetrahydro-quinolinyl, indolinyl, octahydro-quinoliziny, dihydro-indoliziny,

octahydro-indoliziny, octahydro-indolyl, decahydroquinazoliny, decahydroquinoxaliny, 1,2,3,4-tetrahydroquinazoliny or 1,2,3,4-tetrahydroquinoxaliny, aza-bicyclo[3.2.1]octane, aza-bicyclo[2.2.1]heptane, aza-bicyclo[2.2.2]octane, aza-bicyclo[3.2.2]nonane, aza-bicyclo[2.1.1]hexane, aza-bicyclo[3.1.1]heptane, aza-bicyclo[3.3.2]decane and 2-oxa or 2-thia-5-aza-bicyclo[2.2.1]heptan each heterocyclic ring being substituted with one or more R_5 . The substituent R_5 is defined above.

As used herein above and throughout this application, "nitrogen" and "sulfur" include any oxidized form of nitrogen and sulfur and the quaternized form of any basic nitrogen.

10

In order that this invention be more fully understood, the following examples are set forth. These examples are for the purpose of illustrating preferred embodiments of this invention, and are not to be construed as limiting the scope of the invention in any way.

15

The examples which follow are illustrative and, as recognized by one skilled in the art, particular reagents or conditions could be modified as needed for individual compounds. Starting materials used in the scheme below are either commercially available or easily prepared from commercially available materials by those skilled in the art.

20

GENERAL SYNTHETIC METHODS

The invention also provides processes of making the present novel compounds.

Compounds of the invention may be prepared by methods described below. Standard peptide coupling, protection and deprotection reactions (see for example M. Bodanszky, 1984, The Practice of Peptide Synthesis, Springer-Verlag) are employed in these syntheses and are incorporated herein by reference in their entirety.


25

COMPOUNDS OF THE FORMULAS (I) AND (II):

30

Compounds of the invention having formulas (I) and (II) may be prepared by Method A as illustrated in Scheme I.

Scheme I (Method A)



5

According to Method A a suitably protected amino acid bearing "Het" is allowed to react with ammonia under standard coupling conditions. An example of a suitable protecting group is the *t*-butoxycarbonyl (BOC) group. An example of standard coupling conditions would be combining the starting materials in the presence of a coupling reagent such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) with 1-hydroxybenzotriazole (HOBT), in a suitable solvent such as DMF or methylene chloride. A base such as N-


10

methyldmorpholine may be added. This is followed by deprotection to give amino acid amide III. An amino acid ester (IV) bearing R_2 , R_3 and optionally R_4 other than H is then reacted with an activated acid [$R_1C(O)L$] such as acid chloride ($L = Cl$) in the presence of a suitable base such as N,N-diisopropylethylamine to provide V. Alternately, one may
5 use the carboxylic acid [$R_1C(O)L$, $L = OH$] and activate using standard peptide coupling conditions, such as EDC and HOBT as described above. If R_4 is H in V, one may optionally react V with an alkyl halide in the presence of a suitable base such as sodium hydride, in a suitable solvent such as DMF or THF to provide V in which R_4 is alkyl. Conversion to the carboxylic acid provides VI. Standard peptide coupling of III and VI,
10 followed by dehydration of the amide provides the desired nitrile I or II. An example of suitable dehydration conditions is cyanuric chloride in DMF.

In a variation (Method B) illustrated in Scheme II, an amino acid amide bearing "Het" is coupled with an amine-protected amino acid bearing R_2 and R_3 . A suitable protecting
15 group and coupling conditions would be as described above. Deprotection is then followed by reaction with $R_1C(O)L$ (as described in Method A). Conversion of the amide to the nitrile as above provides I or II.

Scheme II (Method B)


20



Compounds of the invention having formula (I) and (II) may also be prepared by Method

5 C as illustrated in Scheme III.

Scheme III (Method C)




10

In this variation (Method C) an amino nitrile bearing "Het" is coupled with an amine protected amino acid bearing R₂ and R₃. A suitable protecting group and coupling

conditions are described above. Deprotection is then followed by reaction with $R_1C(O)L$ as described above to furnish the nitrile (I/II).

Compounds of the invention having formulas (I) and (II) may also be prepared by as
5 outlined below in Scheme IV (Method D).

Scheme IV (Method D)




10

In a further variation (Method D) illustrated in Scheme IV, an amino acid ester (IV) bearing R_2 , R_3 and optionally R_4 other than H is reacted with $R_1C(O)L$ as described in Method A. Conversion to the carboxylic acid provides VI. Standard peptide coupling of
15 an amino nitrile bearing "Het" with VI yields the desired nitrile (I/II).

The intermediate aminonitrile used in Methods C, and D above may be prepared as outlined in Scheme V

20 Scheme V



In this method, a ketone bearing "Het" is reacted with an a primary amine or an ammonium salt, such as ammonium chloride, and a cyanide salt, such as potassium cyanide or sodium cyanide, in a suitable solvent, such as water or a solution of ammonia in methanol, at about room temperature to reflux temperature.

In each of the methods described above, required starting materials are either commercially available or easily prepared by those skilled in the art, for example see:

10 Leung, M.-k.; Lai, J.-L.; Lau, K.-H.-; Yu, H.-h.; Hsiao, J.-J. *J. Org. Chem.* **1996**, *61*, 4175-4179.

Mee, J. D. *J. Org. Chem.* **1975**, *40*, 2135-2136.

Micovic, I. V.; Roglic, G. M.; Ivanovic, M. D.; Dosen-Micovic, L.; Kiricojevic, V. D.;

Popovic, J. B. *J. Chem. Soc., Perkin Trans. 1*, **1996**, 2041-2050.

15 Tornus, I.; Schaumann, E. *Tetrahedron* **1996**, *52*, 725-732.

Jadhav, P. K.; Woerner, F. J. *Tetrahedron Letters* **1995**, *36*, 6383-6386.

Kochhar, K. S.; et al. *Tetrahedron Letters* **1984**, *25*, 1871-1874.

Fordon, K. J.; Crane, C. G.; Burrows, C. J. *Tetrahedron*


Letters **1994**, *35*, 6215-6216. These references are incorporated herein by reference in

20 their entirety,


COMPOUNDS OF THE FORMULAS (Ia) AND (Ib):

The invention also provides processes of making the present novel compounds of formula

25 (Ia) and (Ib). Compounds of the invention may be prepared by methods described below.




(Ia)



(Ib)

A key intermediate in the preparation of compounds of formula (Ia) and (Ib) is the
 5 dipeptide nitrile intermediate (VII).




(VII)

10 The synthesis of intermediates of formula (VII) is described in US provisional patent application no. 60/153,738 and outlined below in Schemes VI and VII.

Scheme VI


15



As illustrated in Scheme VI, an amino acid bearing a suitable protecting group R' (VIII), is reacted with an amino nitrile (IX) under suitable coupling conditions. An example of a suitable protecting group is the *t*-butoxycarbonyl (BOC) group. An example of standard coupling conditions would be combining the starting materials in the presence of a coupling reagent such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) with 1-hydroxybenzotriazole (HOBT), in a suitable solvent such as DMF or methylene chloride. A base such as N-methylmorpholine may be added. This is followed by deprotection to give amino acid nitrile VII.

- 10 The intermediate aminonitrile (IX) used in Scheme VI above may be prepared as outlined in Scheme VII.

Scheme VII




- 15 In this method, a ketone bearing "Het" (XI) is reacted with an a primary amine or an ammonium salt, such as ammonium chloride, and a cyanide salt, such as potassium cyanide or sodium cyanide, in a suitable solvent, such as water or a solution of ammonia in methanol, at about room temperature to reflux temperature.

20

Compounds having formula (Ia/Ib) may be prepared by Methods E-H, as illustrated in Schemes VIII-IX.


- 25 Scheme VII (Method E)



According to Method E, a dipeptide nitrile intermediate (VII), or a basic salt thereof, is allowed to react with (XII) in the presence of a suitable coupling agent to provide the desired product (Ia/Ib). Suitable reaction conditions are known to those skilled in the art and some examples of suitable coupling agents include 2-chloro-1-methylpyridinium iodide (Yong, Y.F. et al., J. Org. Chem. 1997, 62, 1540), phosgene or triphosgene (Barton, D.H. et al., J. Chem. Soc. Perkin Trans. I, 1982, 2085), alkyl halides (Brand, E. and Brand, F. C., Org. Synth., 1955, 3, 440) carbodiimides (Poss, M. A. et al., Tetrahedron Lett., 1992, 40, 5933) and mercury salts (Su, W., Synthetic Comm., 1996, 26, 407 and Wiggall, K. J. and Richardson, S. K. J., Heterocyclic Chem., 1995, 32, 867).

Compounds having formulas (Ia) and (Ib) may also be prepared by Method B as illustrated in Scheme IV, where R is an alkyl or aryl group.

Scheme VIII (Method F)




According to Method F a dipeptide nitrile intermediate (VII), or a basic salt thereof, is allowed to react with XII, with or without an added base such as triethylamine, to provide the desired product (Ia/Ib). Suitable reaction conditions are known to those skilled in the art and examples of such amine additions may be found in the chemical literature, for example Haake, M. and Schummelfeder, B., Synthesis, 1991, 9, 753; Dauwe, C. and Buddrus, J., Synthesis 1995, 2, 171; Ried, W. and Piechaczek, D., Justus Liebigs Ann. Chem. 1966, 97, 696 and Dean, W. D. and Papadopoulos, E. P., J. Heterocyclic Chem., 1982, 19, 1117.

The intermediate XII is either commercially available or can be synthesized by methods known to those skilled in the art and described in the literature, for example Francesconi, I. et. al., J. Med. Chem. 1999, 42, 2260; Kurzer, F., Lawson, A., Org. Synth. 1963, 645,
 5 and Gutman, A. D. US 3984410, 1976.

In a similar reaction, intermediate XIV having a halogen or other suitable leaving group (X) may be used in place of intermediate XIII, as illustrated in Method G, Scheme IX.:


10 Scheme IX (Method G)



According to Method G, a dipeptide nitrile intermediate, or a basic salt thereof, is allowed
 15 to react with intermediate XIV, with or without an added base such as triethylamine, to provide the desired product (Ia/Ib). Procedures for accomplishing this reaction are known to those skilled in the art and described in the chemical literature (for example, Dunn, A. D., Org. Prep. Proceed. Int., 1998, 30, 709; Lindstroem, S. et al., Heterocycles, 1994, 38, 529; Katritzky, A. R. and Saczewski, F., Synthesis, 1990, 561; Hontz, A. C.
 20 and Wagner, E. C., Org Synth., 1963, IV, 383; Stephen, E. and Stephen, H., J. Chem. Soc., 1957, 490).


Compounds having formula (Ia/Ib) in which R₁ is an amine may also be prepared by
 25 Method H as illustrated in Scheme X.

Scheme X (Method H)



According to Method H, a carbodiimide (XV) derivative of (VII) is allowed to react with an amine (R_1) to provide the desired guanidine (Ia/Ib) product. The conversion of amines to carbodiimides is known to those in the art and described in the literature (for example, Pri-Bar, I. and Schwartz, J., J. Chem. Soc. Chem. Commun., 1997, 347; Hirao, T. and Saegusa, T., J. Org. Chem., 1975, 40, 298). The reaction of carbodiimides with amine nucleophiles is also described in the literature (for example, Yoshiizumi, K. et al., Chem. Pharm. Bull., 1997, 45, 2005; Thomas, E. W. et al., J. Med. Chem., 1989, 32, 228; Lawson, A. and Tinkler, R. B., J. Chem. Soc. C, 1971, 1429).

In a modification of Method H, one may start with the thiourea XVI (formed by reaction of the corresponding amine with an isothiocyanate $R_6N=C=S$) and then form the corresponding carbodiimide (XV) *in situ* by reaction with a suitable desulfurizing agent, such as $HgCl_2$, in a suitable solvent such as DMF or acetonitrile.




Compounds of formula (Ib), where R_1 is an amine may be prepared using a general procedure described by M. Haake and B. Schummfelder (Synthesis, 1991, 753).

According to this procedure (Method I, Scheme XI), intermediate XVII bearing two suitable leaving groups Z, such as phenoxy groups, is reacted sequentially with amines R_1 and R_6R_3NH in a suitable solvent such as methanol or isopropanol to provide the desired

product. Reaction of the first amine may be carried out at about room temperature and reaction of the second amine is preferentially carried out with heating at the reflux temperature of the solvent. If XIII is allowed to react with a bifunctional nucleophile intermediate XVIII, where Y is a nucleophilic heteroatom such as N, O or S, one may obtain the product of formula (Ib) where R₁ and R₆ form a heterocyclic ring.

Intermediate XVII may be prepared by reaction of VII (R₄ = H) with dichlorodiphenoxymethane, which in turn, may be prepared by heating diphenyl carbonate with PCl₅ (R.L. Webb and C.S. Labow, J. Het. Chem., 1982, 1205).

10 Scheme XI (Method I)



15 In order that this invention be more fully understood, the following examples are set forth. These examples are for the purpose of illustrating embodiments of this invention, and are not to be construed as limiting the scope of the invention in any way.

The examples which follow are illustrative and, as recognized by one skilled in the art, particular reagents or conditions could be modified as needed for individual compounds.
 20 Starting materials used in the scheme below are either commercially available or easily prepared from commercially available materials by those skilled in the art.

5

SYNTHETIC EXAMPLES

EXAMPLE 1

10 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

(a) 4-Amino-4-cyano-1-methylpiperidine.

A solution of ammonium chloride (1.89 g, 35.37mmol) and potassium cyanide (2.30 g,
15 35.37 mmol) was prepared in 50 mL of water. 1-Methyl-4-piperidone (1.0 g, 8.84 mmol)
was added to the solution and stirred for 2 days. The solution was brought to pH 11 with
solid sodium carbonate and the reaction solution was extracted 3 x 100 mL of EtOAc.
The organic layer was dried over anhydrous Na₂SO₄, decanted and concentrated to an
orange oil (857 mg). ¹H NMR showed that the oil was a 2:1:1 mixture of the desired
20 aminonitrile, cyanohydrin and starting ketone. The crude mixture was used in the next
step without further purification.

(b) N-(4-morpholinecarbonyl)-L-cyclohexyl alanine methyl ester.

25 Methyl L-β-cyclohexylalanine hydrochloride (1.45 g, 6.54 mmol) was dissolved in 20
mL of DMF and 10 mL of Hunig's base was added to give a clear colorless solution. 4-
Morpholinecarbonyl chloride (1.17 g, 7.85 mmol) was added and the resulting reaction
was stirred at ambient temperature for 6 h. The reaction mix was concentrated *in vacuo*
and the residue was taken up in 200 mL of CH₂Cl₂ and washed with 1x100 mL of EtOAc
30 and 2x100 mL of brine. The organic layer was dried over Na₂SO₄, decanted, and
concentrated to a semi-solid (1.86 g) which was used in the next step without further
purification.

(c) N-(4-morpholinecarbonyl)-L-cyclohexyl alanine

N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine methyl ester (1.86 g, 6.23 mmol) was dissolved in 50 mL of MeOH to which was added 50 mL THF and 50 mL of water.

LiOH monohydrate (2.61 g, 62.3 mmol) was added to the reaction solution and the

5 reaction was monitored at 5 min and every 20 min thereafter using 5% MeOH in CH₂Cl₂.

The starting material was consumed at 2 h and the reaction was washed with 150 mL of diethyl ether with the organic layer being discarded. The aqueous layer was brought to pH 1 with concentrated HCl and the product was extracted with 2x100 mL of EtOAc.

The combined organic layers were dried over Na₂SO₄, decanted and concentrated to
10 white solid foam (1.63 g): ¹H NMR (CDCl₃) δ 8.90-7.90 (br, 1H), 5.05-4.99 (m, 1H), 4.55-4.39 (m, 1H), 3.71-3.62 (m, 4H), 3.50-3.36 (m, 4H), 1.90-0.83 (m, 13H).

(d) Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

15

N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine (350 mg, 1.23 mmol) was dissolved in 15 mL of DMF. EDC (235 mg, 1.23 mmol) and HOBT (166 mg, 1.23 mmol) were added and the resulting mixture was stirred at ambient temperature for 20 min during which time the solids went into solution. 4-Amino-4-cyano-1-methylpiperidine (310 mg of the

20 2:1:1 mixture of aminonitrile:cyanohydrin:ketone, \cong 1.1 mmol aminonitrile) was

dissolved in 5 mL of DMF, N-methylmorpholine was added to this solution (497 mg, 4.92 mmol), and the resultant solution added to the solution of the activated ester. The resulting mixture was stirred at ambient temperature for 16 h. The volatiles were

removed *in vacuo* and the resulting residue was dissolved in 200 mL of EtOAc and

25 washed sequentially with 2x200 mL saturated sodium bicarbonate and 1x100 mL brine.

The organic layer was dried over anhydrous Na₂SO₄, decanted, and concentrated to a thick oil. The oil was purified by column chromatography on SiO₂ using as eluent 100% CH₂Cl₂ to 12% MeOH in CH₂Cl₂ to give the desired product as a white powder (225 mg): ¹H NMR (CDCl₃) δ 7.55 (s, 1H), 5.13-5.08 (m, 1H), 4.40-4.20 (m, 1H), 3.77-3.62

30 (m, 4H), 3.51-3.33 (m, 4H), 2.88-2.55 (m, 2H), 2.53-2.39 (m, 2H), 2.30 (s, 3H), 2.10-0.83 (m, 17H).

Following the above procedures the following compounds can be synthesised;

5 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-naphthalen-2-yl-ethyl]-amide

Morpholine-4-carboxylic acid [2-(3-chloro-phenyl)-1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl) -ethyl]-amide

10 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-(3,4-dichloro-phenyl) -ethyl]-amide

Morpholine-4-carboxylic acid [2-(4-chloro-phenyl)-1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl) -ethyl]-amide

15 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl) -pentyl]-amide

20 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl) -3-methyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-1-phenyl-2,6-dioxo-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

25 Morpholine-4-carboxylic acid [1-(4-cyano-1-phenyl-2,6-dioxo-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-2-oxo-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

30 Morpholine-4-carboxylic acid [1-(4-cyano-2-oxo-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-2-oxo-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

- 5 Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-2-oxo-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(5-cyano-1,1-dioxo-1 λ^6 -[1,2]thiazinan-5-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

10

Morpholine-4-carboxylic acid [1-(5-cyano-1,1-dioxo-1 λ^6 -[1,2]thiazinan-5-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

- 15 Morpholine-4-carboxylic acid [1-(5-cyano-2-methyl-1,1-dioxo-1 λ^6 -[1,2]thiazinan-5-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

Morpholine-4-carboxylic acid [1-(5-cyano-2-methyl-1,1-dioxo-1 λ^6 -[1,2]thiazinan-5-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

- 20 Morpholine-4-carboxylic acid [1-(5-cyano-2-oxo-hexahydro-pyrimidin-5-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(5-cyano-1,3-dimethyl-2-oxo-hexahydro-pyrimidin-5-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

25

Morpholine-4-carboxylic acid [1-(4-cyano-1,1-dioxo-1 λ^6 -[1,2,6]thiadiazinan-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

- 30 Morpholine-4-carboxylic acid [1-(4-cyano-2,6-dimethyl-1,1-dioxo-1 λ^6 -[1,2,6]thiadiazinan-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

Morpholine-4-carboxylic acid [1-(3-cyano-5-oxo-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

5 Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-5-oxo-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

Morpholine-4-carboxylic acid [1-(3-cyano-5-oxo-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

10 Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-5-oxo-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

15 **EXAMPLE 2**

Morpholine-4-carboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

20 **(a) 4-Amino-4-cyano-tetrahydropyran.**

A solution of ammonium chloride (2.12 g, 39.57mmol) and potassium cyanide (2.58 g, 39.57 mmol) was prepared in 50 mL of water. Tetrahydropyran-4-one (1.0 g, 9.89 mmol) was added to the solution and stirring was continued for 2 days. The solution was brought to pH 11 with solid sodium carbonate and the reaction solution was extracted 3 x 100 mL of EtOAc. The organic layer was dried over anhydrous Na₂SO₄, decanted, and concentrated to an clear oil (1.02 g). ¹H NMR showed that the oil was a 7 to 1 mixture of the desired aminonitrile and cyanohydrin. The crude mixture was used in the next step without further purification.

30

(b) Morpholine-4-carboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

N-(4-morpholinecarbonyl)-L-cyclohexyl alanine (350 mg, 1.23 mmol) was dissolved in 15 mL of DMF. EDC (235 mg, 1.23 mmol) and HOBT (166 mg, 1.23 mmol) were added and the resulting mixture was stirred at ambient temperature for 20 min during which time the solids went into solution. 4-Amino-4-cyano-tetrahydropyran (161 mg of the 7:1 mixture of aminonitrile, \cong 1.1 mmol aminonitrile) was dissolved in 5 mL of DMF, N-methylmorpholine was added to this solution (497 mgs, 4.92 mmol), and the resultant solution added to the solution of the active ester. The resulting mixture was stirred at ambient temperature for 16 h. The volatiles were removed *in vacuo* and the resulting residue was vigorously stirred for 30 min with 100 mL of a 1 to 1 mixture of water and saturated sodium bicarbonate to give a fluffy white solid that was collected by filtration. The solid was washed with 3x50 mL of water and dried to give the desired product (210 mg): ^1H NMR (CDCl_3) δ 7.80 (s, 1H), 5.25-5.15 (m, 1H), 4.41-2.0 (m, 1H), 3.97-3.62 (m, 8H), 3.50-3.41 (m, 4H), 2.50-2.37 (m 1H), 2.35-2.20 (m, 1H), 2.05-1.88 (m, 2H), 1.79-0.75 (m, 13H).

EXAMPLE 3

4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester

(a) 4-Amino-4-cyano-piperidine-1-carboxylic acid ethyl ester.

A solution of ammonium chloride (31 g, 584 mmol) and potassium cyanide (7.61 g, 116.8 mmol) was prepared in 250 mL of water. 1-(Ethoxycarbonyl)-4-piperidone (10 g, 58.4 mmol) was added to the solution followed by 50 mL of MeOH and stirring was continued for 3 days. The solution was brought to pH 11 with solid sodium carbonate (20g) and the reaction solution was extracted with 3 x 250 mL of EtOAc. The organic layers were combined, dried over Na_2SO_4 , decanted and concentrated to an thick oil. The oil was triturated with 500 mL of hexane and the resulting solid was collected by filtration (8.3g). ^1H NMR showed that the product was better than 95% pure.

(b) **4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester**

N-(4-morpholinecarbonyl)-L-cyclohexyl alanine (555 mg, 1.95 mmol) was dissolved in 15 mL of DMF. EDC (373 mg, 1.95 mmol) and HOBt (264 mg, 1.95 mmol) were added and the resulting mixture was stirred at ambient temperature for 20 min during which time the solids went into solution. 4-Amino-4-cyano-piperidine-1-carboxylic acid ethyl ester (350 mg, 1.77 mmol) was dissolved in 5 mL of DMF and added to the solution of the active ester followed by addition of 2 mL of N-methylmorpholine. The resulting mixture was stirred at ambient temperature for 16 h. The volatiles were removed *in vacuo* and the resulting residue was dissolved in 200 mL of EtOAc and washed sequentially with 2x200 mL saturated sodium bicarbonate, 1x100 mL brine. The organic layer was dried over Na₂SO₄, decanted, and concentrated to a white solid. The solid was purified by column chromatography on SiO₂ using as eluent 100% CH₂Cl₂ to 5% MeOH in CH₂Cl₂ to give the title compound as a white powder (511 mg): m.p. 140-143 °C.

EXAMPLE 4

Morpholine-4-carboxylic acid [1-(4-cyano-1-phenethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

- (a) 4-Amino-4-cyano-1-phenethylpiperidine was prepared according to the procedure from Example 1, step a, starting with 1-phenylethyl-4-piperidone.
- (b) The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-1-phenethylpiperidine according to the procedure from Example 2, step b, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 496 = M+1.

EXAMPLE 5

Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

5

(a) 4-Amino-4-cyano-1-benzylpiperidine was prepared according to the procedure from Example 1, step a, starting with 1-benzyl-4-piperidone

(b) The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-1-benzylpiperidine according to the procedure from Example 2, step b, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 482 = M+1.

15

EXAMPLE 6

Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

(a) 4-Amino-4-cyano-1-propylpiperidine was prepared according to the procedure from Example 1, step a, starting with 1-propyl-4-piperidone.

(b) The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-1-propylpiperidine according to the procedure from Example 2, step b, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 434 = M+1.

30

EXAMPLE 7

4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid benzyl ester

(a) A solution of sodium cyanide (1052 mg, 21.5 mmol), ammonium chloride (1265 mg, 23.65 mmol), and benzyl 4-oxo-1-piperidine-carboxylate (5.0 gm, 21.5 mmol), was prepared in 5 M ammonia in methanol (8.6 mL, 43 mmol). The solution was brought to reflux for 4 h and then allowed to cool to room temperature. The solution was then filtered and washed with methanol (100 mL) and the filtrate was concentrated *in vacuo*. The resulting oil was taken up in MTBE (250 mL) and filtered again. The filter cake was washed with MTBE (100 mL) and the filtrate was concentrated *in vacuo* to yield 4-amino-4-cyano-piperidine-1-carboxylic acid benzyl ester as a clear oil (3.5 g) which was used without further purification.

(b) The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-piperidine-1-carboxylic acid benzyl ester according to the procedure from Example 2, step b, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 526 = M+1.

EXAMPLE 8

Morpholine-4-carboxylic acid [1-(4-cyano-tetrahydro-thiopyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

(a) 4-Amino-tetrahydro-thiopyran-4-carbonitrile was prepared according to the procedure from Example 7, step a, starting from tetrahydrothiopyran-4-one.

(b) The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-tetrahydrothiopyrane according to the procedure from Example 2, step b, except that the compound was purified by reverse phase HPLC

using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 409 = M+1.

EXAMPLE 9

5

Morpholine-4-carboxylic acid [1-(4-cyano-1-pyrimidin-2-yl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

(a) 4-Amino-4-cyano-1-pyrimidin-2-yl-piperidine was prepared according to the procedure from Example 7, step a, starting with 1-(pyrimidin-2-yl)-4-piperidone with the exception that a 2 M ammonia in methanol solution replaced the 5 M ammonia in methanol solution.

(b) The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-1-pyrimidin-2-yl-piperidine according to the procedure from Example 2, step b, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 469 = M+1.

20

EXAMPLE 10

Morpholine-4-carboxylic acid [1-(4-cyano-2,6-diphenyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

(a) 4-Amino-4-cyano-2,6-diphenyl-piperidine was prepared according to the procedure from Example 9, step a, starting from 2,6-diphenyl-4-piperidone.

(b) The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-2,6-diphenyl-piperidine according to the procedure from Example 2, step b, except that the compound was purified by HPLC

30

using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 544 = M+1.

EXAMPLE 11

5

Morpholine-4-carboxylic acid [1-(4-cyano-2,6-diphenyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

(a) 2-Amino-4,4-dimethyl-pentanoic acid methyl ester

10

2-Amino-4,4-dimethyl-pentanoic acid (1.00 g, 6.84 mmol) was suspended in 50 mL of methanol and cooled in an ice bath. Thionyl chloride (1.82 g, 15.0 mmol) was added dropwise, at which time all the acid went into solution. The reaction was then removed from the ice bath and heated to reflux for 3.5 h. The reaction mixture was concentrated *in vacuo* and the resulting solid (1.10 g) was used in the next step without further purification. MS, m/z 159.9 = M+1

15

(b) 4,4-Dimethyl-2-[(morpholine-4-carbonyl)-amino]-pentanoic acid methyl ester

20

2-Amino-4,4-dimethyl-pentanoic acid methyl ester (5.35 g, 27.4 mmol) was dissolved in 100 mL of dichloromethane. Hunig's base (7.07 g, 54.7 mmol) and 4-morpholinecarbonyl chloride (4.08 g, 27.4 mmol) were added and the reaction was stirred at ambient temperature 16 h. The reaction mixture was concentrated *in vacuo* and taken up in 150 mL EtOAc. A white precipitate formed and was filtered and washed with EtOAc. EtOAc solutions were combined and washed with 3 x 50 mL 1 N HCl (aq), 3 x 50 mL saturated NaHCO₃ (aq), and 1 x 50 mL brine. The organic layer was dried over Na₂SO₄, decanted, and concentrated to a white solid (6.33 g). MS, m/z 273 = M+1

25

(c) N-(4-morpholinecarbonyl)-L-neopentyl glycine

30

4,4-Dimethyl-2-[(morpholine-4-carbonyl)-amino]-pentanoic acid methyl ester (6.33 g, 23.2 mmol) was dissolved in 100 mL of THF and 50 mL of methanol. The solution was cooled on an ice bath and lithium hydroxide monohydrate (5.80 g, 116 mmol) was added as a suspension in 50 mL of water. The reaction was stirred at ambient temperature for 1 h. Additional water was added to the reaction (25 mL) and the mixture was extracted with diethyl ether 2 x 75 mL. The organic layer was discarded. The aqueous layer was acidified to pH 2 with 20% HCl (aq) and the product was extracted with 3 x 75 mL EtOAc. The EtOAc layer was washed with 1 x 50 mL brine and dried over Na₂SO₄, decanted, and concentrated *in vacuo* to a white solid (5.85 g). MS, m/z 259 = M+1

(d) Morpholine-4-carboxylic acid [1-(4-cyano-2,6-diphenyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

N-(4-morpholinecarbonyl)-L-neopentyl glycine (214 mg, 0.83 mmol) was dissolved in 25 mL of dichloromethane. EDC (175 mg, 0.91 mmol), HOBT (123 mg, 0.91 mmol), 4-amino-4-cyano-2,6-diphenylpiperidine (Example 10) (278 mg, 0.91 mmol), and N-methylmorpholine (420 mg, 4.2 mmol) were added to the solution. The reaction was stirred at ambient temperature for 16 h. The reaction was concentrated *in vacuo* and the resulting residue was dissolved in 150 mL of EtOAc. The EtOAc layer was washed with 2 x 50 mL saturated NaHCO₃, 1 x 50 mL brine, then dried over Na₂SO₄, decanted, and concentrated to an oil. Product was recrystallized from EtOAc/ hexanes to yield a white solid (42 mg). MS, m/z 518 = M+1

EXAMPLE 12

Morpholine-4-carboxylic acid [1-(1-acetyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

(a) 4-Amino-4-cyano-1-acetylpiperidine was prepared according to the procedure from Example 9, step a, starting with 1-acetyl-4-piperidone.

- (b) The title compound was prepared starting from N-(4-morpholinecarbonyl)-L-cyclohexyl alanine and 4-amino-4-cyano-1-acetylpiperidine according to the procedure from Example 2, step b, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 433 = M+1.

EXAMPLE 13

Morpholine-4-carboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

- (a) 4-Amino-4-cyano-tetrahydropyran was prepared according to the procedure from Example 1, step a, starting from tetrahydropyran-4-one.
- (b) The title compound was prepared starting from N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) and 4-amino-4-cyano-tetrahydropyrane according to the procedure from Example 2, step b, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 367 = M+1.

EXAMPLE 14

Morpholine-4-carboxylic acid [1-(4-cyano-tetrahydro-thiopyran-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

- The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) and 4-amino-4-cyano-tetrahydrothiopyran (Example 8) according to the procedure from Example 1, step d, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 383 = M+1.

EXAMPLE 15

Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

5

The title compound was prepared starting from N-(4-Morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) and 4-amino-4-cyano-1-benzylpiperidine (Example 5, step a) according to the procedure from Example 1, step d, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being

10 30% acetonitrile in water to 100% acetonitrile. MS, m/z 456 = M+1.

EXAMPLE 16

Morpholine-4-carboxylic acid [1-(1-isopropyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

15

(a) 4-Amino-4-cyano-1-isopropylpiperidine was prepared according to the procedure from Example 1, step a, starting from 1-*i*-propyl-4-piperidone.

20 (b) The title compound was prepared starting from N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) and 4-amino-4-cyano-1-isopropylpiperidine according to the procedure from Example 1, step d, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 456 = M+1.

25

EXAMPLE 17

Morpholine-4-carboxylic acid [1-(1-phenethyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

30

The title compound was prepared starting from N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) and 4-amino-4-cyano-1-phenethylpiperidine (Example 4) according to the procedure from Example 1, step d, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 470 = M+1.

EXAMPLE 18

10 **Morpholine-4-carboxylic acid [1-(1-n-propyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide**

The title compound was prepared starting from N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) and 4-amino-4-cyano-1-n-propylpiperidine (Example 6) according to the procedure from Example 1, step d, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 408 = M+1.

20 **4-Cyano-4-{3,3-dimethyl-2-[(morpholine-4-carbonyl)-amino]-pentanoylamino}-piperidine-1-carboxylic acid benzyl ester**

25 The title compound was prepared starting from N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) and 4-amino-4-cyano-piperidine-1-carboxylic acid benzyl ester (Example 7, step a) according to the procedure from Example 1, step d, except that the compound was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 30% acetonitrile in water to 100% acetonitrile. MS, m/z 500 = M+1.

30

EXAMPLE 20

Morpholine-4-carboxylic acid [1-(1-acetyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

5

(a) 1-Acetyl-4-amino-piperidin-4-carbonitrile

1-Acetyl-4-amino-piperidin-4-carbonitrile was prepared from N-acetyl-4-piperidone according to the procedure from Example 9, step a.

- 10 (b) The title compound was prepared starting from 1-Acetyl-4-amino-piperidine-4-carbonitrile and N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) according to the procedure from Example 11, step d and purified by reverse phase HPLC (43 mg). MS, m/z 408 = M+1.

15

EXAMPLE 21

Morpholine-4-carboxylic acid [1-(1-benzoyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

20

(a) 4-Amino-1-benzoyl-piperidine-4-carbonitrile was prepared from N-benzoyl-4-piperidone according to the procedure from Example 9, step a. MS, m/z 168 = M+1

- 25 (b) The title compound was prepared starting from 4-amino-1-benzoyl-piperidine-4-carbonitrile and N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) according to the procedure from Example 11, step d and purified by reverse phase HPLC (66 mg). MS, m/z 470 = M+1

30

EXAMPLE 22

4-Cyano-4-{4,4-dimethyl-2-[(morpholine-4-carbonyl)-amino]-pentanoylamino}-piperidine-1-carboxylic acid ethyl ester

5 (a) 4-Amino-4-cyano-piperidine-1-carboxylic acid ethyl ester was prepared according to the procedure from Example 1, step a, from 4-oxopiperidine-1-carboxylic acid ethyl ester.

(b) The title compound was prepared starting from 4-Amino-4-cyano-piperidine-1-carboxylic acid ethyl ester and N-(4-morpholinecarbonyl)-L-neopentyl glycine (Example 11, step c) according to the procedure from Example 11, step d and purified by reverse phase HPLC (67 mg). MS, m/z 438 = M+1

EXAMPLE 23

15

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-dimethylamino-acetyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

The title compound was prepared from N,N-dimethylaminoglycine and morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride using the coupling method described in Example 1-part (d). The product was purified by reverse phase preparative HPLC to give the title compound as an off-white solid; MS, m/z 477 = M+1.

20

EXAMPLE 24

25

4-Acetylamino-N-[1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide

(a) *t*-Butoxycarboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

30

t-Butoxycarboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide was prepared from N-Boc-L-cyclohexylalanine and 4-amino-4-cyano-tetrahydropyran by the method of Example 2-part (b). The product was used in the next step without further purification.

5

(b) [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amine hydrochloride.

t-Butoxycarboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide (1000 mg, 2.62 mmol) was dissolved in 15 mL of 4 M HCl in dioxane. The solution was stirred at ambient temperature for 1 hr. The volatiles were removed *in vacuo* and the resulting paste was triturated with 25 mL of diethyl ether to give a fine white solid that was collected by filtration and dried *in vacuo*. The product was used without further purification.

(c) 4-Acetylamino-N-[1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide.

4-Acetamidobenzoic acid (353 mg, 1.98 mmol), EDC (378 mg, 1.98 mmol), and HOBt (268 mg, 1.98 mmol) were combined in 15 mL of DMF and stirred for 20 min. Solid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amine hydrochloride (625 mg, 1.98 mmol) was added. The reaction was stirred for 16 hours. The volatiles were removed with a pump and the resulting residue was triturated, with rapid stirring, with 250 mL of saturated aqueous sodium bicarbonate. The resulting solid was collected by filtration and washed with 250 mL of water. The solid was dried *in vacuo* to give the title compound (250 mg); MS, m/z 441 = M+1.

25

Following the above procedures the following compounds can be synthesised;

4-Chloro-N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide

30

N-[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-methoxy-benzamide

N-[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-isonicotinamide

5 Pyrazine-2-carboxylic acid [1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

N-[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-3-phenoxy-benzamide

10 Furan-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

15 Thiophene-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

5-Chloro-thiophene-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

20 N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(2-thiophen-2-yl-acetylamino)-propionamide.

EXAMPLE 25

25 **Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide**

30 (a) ***t*-Butoxycarboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.**

t-Butoxycarboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide was prepared from N-Boc-L-neopentylglycine and 4-amino-4-cyano-1-methyl-piperidine by a method analogous to that of Example 2-part (b). The product was used without further
35 purification.

(b) **[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amine dihydrochloride.**

[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amine dihydrochloride was prepared by a method analogous to that of Example 24-part (b). The product was used without further purification.

5

(c) **Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.**

[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amine dihydrochloride (350 mg, 1.03 mmol) was mixed in 10 mL of DMF to which was added 1 mL of N-methyl morpholine followed by addition of 4-morpholine carbonyl chloride (180 mg, 1.20 mmol) as a solution in 5 mL of DMF. The reaction was stirred for 16 hours at which time the volatiles were removed *in vacuo*. The residue was redissolved in 150 mL of EtOAc and washed sequentially with 50 mL of saturated aqueous bicarbonate and 50 mL of brine. The organic layer was dried over sodium sulfate, decanted and concentrated. The product was purified by flash chromatography on silica gel using 100% methylene chloride to 12 % methanol in methylene chloride as eluent to give the title compound as a thick oil (85 mg); MS, m/z 380 = M+1.

15

Following the above procedures the following compounds can be synthesised;

20

4-Chloro-N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide

25

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methoxy-benzamide

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-isonicotinamide

30

Pyrazine-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Furan-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

5 Thiophene-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

4,4-Dimethyl-2-(2-thiophen-2-yl-acetylamino)-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

10 N-[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-3-phenoxy-benzamide

5-Chloro-thiophene-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.

15

EXAMPLE 26

4-Acetylamino-N-1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl)-benzamide

20

The title compound was prepared by a method analogous to that of Example 24; MS, m/z 454 = M+1.

EXAMPLE 27

25

4-Acetylamino-N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide

30 The title compound was prepared by a method analogous to that of Example 24; MS, m/z 428 = M+1.

EXAMPLE 28

4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid *t*-butyl ester

(a) 4-Amino-4-cyano-piperidine-1-carboxylic acid *t*-butyl ester.

4-Amino-4-cyano-piperidine-1-carboxylic acid *t*-butyl ester was prepared by a method analogous to that of Example 3-part (a). The product was used without further purification.

(b) 4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid *t*-butyl ester.

4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid *t*-butyl ester was prepared by a method analogous to that of Example 3-part (b); MS, *m/z* 391, M- *t*-butoxycarbonyl).

EXAMPLE 29

Morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride

4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid *t*-butyl ester (1000 mg, 2.03 mmol) was dissolved in 20 mL of 4 M HCl in dioxane and stirred for 1 hour at which time the volatiles were removed in vacuo. The resulting residue was triturated with 100 mL of diethyl ether and the resulting solid was collected by filtration under inert atmosphere (the solid is very hygroscopic) and washed 2x50 mL of diethyl ether and dried in vacuo to yield the title compound as a bright white powder (802 mg); MS, *m/z* 392, M-35).

EXAMPLE 30

Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

5 **(a) 4-Amino-4-cyano-1-(1-methyl-ethyl)-piperidine.**

4-Amino-4-cyano-1-(1-methyl-ethyl)-piperidine was prepared by a method analogous to that of Example 1-part (a). The product was used without further purification.

10 **(b) Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide.**

Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide was prepared by a method analogous to that of Example 1-part (d); MS,

15 m/z 434 = M+1.

EXAMPLE 31

Morpholine-4-carboxylic acid {1-[3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

20

(a) 3-Amino-3-cyano-1-benzylpyrrolidine.

3-Amino-3-cyano-1-benzylpyrrolidine was prepared by a method analogous to that of Example 1-part (a) with the exception that no sodium carbonate was added to the reaction mixture. The product was extracted from the crude reaction with 3x100 mL of EtOAc and was used without purification.

25

(b) Morpholine-4-carboxylic acid {1-[3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide. Separated diastereomers.

30

Diastereomeric morpholine-4-carboxylic acid {1-[3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide was prepared by a method analogous to that of Example 1-part (d). The purification was done by reverse phase preparative HPLC (Hypersil HyPURITY™, C18 column, 250 x 21.2 5μ) to separate the two diastereomers; MS, m/z 468 = M+1.

5

EXAMPLE 32

Morpholine-4-carboxylic acid [1-(4-cyano-2,6-dimethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

10

(a) Cis-2,6-dimethyl-4-piperidone.

Into a mixture of dimethyl acetonedicarboxylate (10 g, 57.4 mmol) and acetaldehyde (4.4 g, 100 mmol) maintained at -25 °C was bubbled ammonia until the solution was saturated (careful bubbling required due to exothermic dissolution of NH₃). The resulting solution was stored at 0 °C for 20 hours, by which time it was a white sludge. To this was added 25 mL of 3 N hydrochloric acid and the solution was heated on the steam-bath. Carbon dioxide began to evolve soon, but after 24 hours was still evolving very slowly. The solution was evaporated almost to dryness. To the tan heavy precipitate was added 25 mL of water and the solution was again evaporated. To the residue was added a solution of 10 g of sodium carbonate in 45 mL of water and 20 mL of chloroform. The layers were shaken and separated. The water layer was extracted six times with 20 mL portions of methylene chloride. The organic layers were dried over magnesium sulfate and concentrated to give the desired crude product which was used without further purification.

25

(b) 4-Amino-4-cyano-2,6-dimethyl-piperidine.

To a mixture of ammonium chloride (0.58 g, 9.98 mmol), sodium cyanide (0.50 g, 11.0 mmol), ammonium hydroxide (2 mL) was added a solution of cis-2,6-dimethylpiperidone (1.27 g, 9.98 mmol) in 5 mL of methanol. The resulting mixture was refluxed for 4 hours. The reaction mixture was evaporated to dryness and the residue was taken up in 50 mL EtOAc, washed with

30

saturated sodium bicarbonate 3 x 50 mL. The organic layer was evaporated to dryness and purified by flash chromatography on silica gel using 90 to 9 methylene chloride and methanol to give the desired product.

- 5 (c) **Morpholine-4-carboxylic acid [1-(4-cyano-2,6-dimethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.**

The title compound was prepared by the standard method of Example 1-part (d); MS, m/z 420 = M+1.

10

EXAMPLE 33

Morpholine-4-carboxylic acid [1-(4-cyano-1,3-dimethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

15

- (a) **1,3-Dimethyl-4-piperidone hydrochloride.**

To a solution of methylamine (100 mL of 2.0 M solution in methanol) was added, over the course of 1 hour at 0 °C, a solution of methyl methacrylate (30.2 g, 300 mmol) in 20 mL of
20 methanol. The resulting solution was allowed to stand for three days, at which time the volatiles were removed on a rotovap and the residue was vacuum distilled to give the desired product as a clear oil, b.p. 48-49 C at 8.5 mm. The oil was dissolved in 100 mL of methanol and methyl acrylate (14.8 g, 200 mmol) was added and the reaction was allowed to stand for 3 days. The volatiles were removed.

25

30 mL of Xylene was prepared over sodium (2.42 g) and refluxed for 2 hours and cooled to 60 °C. To this mixture was added the diester and the reaction was refluxed until the sodium particles had disappeared. The resulting dark red liquid was cooled and poured into 150 mL of ice water. The phases were separated and the xylene extracted with 50 mL of concentrated
30 hydrochloric acid and, after washing with 50 mL of isopropyl ether, the aqueous layer was cooled, basified with potassium carbonate and extracted eight times with 75 mL portions of ethyl

ether. The combined ethereal extracts were dried over potassium carbonate and treated with excess dry ethereal hydrogen chloride; the resulting salt was filtered and dried.

The salt was taken up in 60 mL of 6 N hydrochloric acid and heated on a water bath for three hours, at the end of which time the initially vigorous carbon dioxide evolution had become negligible. The resulting solution was evaporated to dryness and dried *in vacuo*, to yield 1,3-dimethyl-4-piperidone hydrochloride (5g) which was used in the next step without further purification.

10 (b) **4-Amino-4-cyano-1,3-dimethylpiperidine.**

The title compound was prepared as described in the previous example for 4-amino-4-cyano-2,6-dimethylpiperidine. The crude product was used in the next step without further purification.

15 (c) **Morpholine-4-carboxylic acid [1-(4-cyano-1,3-dimethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.**

The title compound was prepared as in Example 1-part (d); MS, m/z 420 = M+1.

20 **EXAMPLE 34**

4-Cyano-4-{3-cyclohexyl-2-[(4-acetylamino)-phenyl-1-carbonyl]-aminol-propionylamino}-piperidine-1-carboxylic acid ethyl ester

25 The title compound was prepared by a method analogous to that of Example 24; MS, m/z 512 = M+1.

EXAMPLE 35

30 **4-Acetylamino-N-[1-(4-cyano-1-benzyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide**

The title compound was prepared by a method analogous to that of Example 24; MS, m/z 530 = M+1.

5

EXAMPLE 36

4-Acetyl-amino-N-{1-[4-cyano-1-(1-methyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-benzamide

10 The title compound was prepared by the method of Example 24; MS, m/z 482 = M+1.

EXAMPLE 37

15

Morpholine-4-carboxylic acid {1-[3-cyano-1-benzyl-piperidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

The title compound, separated into two diastereomers, was prepared by a method analogous to that of Example 31; MS, m/z 482 = M+1.

20

EXAMPLE 38

4-Cyano-4-{3-cyclohexyl-2-[(4-acetyl-amino)-phenyl-1-carbonyl]-aminol-propionyl-amino}-piperidine-1-carboxylic acid benzyl ester

25 The title compound was prepared by a method analogous to that of Example 24; MS, m/z 574 = M+1.

EXAMPLE 39

30 N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide

The title compound was prepared by a method analogous to that of Example 24; MS, m/z 397 = $M+1$.

EXAMPLE 40

5

4-Acetylamino-N-{1-[4-cyano-1-(2-phenyl-ethyl)-piperidin-4-yl]carbamoyl}-2-cyclohexyl-ethyl}-benzamide

The title compound was prepared by a method analogous to that of Example 24; MS, m/z 544 = $M+1$.

10

EXAMPLE 41

15

4-(Acetylamino-methyl)-N-[1-(4-cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl}-benzamide

(a) 4-(Acetylamino-methyl)-benzoic acid.

Methyl-4-(acetylamino-methyl)-benzoate was prepared from acetic acid and methyl 4-(aminomethyl)benzoate using a method analogous to that of Example 1-part (d). The crude N-acyl ester was saponified using a method analogous to that of Example 1-part (c). The crude product was used without further purification.

20

(b) 4-(Acetylamino-methyl)-N-[1-(4-cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl}-benzamide.

25

The title compound was prepared by a method analogous to that of Example 24-part (c); MS, m/z 468 = $M+1$.

30

EXAMPLE 42

Morpholine-4-carboxylic acid [1-(3-cyano-8-methyl-8-aza-bicyclo[3.2.1]oct-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

(a) 3-Amino-3-cyano-8-methyl-8-aza-bicyclo[3.2.1]octane.

5

The aminonitrile was prepared from tropinone using a method analogous to that of Example 1-part (a).

(b) Morpholine-4-carboxylic acid [1-(3-cyano-8-methyl-8-aza-bicyclo[3.2.1]oct-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

10

The title compound was prepared using a method analogous to that of Example 1-part (d); MS, m/z 432 = M+1.

15

EXAMPLE 43

Morpholine-4-carboxylic acid [1-(1-carbamimidoyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide *p*-toluenesulfonate

(a) 1-Carbamimidoyl-1,2,3-benztriazole *p*-toluenesulfonate.

20

A mixture of benztriazole (11.9 g, 100 mmol), cyanamide (4.2 g, 100 mmol), and *p*-toluene sulfonic acid hydrate (19.2 g, 100 mmol) in dioxane was refluxed for 24 hours. The reaction mixture was allowed to cool to room temperature and was diluted with ether, stirred vigorously, then filtered. The filter cake was washed with ether and recrystallized from ethanol to give the desired product as a white solid.

25

(b) Morpholine-4-carboxylic acid [1-(1-carbamimidoyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide *p*-toluenesulfonate.

30

Morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride (0.2 g, 0.47 mmol) was dissolved in 3 mL of DMF and 2 equiv of Hunig's base was added followed by 1-carbamimidoyl-1,2,3-benzotriazole *p*-toluenesulfonate (0.16 g, 0.47 mmol). The reaction was stirred 24 hours at which time the solvent was removed in vacuo. The resulting paste was purified by preparative HPLC to give the title compound; MS, *m/z* 434, M+1- *p*-toluene sulfonate).

EXAMPLE 44

10 **4-Acetylamino-N-[1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide**

4-Amino-4-cyano-tetrahydropyran prepared according to the procedure from Example 1, step a, starting from tetrahydropyran-4-one.

15

The title compound was prepared from 4-amino-4-cyano-tetrahydropyran, L-neopenyyl glycine and 4-acetylamino benzoic acid analogous to the procedure described in Example 24.

20

EXAMPLE 45

Morpholine-4-carboxylic acid [1-(4-cyano-1-methanesulfonyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

25 The title compound is prepared by treatment of morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride with methanesulfonyl chloride and a tertiary amine base such as *N*-methylmorpholine in a solvent such as methylene chloride.

30

EXAMPLE 46

4-Acetylamino-piperidine-1-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

The title compound is prepared by a method analogous to that of Example 24.

5

EXAMPLE 47

Morpholine-4-carboxylic acid {1-[1-(2-chloro-benzyl)-3-cyano-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide.

10

The title compound is prepared by a method analogous to that of Example 31.

EXAMPLE 48

15 **Morpholine-4-carboxylic acid [1-(1-benzylcarbamoyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.**

The title compound is prepared from morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide hydrochloride and benzyl isocyanate in the presence of
20 a tertiary amine base such as *N*-methylmorpholine in a solvent such as methylene chloride.

EXAMPLE 49

25 **Morpholine-4-carboxylic acid [1-(1-phenylcarbamoyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.**

The title compound is prepared from morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide hydrochloride and phenyl isocyanate in the presence of
30 a tertiary amine base such as *N*-methylmorpholine in a solvent such as methylene chloride.

EXAMPLE 50

Morpholine-4-carboxylic acid {1-[4-cyano-1-(morpholine-4-carbonyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl]-amide.

- 5 The title compound is prepared from morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide hydrochloride and 4-morpholine carbonyl chloride in the presence of a tertiary amine base such as *N*-methylmorpholine in a solvent such as methylene chloride.

10 EXAMPLE 51

Morpholine-4-carboxylic acid (1-{4-cyano-1-[(pyridin-3-ylmethyl)-carbamoyl]-piperidin-4-ylcarbamoyl}-3,3-dimethyl-butyl)-amide.

- 15 The title compound is prepared from morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide hydrochloride and 3-pyridyl-methyl isocyanate in the presence of a tertiary amine base such as *N*-methylmorpholine in a solvent such as methylene chloride.

20 EXAMPLE 52

Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-methyl-piperazine-1-carbonyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl]-amide.

- 25 The title compound is prepared from morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride and 4-methyl-piperzine carbonyl chloride in the presence of a tertiary amine base such as *N*-methylmorpholine in a solvent such as methylene chloride.

30 EXAMPLE 53

4-(4-Cyano-4-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-piperidin-1-yl)-butyric acid.

The title compound is prepared from morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride and 4-bromo-butyric acid in the presence of a hindered tertiary amine base such as Hunig's base in a solvent such as methylene chloride.

EXAMPLE 54

Morpholine-4-carboxylic acid [1-(4-cyano-1-cyclopropyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

The title compound may be prepared from morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride and 1-ethoxy-1-trimethylsilyloxy-cyclopropane using a reducing agent such sodium cyanoborohydride in a solvent system such as acetic acid in methanol.

EXAMPLE 55

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-dimethylamino-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide.

The title compound is prepared by the method of Example 33.

EXAMPLE 56

Morpholine-4-carboxylic acid [1-(4-cyano-1-phenyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

The title compound is prepared by a method analogous to that of Example 58.

EXAMPLE 57

5 Morpholine-4-carboxylic acid {1-[4-cyano-1-(1,1-dimethyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide.


The title compound can be prepared by a method analogous to that of the method of Example 59.

EXAMPLE 58

10

Morpholine-4-carboxylic acid [1-(4-cyano-1-phenyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.

(a) *N*-Phenyl-4-piperidone.




15

1,4-Dioxo-8-azaspiro[4.5]-decane (2.0 g, 14.0 mmol, 1.0 equiv), Pd₂(DBA)₃ (0.31 g, 0.34 mmol, 0.024 equiv), BINAP (0.64 g, 1.0 mmol, 0.073 equiv), NaO-*t*-Bu (3.9g, 41 mmol, 3.0 equiv) and bromobenzene (2.6 g, 17.7 mmol, 1.3 equiv) were combined under Ar in 50 mL of dry toluene. The resulting mixture was refluxed under Ar for 4 h. The reaction
20 mix was cooled and poured into 250 mL of saturated sodium bicarbonate solution. The product was extracted with 3 x 100 mL CH₂Cl₂. The organic extracts were combined and concentrated. The product was purified by flash chromatography on SiO₂ using 50% hexanes in CH₂Cl₂ to pure CH₂Cl₂ to give the *N*-phenyl ketal (2.9 g). The purified ketal was dissolved in mixture of 50 mL 1,4-dioxane, 50 mL water, and 20 mL concentrated
25 HCl. The mixture was refluxed for 3 h at which time mass spectrometry showed disappearance of the starting ketal. The cooled mixture was carefully poured into 600

mL of saturated sodium bicarbonate solution and the product extracted with 3 x 200 mL EtOAc. The combined organic extracts were combined and dried over Na₂SO₄, decanted and concentrated to a red oil (2.3 g) which was used without further purification; MS, m/z 176=M+1.

5


(b) 4-Amino-4-cyano-1-phenyl-piperidine.



N-Phenyl-4-piperidone (2.3 g, 13 mmol, 1.0 equiv) was dissolved in 26 mL of 2 M NH₃ in MeOH and NaCN (0.76 g, 15 mmol, 1.2 equiv) and NH₄Cl (0.80 g, 15 mmol, 1.2 equiv) were added and the mixture was refluxed for 2 h at which time an additional 26 mL of 2 M NH₃/MeOH was added followed by another 2 h of reflux. The reaction mixture was cooled and filtered. The filtrate was concentrated. The crude product was purified by flash chromatography on SiO₂ eluting with 100% CH₂Cl₂ and 2% MeOH in CH₂Cl₂ to give the pure product (1.92 g) as a thick yellow oil which solidified on standing; MS, m/z 202=M+1.

15

(c) Morpholine-4-carboxylic acid [1-(4-cyano-1-phenyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.



5 *N*-(4-morpholinecarbonyl)-L-neopenylglycine (0.2 g, 0.97 mmol, 1.0 equiv) and EDC (0.19 g, 0.97 mmol, 1.0 equiv) were combined in 10 mL of CH₂Cl₂ and stirred for 15 min at room temperature. A solution of 4-amino-4-cyano-1-phenyl-piperidine (0.20 g, 0.97 mmol, 1.0 equiv) in 5 mL of CH₂Cl₂ and *N*-methyl-morpholine (0.31 g, 3.1 mmol, 4.0 equiv) were added and stirring was continued for 16 h. The reaction was concentrated *in vacuo* and the residue was triturated with 100 mL saturated sodium bicarbonate solution with rapid stirring for 2 h. The resulting solid was collected by filtration and recrystallized from CH₃CN and water (2 to 1) to yield the title compound as an off-white solid (165 mg, 39%); MS, *m/z* 443=M+1.

Following the above procedures the following compounds can be synthesized:

15 Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-methoxy-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-methoxy-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

20 Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-methoxy-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-methyl-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-methyl-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

- 5 Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-methyl-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-phenyl-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

10

Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-phenyl-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

15

Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-phenyl-phenyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-methoxy-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

20

Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-methoxy-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-methoxy-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

25

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-methyl-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

30

Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-methyl-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-methyl-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

5 Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-phenyl-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-phenyl-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide


10 Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-phenyl-phenyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

EXAMPLE 59

15

Morpholine-4-carboxylic acid [1-(1-*tert*-butyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.


(a) *N*-Methoxy-*N*-methyl-acrylamide.



20


Acrolyl chloride (20 g, 221 mmol, 1.0 equiv) was dissolved in 500 mL of CH₂Cl₂ and cooled to 0 °C. Solid *N,O*-dimethyl-hydroxylamine hydrochloride (21.5 g, 221 mmol, 1.0 equiv) was added all at once. Et₃N was added dropwise over a 2 h period to give a thick yellow mixture. The reaction was stirred for an additional hour during which time
25 it was allowed to warm to room temperature. The mixture was poured into 1 L of water. Layers were separated and the organic layer was washed with 1 x 500 mL water, 1 x 500 mL brine and dried over Na₂SO₄. The solution was decanted and concentrated *in vacuo* to give the desired product as a yellow oil (23 g, 90%) which was used without further purification.

(b) 3-(*t*-Butyl-amino)-*N*-methoxy-*N*-methyl-propanamide.




N-Methoxy-*N*-methyl-acrylamide (5 g, 43.4 mmol, 1.0 equiv) was dissolved in *t*-butylamine (3.36 g, 46 mmol, 1.06 equiv). The resulting solution was stirred at room temperature for 48 h. The excess primary amine was removed *in vacuo* and the crude product was purified by flash chromatography on silica using 100% CH₂Cl₂ to 2% MeOH in CH₂Cl₂ to give the desired product as a light yellow oil (5.7g, 70%); MS, *m/z* 189=M+1.

(c) 1-*t*-Butyl-4-piperidone.




3-(*t*-Butyl-amino)-*N*-methoxy-*N*-methyl-propanamide (5 g, 26.6 mmol, 1.0 equiv) was dissolved in dry THF (50 mL) under Ar. The solution was cooled to -78 °C and a 1 M solution of vinylmagnesium bromide (66.5 mL, 66.5 mmol, 2.5 equiv) was added dropwise over a 20 min period. The reaction was then stirred at -78 °C for 30 min and at 0 °C for 30 min at which time the reaction solution was transferred via a double-ended cannula into ice-cold saturated sodium bicarbonate solution under Ar. The mixture was stirred for 10 min and the crude product was extracted 2 x 150 mL EtOAc. The organic extracts were combined and concentrated *in vacuo* to a red oil. Purification was done by flash chromatography on silica using 100% CH₂Cl₂ through 4, 8, and 16% MeOH in CH₂Cl₂. The product was isolated as an orange oil (1.3 g, 32%); MS, *m/z* 156=M+1.

(d) 4-Amino-1-*t*-butyl-4-cyano-piperidine.



1-*t*-Butyl-4-piperidone (1.3 g, 8.4 mmol, 1.0 equiv), NaCN (0.61 g, 12.6 mmol, 1.5 equiv), and NH₄Cl (0.67 g, 12.6 mmol, 1.5 equiv) were combined in 34 mL of 2 M NH₃ in MeOH. The mixture was refluxed for 2 h at which time an additional 34 mL of 2 M NH₃ in MeOH was added followed by another 2 h at reflux. The mixture was cooled and filtered. The filtrate was concentrated *in vacuo* and the residue was triturated with CH₂Cl₂ and filtered again. The solution was concentrated to a thick red oil which was used without further purification; MS, *m/z* 182=*M*+1.

(e) Morpholine-4-carboxylic acid [1-(1-*tert*-butyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.



N-(4-morpholinecarbonyl)-*L*-neopenylglycine (0.070 g, 0.27 mmol, 1.0 equiv) and EDC (0.057 g, 0.30 mmol, 1.1 equiv) were combined in 10 mL of DMF and stirred for 15 min at room temperature. A solution of 4-amino-1-*t*-butyl-4-cyano-piperidine (0.054 g, 0.30 mmol, 1.1 equiv) in 5 mL of DMF and *N*-methyl-morpholine (0.11 g, 1.1 mmol, 4.0 equiv) were added and stirring was continued for 16 h. The reaction was diluted with 50 mL of saturated sodium bicarbonate solution and the product was extracted with 3 x 50 mL EtOAc. The organic extracts were combined and concentrated *in vacuo*. The

product was purified by semi-prep reverse-phase HPLC using 20 to 60% CH₃CN in water over a 25 min gradient to yield the title compound as a white solid after concentration (25 mg, 22%); MS, m/z 422=M+1.


5

EXAMPLE 60

Morpholine-4-carboxylic acid [1-(4-cyano-1,2-dimethyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.

10


(a) 3-(Benzyl-methyl-amino)-butyric acid methyl ester.



Benzyl-methyl-amine (20 g, 165 mmol, 1.0 equiv) was added neat to methyl crotonate (19.8 g, 198 mmol, 1.2 equiv). The resulting solution was stirred at room temperature for 72 h. The excess crotonic ester was removed *in vacuo* to yield the desired product (40.3 g, ~100%) which was used without further purification; MS, m/z 222=M+1.

15

(b) 3-Methylamino-butyric acid methyl ester.




3-(Benzyl-methyl-amino)-butyric acid methyl ester (15 g, 67.8 mmol, 1.0 equiv) was placed in a Parr hydrogenation bottle and dissolved in 50 mL of MeOH. 20% Palladium hydroxide on carbon (0.5 g, 0.94 mmol, 0.014 equiv) was added and the mixture was shaken at 50 psi H₂ for 16 h. The reaction was judged as complete when the uptake of H₂ had stopped. The bottle was opened and 10 g of diatomaceous earth in 100 mL of MeOH was added. The mixture was filtered on a pad of diatomaceous earth which was then

25

washed with 2 x 100 mL of MeOH. The filtrates were combined and concentrated *in vacuo* to yield the desired product as an oil that is somewhat volatile (7.6 g, 85%). The crude product was used without further purification; MS, m/z 132=M+1.


5 (c) 3-[(2-Methoxycarbonyl-ethyl)-methyl-amino]-butyric acid methyl ester.



3-Methylamino-butyric acid methyl ester (7.6g, 58 mmol, 1.0 equiv) was added neat to methyl acrylate (7.5 g, 87 mmol, 1.5 equiv). The resulting solution was refluxed for 16 h. The reaction was cooled and diluted with hexanes (200 mL) and an insoluble polymer
10 separated out. The hexane solution was decanted and the polymer washed 2 x 100 mL hexanes with vigorous stirring. The combined hexane solutions were then concentrated *in vacuo*. The crude product was purified by flash chromatography on SiO₂ using pure CH₂Cl₂ as an eluent. The pure product was isolated as a clear colorless oil (7.3g, 58%); MS, m/z 218=M+1.

15


(d) 1,2-Dimethyl-4-piperidone.



A 1 M solution of TiCl₄ in CH₂Cl₂ (23 mL, 23 mmol, 1.0 equiv) was added to a flask under Ar and cooled to -15 °C with a MeOH/ice water bath. 3-[(2-Methoxycarbonyl-ethyl)-methyl-amino]-butyric acid methyl ester (5 g, 23 mmol, 1.0 equiv) was added
20 dropwise over a 25 min period as a solution in 75 mL of dry CH₂Cl₂ to give a dark red mixture that was difficult to stir with a magnetic stir bar. Stirring was continued an additional 1 h and then Et₃N (5.1 g, 50.6 mmol, 2.2 equiv) was added dropwise over a 30 min period and then the reaction was stirred an additional 1.5 h at -15 °C. The reaction


mix was poured into 150 mL of brine and 150 mL of CH_2Cl_2 was added. After thorough mixing, the pH of the water was brought to 8-9 with Et_3N . The mix was filtered and the gel-like solid was washed 3 x 100 mL CH_2Cl_2 . The filtrate layers were separated and the aqueous layer was washed 3 x 50 mL CH_2Cl_2 . All of the organic layers were combined and concentrated to a thick red oil. The residue was taken up in 150 mL of concentrated HCl and the solution was refluxed 4 h. The cooled reaction solution was evaporated to dryness and the residue was dissolved in 200 mL of saturated sodium bicarbonate solution. The product ketone was extracted with 2 x 100 mL of EtOAc. The organic layers were combined and dried over Na_2SO_4 . The product was purified by flash chromatography on SiO_2 using pure CH_2Cl_2 to 4% MeOH in CH_2Cl_2 as eluent. The product was isolated as an orange oil (1.23g, 42%); MS, m/z 128=M+1.

(e) **4-Amino-4-cyano-1,2-dimethyl-piperidine.**



1,2-Dimethyl-4-piperidone (1.23g, 9.67 mmol, 1.0 equiv) was dissolved in 39 mL of 2 M NH_3 in MeOH (8 equiv NH_3). To this solution was added NaCN (0.52g, 10.6 mmol, 1.1 equiv) and NH_4Cl (0.57g, 10.6 mmol, 1.1 equiv). The resulting mixture was refluxed for 2 h at which time an additional 39 mL of 2 M NH_3 in MeOH was added followed by an additional 2 h of reflux. The reaction was cooled and filtered. The filtrate was concentrated and taken up in 100 mL of CH_2Cl_2 giving more salt precipitate which was removed by a second filtration. The filtrate was then concentrated to thick orange oil (1.32g, 89%). ^1H NMR showed a 3 to 1 mixture of diastereomers of unknown configuration. The crude product was used without further purification; MS, m/z 154=M+1.

(f) **Morpholine-4-carboxylic acid [1-(4-cyano-1,2-dimethyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.**



N-(4-morpholinecarbonyl)-L-neopenylglycine (0.20g, 0.77 mmol, 1.0 equiv) and EDC (0.15g, 0.77 mmol, 1.0 equiv) were combined in 10 mL of DMF and stirred for 15 min at room temperature. A solution of 4-amino-4-cyano-1,2-dimethyl-piperidine (0.11g, 0.74
 5 mmol, 0.95 equiv) in 5 mL of DMF and *N*-methyl-morpholine (0.31g, 3.1 mmol, 4.0 equiv) were added and stirring was continued for 16 h. The reaction was diluted with 50 mL of saturated sodium bicarbonate solution and the product was extracted with 3 x 50 mL of EtOAc. The organic layers were combined and concentrated. The crude product was purified by semi-prep reverse-phase HPLC using 20% CH₃CN in water to 60%
 10 CH₃CN in water over a gradient of 16 min to give two peaks (diastereomers) eluting at 13.1 and 14.0 min (49mg and 20mg respectively); MS, *m/z* 394=*M*+1 for each peak.

Following the above procedures the following compounds can be synthesised:

- 15 Morpholine-4-carboxylic acid [1-(4-cyano-2-methyl-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-2-ethyl-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

20

Morpholine-4-carboxylic acid [1-(4-cyano-1,2-dipropyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

- Morpholine-4-carboxylic acid [1-(2-butyl-4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide
 25

Morpholine-4-carboxylic acid [1-(2-benzyl-4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

- 5 Morpholine-4-carboxylic acid [1-(4-cyano-2-cyclohexyl-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-2-cyclohexylmethyl-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

10

Morpholine-4-carboxylic acid [1-(4-cyano-2-methyl-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-2-ethyl-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

15

Morpholine-4-carboxylic acid [1-(4-cyano-1,2-dipropyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

- 20 Morpholine-4-carboxylic acid [1-(2-butyl-4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

Morpholine-4-carboxylic acid [1-(2-benzyl-4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

25

Morpholine-4-carboxylic acid [1-(4-cyano-2-cyclohexyl-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-2-cyclohexylmethyl-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide


30

EXAMPLE 61

Morpholine-4-carboxylic acid [1-(4-cyano-1-cyclohexyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.


5

(a) 4-Amino-4-cyano-piperidine-1-carboxylic acid *t*-butyl ester.



t-Butyl 4-oxo-1-piperidine-carboxylate (10 g, 50 mmol, 1.0 equiv) was dissolved 100 mL of 2 M NH₃ in MeOH. NaCN (2.7 g, 55 mmol, 1.1 equiv) and NH₄Cl (3 g, 55 mmol, 1.1 equiv) were added and the resulting mixture was refluxed for 2 h at which time an additional 100 mL of 2 M NH₃ in MeOH was added followed by another 2 h of reflux. The reaction mixture was cooled and filtered. The MeOH was removed in vacuo and the residue triturated with 100 mL of CH₂Cl₂ and filtered again. The filtrate was concentrated by about 75% and 200 mL of hexanes was added to give a tan precipitate that was collected by filtration to yield, after drying under vacuum, the desired product as cream-colored solid (10.1 g) which was used without further purification.


(b) 4-Cyano-4-{3,3-dimethyl-2-[(morpholine-4-carbonyl)-amino]-pentanoylamino}-piperidine-1-carboxylic acid *t*-butyl ester.



20


N-(4-morpholinecarbonyl)-*L*-neopenylglycine (1.00 g, 3.87 mmol, 1.00 equiv) and EDC (0.739 g, 3.87 mmol, 1.00 equiv) were mixed in 20 mL of CH₂Cl₂ and stirred for 15 min. A solution of 4-amino-4-cyano-piperidine-1-carboxylic acid *t*-butyl ester (0.872 g, 3.87 mmol, 1.00 equiv) in 10 mL CH₂Cl₂ and *N*-methyl-morpholine (1.56g, 15.5 mmol, 4.0 equiv) were added and the resulting solution was stirred at room temperature for 16 h. The reaction was diluted with 100 mL CH₂Cl₂ and 100 mL saturated sodium bicarbonate solution. The layers were separated and the aqueous was washed 2 x 50 mL CH₂Cl₂. The organic extracts were combined and dried over Na₂SO₄. The solution was decanted and concentrated to a white solid. The solid was dissolved in 20 mL of CH₃CN and water (100 mL) was added to precipitate the product. The fluffy white solid was collected by filtration and dried under high vacuum to yield the desired compound as a white powder (1.51 g).

(c) Morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide hydrochloride.



4-Cyano-4-{3,3-dimethyl-2-[(morpholine-4-carbonyl)-amino]-pentanoylamino}-piperidine-1-carboxylic acid *t*-butyl ester (1.51g, 3.24 mmol, 1.0 equiv) was dissolved in 50 mL of dry Et₂O under Ar. 4 M HCl in 1,4-dioxane (16 mL, 20 equiv) was added and the mixture was stirred for 20 min. A white solid precipitated almost immediately upon addition of the acid. The mixture was filtered under Ar and the solid was washed 2 x 25 mL of dry Et₂O. The solid was dried under high vacuum to a bright white powder (1.25g, 96%) which was used without further purification.

(d) Morpholine-4-carboxylic acid [1-(4-cyano-1-cyclohexyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide.



Morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide hydrochloride (0.050 g, 0.12 mmol, 1.0 equiv), cyclohexanone (0.015 g, 0.15 mmol, 1.2 equiv), and Na(OAc)₃BH (0.046 g, 0.22 mmol, 1.75 equiv) were mixed in 15 mL of 1% AcOH in THF. The reaction was stirred at room temperature for 16 h. The reaction was diluted with 25 mL of saturated sodium bicarbonate solution and the product was extracted 4 x 25 mL EtOAc. The organic extracts were combined and concentrated. The crude product was purified by semi-prep reverse-phase HPLC using 20 to 80% CH₃CN in water over a gradient of 25 min to yield the desired product, pure, as a white solid (0.012g, 21%); MS, m/z 448=M+1.

Following the above procedures the following compound was synthesised:

Morpholine-4-carboxylic acid {1-[4-cyano-1-(tetrahydro-pyran-4-yl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide; MS, m/z 450=M+1

Following the above procedures the following compounds can be synthesized;

Morpholine-4-carboxylic acid [1-(1-butyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-1-pentyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-1-hexyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

5 Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-ethyl-propyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-methyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

10 Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-methyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-methyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

15 Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-phenyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

20 Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-phenyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-phenyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

25 Morpholine-4-carboxylic acid {1-[4-cyano-1-(cyclohexyl-methyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

30 Morpholine-4-carboxylic acid {1-[4-cyano-1-(cyclopropyl-methyl)-piperidin-4-ylcarbamoyl]-3,3-dimethyl-butyl}-amide

Morpholine-4-carboxylic acid [1-(1-butyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

5 Morpholine-4-carboxylic acid [1-(4-cyano-1-pentyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

Morpholine-4-carboxylic acid [1-(4-cyano-1-hexyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide

10 Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-ethyl-propyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-methyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

15 Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-methyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

20 Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-methyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(2-phenyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

25 Morpholine-4-carboxylic acid {1-[4-cyano-1-(3-phenyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(cyclopropyl-methyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

30


Morpholine-4-carboxylic acid {1-[4-cyano-1-(4-phenyl-cyclohexyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

Morpholine-4-carboxylic acid {1-[4-cyano-1-(cyclohexyl-methyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide

EXAMPLE 62


Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopropylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

(a) 1-Cyclopropylmethyl-3-hydroxy-pyrrolidine.




3-Hydroxy-pyrrolidine (5.65 g, 65 mmol, 1.0 equiv) was dissolved in 100 mL of 1% AcOH in THF and cooled to 0 °C. Na(OAc)₃BH (24 g, 114 mmol, 1.75 equiv) and cyclopropylcarboxaldehyde (5.0 g, 71 mmol, 1.1 equiv) were added and the resulting mixture was stirred at 0 °C for 1 h and room temperature overnight (16 h). The reaction was diluted with 200 mL of 2 N NaOH, and the product was extracted 3x200 mL of CH₂Cl₂. The organic extracts were combined, dried over Na₂SO₄, decanted and concentrated to yield the desired product as a free-flowing oil (7.26 g, 79%) which was used without further purification; MS, m/z 142=M+1.

(b) 1-Cyclopropylmethyl-pyrrolidin-3-one.




A solution of oxalyl chloride (13.1 g, 103 mmol, 2.0 equiv) was prepared in 200 mL of dry CH_2Cl_2 and cooled under Ar to -78°C . DMSO (16.1 g, 206 mmol, 4.0 equiv) was added as a solution in 20 mL of CH_2Cl_2 dropwise over a 30 min period giving vigorous gas formation. After addition, the mixture was stirred for an additional 15 min and then a solution of 1-cyclopropylmethyl-3-hydroxy-pyrrolidine (7.26 g, 52 mmol, 1.0 equiv) in 50 mL of CH_2Cl_2 was added dropwise over a 30 min period. After complete addition the reaction was stirred an additional a hour at -78°C . Et_3N (31 g, 309 mmol, 6.0 equiv) was added over a period of 10 min. The cold-bath was removed and the mixture was stirred while warming for 1 h. The mixture was diluted with 500 mL of water and 100 mL of CH_2Cl_2 . After thorough mixing, the layers were separated and the organic layer was washed with 200 mL of water, dried over Na_2SO_4 , decanted, and concentrated to a yellow oil (6.1 g, 85%) which was used without further purification.

(c) 3-Amino-3-cyano-1-cyclopropyl-pyrrolidine.



1-Cyclopropylmethyl-pyrrolidin-3-one (6.1 g, 44 mmol, 1.0 equiv), NaCN (2.4 g, 48 mmol, 1.1 equiv) and NH_4Cl (2.6 g, 48 mmol, 1.1 equiv) were mixed in 88 mL of 2 M NH_3 in MeOH, and the resulting mixture was refluxed for 2 h at which time another 88 mL of 2 M NH_3 in MeOH was added followed by another 2 h at reflux. The reaction mixture was cooled, filtered, concentrated, and taken up in 100 mL of CH_2Cl_2 . The mixture was filtered a second time and concentrated to a red oil (5.9 g) which was used without further purification.

(d) Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopropylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.



N-(4-morpholinecarbonyl)-L-cyclohexyl alanine (1.00g, 3.52 mmol, 1.00 equiv) and EDC (1.01 g, 4.58 mmol, 1.30 equiv), and HoBt (0.72 g, 4.58 mmol, 1.30 equiv) were mixed 20 mL of DMF for 15 min followed by addition of 3-amino-3-cyano-1-cyclopropyl-pyrrolidine (0.86 g, 5.28 mmol, 1.5 equiv) and *N*-methyl-morpholine (1.42 g, 14.1 mmol, 4.0 equiv). The resulting solution was stirred at room temperature for 16 h. The reaction solution was diluted with 100 mL saturated sodium bicarbonate solution and the product was extracted with 2x100 mL EtOAc. The organic extracts were combined and concentrated. The crude product was purified by semi-prep reverse-phase HPLC using 40 to 90% CH₃CN in water over a gradient time of 30 min to give the desired product in two peaks (diastereomers) eluting at 9.5 min and 10.3 min respectively (128 mg and 98 mg, 15% total purified yield); MS, *m/z* 432=*M*+1 for both peaks.

Following the above procedures the following compounds were also prepared:

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-chloro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, *m/z* 502=*M*+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(cyclohexyl-methyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, *m/z* 474=*M*+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-methyl-ethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide MS, *m/z* 420=*M*+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-benzyloxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, *m/z* 574=*M*+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-benzyloxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 574=M+1

- 5 Morpholine-4-carboxylic acid {1-[3-cyano-1-(3,5-difluoro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 504=M+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2,6-difluoro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 504=M+1

10

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 536=M+1

- 15 Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-phenoxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 560=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 460=M+1

- 20 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-methyl-piperidine-4-yl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 475=M+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-methyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 482=M+1

25

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-phenoxy-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 560=M+1

- 30 Morpholine-4-carboxylic acid {1-[3-cyano-1-(4-fluoro-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 486=M+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2,4,6-trimethyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 510=M+1

5 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1*H*-indol-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 507=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopropyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 418=M+1

10 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-pyridin-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 469=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide; MS, m/z 448=M+1

15

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-2-hydroxymethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide; MS, m/z 472=M+1

20 Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 434=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-isopropyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide; MS, m/z 394=M+1

25 Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide; MS, m/z 408=M+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 448=M+1

30

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide; MS, m/z 422=M+1

5 Morpholine-4-carboxylic acid [1-(3-cyano-1-phenethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 482=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 406=M+1

10 Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide; MS, m/z 394=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 420=M+1

15

Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 474=M+1

20 Morpholine-4-carboxylic acid {1-[3-cyano-1-(*cis*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 474=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide; MS, m/z 420=M+1

25 Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 448=M+1

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 446=M+1

30

1-Benzyl-3-cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-
propionylamino}-pyrrolidine-2-carboxylic acid methyl ester; MS, m/z 526=M+1

5 Morpholine-4-carboxylic acid {1-[3-cyano-1-(*cis*-4-methyl-cyclohexyl)-pyrrolidin-3-
ylcarbamoyl]-3,3-dimethyl-butyl}-amide; MS, m/z 448=M+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-
ylcarbamoyl]-3,3-dimethyl-butyl}-amide; MS, m/z 448=M+1

10 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-(4-
iodo-phenyl)-ethyl]-amide; MS, m/z 580=M+1

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-methoxy-benzyl)-pyrrolidin-3-
ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 498=M+1

15

Morpholine-4-carboxylic acid {1-[3-cyano-1-(naphthalen-2-ylmethyl)-pyrrolidin-3-
ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 518=M+1

20 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentylmethyl-pyrrolidin-3-
ylcarbamoyl)-2-cyclohexyl-ethyl]-amide; MS, m/z 460=M+1


Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-
ylcarbamoyl)-3-methyl-butyl]-amide; MS, m/z 434=M+1

25 [1-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-carbamic
acid benzyl ester; MS, m/z 455=M+1

EXAMPLE 63


30 Morpholine-4-carboxylic acid (1-{3-cyano-1-[1-(toluene-4-sulfonyl)-1*H*-indol-3-
ylmethyl]-pyrrolidin-3-ylcarbamoyl}-2-cyclohexyl-ethyl)-amide.

(a) 1-[1-(Toluene-4-sulfonyl)-1*H*-indol-3-ylmethyl]-3-hydroxy-pyrrolidine.



1-(Toluene-4-sulfonyl)-1*H*-indole-3-carboxaldehyde (prepared as described in Chatterjee,
 5 R. K.; Indian J. Chem Sect. B 1994, 33(1), 32-37) was reacted with 3-hydroxypyrrolidine
 as described for cyclopropylcarboxaldehyde in Example 60 part (a) to provide the desired
 product.

(b) Morpholine-4-carboxylic acid (1-{3-cyano-1-[1-(toluene-4- sulfonyl)-1*H*-indol-3-
 10 ylmethyl]-pyrrolidin-3-ylcarbamoyl}-2-cyclohexyl-ethyl)-amide




The title compound was prepared from the product of part (a) and N-(4-
 morpholinecarbonyl)-L-cyclohexyl alanine by the procedure described in Example 60;
 MS, m/z 661= $M+1$.

EXAMPLE 64

Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-piperidine-4-carbonyl)-piperidin-4-yl]carbamoyl]-2-cyclohexyl-ethyl}-amide.

5



A solution of 1-methyl-piperidin-4-yl carboxylic acid (0.050 g, 0.30 mmol, 1.0 equiv) and EDC (0.057 g, 0.30 mmol, 1.0 equiv) was prepared in 15 mL of DMF. After 15 min morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-
10 amide hydrochloride (0.128 g, 0.30 mmol, 1.0 equiv) was added followed by N-methyl-morpholine (0.12 g, 1.2 mmol, 4.0 equiv) followed by stirring overnight (16 h). The reaction mixture was diluted with 100 mL of saturated sodium bicarbonate solution and the product was extracted with 2x50 mL of EtOAc. The combined organic extracts were concentrated. The crude product was purified by semi-prep reverse-phase HPLC using
15 20 to 80% CH₃CN in water over a gradient of 25 min to yield the desired product as a white solid (39 mg); MS, m/z 517=M+1.


Following the above procedure the following compound was also synthesized;

20 Morpholine-4-carboxylic acid {1-[4-cyano-1-(pyridine-4-carbonyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z 497=M+1

EXAMPLE 65


N-[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl]-4-methanesulfonylamino-benzamide.

- 5 (a) 4-Methanesulfonylamino-benzoic acid.



- Ethyl 4-amino-benzoate (5 g, 30 mmol, 1.0 equiv) was mixed in 50 mL of CH₂Cl₂ with Et₃N (6.1 g, 60 mmol, 2.0 equiv). The solution was cooled to 0 °C and methanesulfonyl chloride (3.8 g, 33 mmol, 1.1 equiv) was added as a solution in 15 mL of CH₂Cl₂ dropwise over a 15 min period. The reaction was stirred for 4 h at which time it was diluted with 50 mL of water. Layers were separated and the organic layer was washed with 50 mL of saturated sodium bicarbonate solution and concentrated. The resulting ester was dissolved in 50 mL of MeOH and treated with 50 mL of 5 N NaOH for 4 h. The reaction solution was extracted with Et₂O and the aqueous layer was acidified to give a white precipitate that was collected by filtration. The solid was dried and used without further purification.

- (b) *N*-[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl]-4-methanesulfonylamino-benzamide.



This compound was prepared from the product of part (a) using the procedure described in Example 24 to yield the desired product as a white solid; MS, m/z 490=M+1.

Following the above procedures the following compounds were also prepared:

N-[1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide; MS, m/z 526= $M+1$


N-[1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-methanesulfonylamino-benzamide; MS, m/z 552= $M+1$

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide; MS, m/z 464= $M+1$

EXAMPLE 66

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-1-oxy-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.

(a) Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-1-oxy-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.



Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide (0.50 g, 1.1 mmol, 1.0 equiv) was dissolved in CH_2Cl_2 (25 mL) and cooled to -78°C under Ar. Solid K_2CO_3 (0.22 g, 1.7 mmol, 1.5 equiv) was added followed by addition of solid *m*-CPBA (0.24 g, 1.1 mmol, 1.0 equiv). The resulting

mixture was stirred at -78°C for 2 h and, then, allowed to warm to room temperature. The reaction mixture was filtered and the solvent removed *in vacuo*. The residue was purified by flash chromatography on silica gel using 10-75% MeOH-EtOAc as a gradient eluent to give the desired product (0.32 g, 62%) as a white solid; MS, m/z 484=M+1.


5

EXAMPLE 66

3-Cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-pyrrolidine-1-carboxylic acid benzyl ester.

10


(a) 3-Hydroxy-pyrrolidine-1-carboxylic acid benzyl ester.



3-Hydroxy-pyrrolidine (10 g, 115 mmol, 1.0 equiv) was dissolved in 2 N NaOH (100 mL) and the mixture was cooled to 0°C . Benzylchloroformate (21 g, 126 mmol, 1.1 equiv) was added dropwise over a 45 min period. After addition, the reaction was stirred at room temperature for 4 h at which time the pH was adjusted to 7-8 using concentrated HCl. The product was extracted with 3x 100 mL of CH_2Cl_2 . The organic extracts were combined and dried over Na_2SO_4 , decanted and concentrated *in vacuo* to yield the desired product as a light yellow oil (24.1 g, 95%) that was used without further purification.


15

(b) 3-Oxo-pyrrolidine-1-carboxylic acid benzyl ester.




A solution of oxalyl chloride (12.6 g, 99 mmol, 2.0 equiv) was prepared in 250 mL of dry CH_2Cl_2 and cooled under Ar to -78°C . DMSO (15.5 g, 199 mmol, 4.0 equiv) was added dropwise over a 15 min period giving vigorous gas formation. After addition, the mixture was stirred for an additional 25 min and then a solution of 3-hydroxy-
5 pyrrolidine-1-carboxylic acid benzyl ester (11 g, 50 mmol, 1.0 equiv) in 20 mL of CH_2Cl_2 was added dropwise over a 10 min period. After complete addition the reaction was stirred an additional a hour at -78°C . Et_3N (55 mL, 398 mmol, 8.0 equiv) was added over a period of 10 min. The cold-bath was removed and the mixture was stirred while warming for 2 h. The mixture was diluted with 500 mL of water. After thorough mixing,
10 the layers were separated and the aqueous layer was extracted 2x150 mL of CH_2Cl_2 . The combined organic layers were washed with 200 mL of sodium bicarbonate solution and 200 mL of brine, dried over Na_2SO_4 , decanted, and concentrated to a yellow oil. The product was purified by flash chromatography on silica gel using CH_2Cl_2 as eluent to yield the desired product as a colorless oil (8.5 g).

15 (c) 3-Amino-3-cyano-pyrrolidine-1-carboxylic acid benzyl ester.



3-Amino-3-cyano-pyrrolidine-1-carboxylic acid benzyl ester was prepared from the ketone from part (b) using the procedure described in Example 1 part (a) to yield the
20 desired product as a 2 to 1 to 1 mixture of amino-nitrile, cyanohydrin and starting ketone that was used without further purification.

(d) 3-Cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-pyrrolidine-1-carboxylic acid benzyl ester.




3-Cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-
pyrrolidine-1-carboxylic acid benzyl ester was prepared from the amine from part (c)
using the procedure described in Example 1 part (a) to yield after purification on silica,
5 the desired product as an off-white hard foam; MS, m/z 512= $M+1$.

Following the above procedures the following compound was also synthesized;

3-Cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-
10 pyrrolidine-1-carboxylic acid (2-propen-1-yl) ester; MS, m/z 462= $M+1$.

EXAMPLE 68

Morpholine-4-carboxylic acid {1-[3-cyano-1-(5,5-dimethyl-3-oxo-cyclohex-1-enyl)-
15 pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide.




3-Cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-pyrrolidine-1-carboxylic acid (2-propen-1-yl) ester (1.35 g, 2.93 mmol, 1.00 equiv) was dissolved in 35 mL of CH₂Cl₂ along with dimedone (3.30 g, 23.5 mmol, 8.03 equiv). Pd(PPh₃)₄ (0.25 g, 0.22 mmol, 0.07 equiv) was added and the suspension was stirred at room temperature for 3.5 h. The reaction mixture was concentrated and taken up into EtOAc (100 mL) and extracted with 1 N HCl (2x50 mL). Concentration of the organic phase and purification of the crude mixture by reverse-phase HPLC provide the product as two separate diastereomers; MS, m/z 500=M+1.

EXAMPLE 69


4-Cyano-4-{3-cyclohexyl-2-[(piperidine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester.

(a) 4-Cyano-4-{3-cyclohexyl-2-[(1-*t*-butoxycarbonyl-piperidine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester.



This intermediate was prepared from 1-*t*-butoxycarbonyl-piperidine carboxylic acid and 4-cyano-4-{[3-cyclohexyl-2-amino]-propionylamino}-piperidine carboxylic acid ethyl ester hydrochloride using the procedure described in Example 24.

(b) 4-Cyano-4-{3-cyclohexyl-2-[(piperidine-4-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester.



The ester from (a) was dissolved in 10 mL of 4 N HCl in 1,4-dioxane at 0 °C for 1 hour. The solution was concentrated *in vacuo* and the salt neutralized by sodium bicarbonate solution and the product extracted with CH₂Cl₂. After concentration of the organic
 5 extract, the crude product was purified by reverse-phase HPLC to yield the desired product; MS, m/z 462=M+1.

Following the above procedures the following compounds were also synthesized:

- 10 4-Cyano-4-{3-cyclohexyl-2-[(4-methyl-piperazine-1-carbonyl)-amino]-propionylamino}-piperidine-1-carboxylic acid ethyl ester; MS, m/z 477=M+1

4-Methyl-piperazine-1-carboxylic acid [1-(4-cyano-tetrahydro-pyran-4-ylcarbamoyl)-2-cyclohexyl-ethyl]amide; MS, m/z 406=M+1.


15

EXAMPLE 70

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-azetidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.


20

(a) 3-Amino-1-benzyl-3-cyano-azetidine.



1-Benzyl-3-oxo-azetidine (1.6 g, 10 mmol, 1.0 equiv), prepared as described in the literature (Katritzky, A. R.; Cundy, D. J.; J. Heterocyclic Chem. 1994, 31 271-275), was dissolved in dry MeOH and the solution was cooled to -78°C . Gaseous ammonia was bubbled through the solution for 30 mins at which time 3 angstrom molecular sieves were added and the mixture transferred to a pressure tube. The solution was heated for 30 min at 60°C . The mixture was cooled to -78°C , the tube opened and KCN (0.65 g, 10 mmol, 1.0 equiv) and NH_4Cl (0.27 g, 5 mmol, 0.5 equiv) were added and the tube was resealed and heated at 60°C for 4 h. The reaction mixture was filtered and the filtrate was evaporated. The crude residue was purified by flash chromatography using 2% MeOH in CH_2Cl_2 to give the desired product (0.11 g, 6%) as a brown oil; MS, m/z 188=M+1.


(b) Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-azetidin-3-ylcarbonyl)-2-cyclohexyl-ethyl]-amide.



The title compound was prepared from 3-amino-1-benzyl-3-cyano-azetidine and *N*-(4-morpholinecarbonyl)-L-cyclohexyl alanine using to the procedure described in Example 1 step (d) to yield the desired product as a white solid; MS, m/z 454=M+1.

EXAMPLE 71

Morpholine-4-carboxylic acid [1-(3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide.




5

3-Cyano-3-{3-cyclohexyl-2-[(morpholine-4-carbonyl)-amino]-propionylamino}-pyrrolidine-1-carboxylic acid benzyl ester (0.1 g, 0.20 mmol, 1.0 equiv) was dissolved in 15 mL of absolute EtOH. 10% Pd on carbon (20 mg) was added and the mixture was stirred under 1 atm of H₂ until the starting material disappeared by TLC (5% MeOH in CH₂Cl₂). The crude mixture was filtered on diatomaceous earth and the filtrate was concentrated. The crude material was purified by reverse-phase HPLC to give two diastereomers; MS, m/z 378=M+1.

15

EXAMPLE 72

Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide.



Reductive amination of morpholine-4-carboxylic acid [1-(3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide with 2,2-dimethyl-2-phenyl-acetaldehyde and $\text{Na}(\text{OAc})_3\text{BH}$ in 1% AcOH in THF provided the desired product; MS, $510=\text{M}+1$.

5

Following the above procedure the following compounds were also synthesized;


Morpholine-4-carboxylic acid {1-[3-cyano-1-(indan-2-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z $508=\text{M}+1$

10

Morpholine-4-carboxylic acid {1-[3-cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide; MS, m/z $488=\text{M}+1$.

EXAMPLE 72

15



2-[[Acetylimino-(4-methoxy-phenyl)-methyl]-amino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide (Method E).


(a) **N-(4-methoxy-thiobenzoyl)acetamide.**

5 A solution of acetyl chloride (4.69 g, 59.8 mmol) in acetone (20 mL) was added dropwise to a solution of 4-methoxythiobenzamide (5.00 g, 29.9 mmol) and pyridine (4.76 g, 60.1 mmol) in acetone (30 mL). The reaction mixture was heated to reflux for 30 min then poured onto ice water. The resulting precipitate was isolated via filtration and dried under vacuum overnight to provide a light yellow/orange solid (4.52 g, 72%). ¹H
10 NMR (400 MHz, CDCl₃) δ 2.56 (s, 3H), 3.87 (s, 3H), 6.89 (dd, *J* = 6.9, 2.0 Hz, 2H), 7.77 (dd, *J* = 6.9, 2.0 Hz, 2H).

(b) **2-[[Acetylimino-(4-methoxy-phenyl)-methyl]-amino]-N-(4-cyano-1-methyl-
15 piperidin-4-yl)-3-cyclohexyl-propionamide.**

2-Chloro-N-methylpyridinium iodide (660 mg, 2.58 mmol), was added to a solution of N-(4-methoxy-thiobenzoyl)acetamide (420 mg, 2.01 mmol), 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt (730 mg, 2.00 mmol),
20 and N,N-diisopropylethylamine (1.05 mL, 6.02 mmol) in dichloromethane (8.0 mL). The reaction mixture was stirred at room temperature for 2 h, then diluted with dichloromethane (100 mL) and washed with 2x150 mL of saturated sodium bicarbonate. The organic phase was dried (MgSO₄) and concentrated. The resulting residue was chromatographed over 100 g of flash silica first using EtOAc, then
25 dichloromethane/methanol 9:1 as the eluant to provide the desired product as an off white solid (377 mg, 40%). ¹H NMR (400 MHz, DMSO-d₆) δ 0.70-0.90 (m, 2H), 1.00-1.30 (m, 4H), 1.35-1.65 (m, 8H), 1.72 (s, 3H), 1.85-2.20 (m, 6H), 2.48-2.60 (m, 1H), 3.78 (s, 3H), 4.20-4.35 (m, 1H), 6.95-6.99 (m, 2 H), 7.33 (d, *J* = 8.4 Hz, 1H), 7.72 (d, *J* = 8.4 Hz, 1H). MS, *m/z* 468 = *M*+1.

EXAMPLE 73




2-[(Acetylimino-phenyl-methyl)-amino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide.

(a) Thiobenzoyl acetamide was prepared according to the procedure from Example 1, step a, starting with thiobenzamide.

(b) The title compound was prepared starting from thiobenzoyl acetamide and 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 72, step b. MS, m/z 438 = $M+1$.


EXAMPLE 74



2- {[Acetylimino-(4-fluoro-phenyl)-methyl]-amino} -N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide.

- 5 (a) N-(4-Fluoro-thiobenzoyl) acetamide was prepared according to the procedure from Example 72, step a, starting with 4-fluorothiobenzamide.
- (b) The title compound was prepared starting from N-(4-fluoro-thiobenzoyl) acetamide and 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 72, step b. MS, m/z 456
- 10 = M+1.

EXAMPLE 75




2-[(Acetylimino-phenyl-methyl)]-amino-N-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide.

15

- (a) The title compound was prepared starting from thiobenzoyl acetamide and 2-amino-N-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 72, step b, except that the
- 20 compound was purified by HPLC using a 20 x 250 mm C18 reverse phase column with the method being 20% acetonitrile in water to 90% acetonitrile in water. MS, m/z 480 = M+1.

EXAMPLE 76



{[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylen}-carbamic acid ethyl ester (Method E).

5

(a) (Morpholine-4-carbothioyl)-carbamic acid ethyl ester.


Morpholine (7.5 mL, 86.0 mmol) was added dropwise to a solution of ethyl isothiocyanato formate (10.0 mL, 84.8 mmol) in tetrahydrofuran (200 mL). The reaction mixture was stirred at room temperature for 2.5 h, then concentrated and dried under vacuum to provide the desired product as a white solid (16.5 g, 89%). This material was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 1.28 (t, J = 7.1 Hz, 3H), 3.61-3.97 (m, 8H), 4.16 (q, 7.1 Hz, 2H), 7.44 (br s, 1H).

(b) {[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylen}-carbamic acid ethyl ester.

2-Chloro-N-methylpyridinium iodide (680 mg, 2.66 mmol), was added to a solution of (Morpholine-4-carbothioyl)-carbamic acid ethyl ester (450 mg, 2.06 mmol), 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt (745 mg, 2.04 mmol), and N,N-diisopropylethylamine (1.10 mL, 6.3 mmol) in dichloromethane (8.0 mL). The reaction was stirred at room temperature for 2.5 h then taken up in 10% citric acid solution and washed with EtOAc. The aqueous phase was

then basified with saturated sodium carbonate and extracted with EtOAc. The organic extract was dried (MgSO_4) and concentrated to provide the desired product as a white solid (250 mg, 26%). This material was further purified by HPLC using a 20 x 250 mm C_{18} reverse phase column with the method being 20% acetonitrile in water to 90% acetonitrile in water. MS, m/z 477 = $M+1$.


EXAMPLE 77



{[1-(4-Cyano-1-propyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethylimino}-morpholin-4-yl-methyl}-carbamic acid ethyl ester.

The title compound was prepared starting from (Morpholine-4-carbothioyl)-carbamic acid ethyl ester and 2-amino -N-(4-cyano-1-propyl-piperidin-4-yl)-3-cyclohexylpropionamide bis hydrochloride salt according to the procedure from Example 76, step b, except that the compound was first purified by chromatography over silica gel using 9:1 methylene chloride : methanol as the eluant prior to reverse phase HPLC purification. MS, m/z 505 = $M+1$.

EXAMPLE 78




{[1-(4-Cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylimino]-morpholin-4-yl-methyl}-carbamic acid ethyl ester.

- 5 The title compound was prepared starting from (Morpholine-4-carbothioyl)-carbamic acid ethyl ester and 2-amino -4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)amide bis hydrochloride salt according to the procedure from Example 76. MS, m/z 460 = M+1.

10

EXAMPLE 79




- 15 ({1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethylamino}-morpholin-4-yl-methylene)-carbamic acid ethyl ester.

The title compound was prepared starting from (Morpholine-4-carbothioyl)-carbamic acid ethyl ester and 2-amino-N-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-

cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 76, step b, . MS, m/z 519 = M+1.

EXAMPLE 80



N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-phenyl-methyl)-amino]-propionamide (Method F).

(a) Benzimidic acid methyl ester.

Benzimidic acid methyl ester hydrochloride (5 g, 29.1 mmol) was partitioned between saturated sodium carbonate solution (200 mL) and diethyl ether (100 mL). The organic layer was dried (MgSO_4) and concentrated to provide the desired product as a colorless liquid (3.20 g, 81%). This material was used without further purification. ^1H NMR (400 MHz, CDCl_3) δ 3.93 (s, 3H), 7.39-7.46 (m, 3H), 7.75 (d, J = 1.1 Hz, 2H).

(b) 1-Ethyl-3-(methoxy-phenyl-methylene)-urea.


A neat mixture of benzimidic acid methyl ester (750 mg, 5.56 mmol) and ethyl isocyanate (808 mg, 11.3 mmol) was stirred at 50 °C for 24 h. Excess isocyanate was removed under vacuum to provide the desired product as a colorless viscous oil (1.09 g, 95%). This material was used without further purification. ^1H NMR (400 MHz, CDCl_3)

δ 1.07 (t, $J = 7.3$ Hz, 3 H), 3.25 (q, $J = 7.3$ Hz, 2 H), 3.87 (s, 3H), 4.97 (br s, 1H), 7.26-7.40 (m, 2H), 7.45 (d, $J = 7.4$ Hz, 1H), 7.69-7.71 (m, 2H). MS, m/z 207 = $M+1$.

(c) N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-phenyl-methyl)-amino]-propionamide.

A solution of 1-ethyl-3-(methoxy-phenyl-methylene)-urea (350 mg, 1.70 mmol), 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt (512 mg, 1.40 mmol) and N,N-diisopropylethylamine (352 mg, 2.73 mmol) in dry methanol (5.0 mL) was stirred at room temperature for 60 h. The reaction mixture was concentrated and the resulting residue was chromatographed over 50 g of flash silica gel using dichloromethane to 5% methanol in dichloromethane as the eluant. This provided the desired product as a light yellow solid (280 mg, 43%) which was further purified by HPLC using a 20 x 250 mm C_{18} reverse phase column with the method being 20% acetonitrile in water to 90% acetonitrile in water. MS, m/z 467 = $M+1$.

EXAMPLE 81



N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(1,1-dioxo-1H-1 λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide (Method G).


(a) A suspension of 3-chloro-benzo[d]isothiazole 1,1-dioxide (300 mg, 1.49 mmol) and 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexylpropionamide bis

hydrochloride salt (500 mg, 1.37 mmol) was prepared in 5.5 mL of acetonitrile.

Triethylamine (575 μ L, 4.10 mmol) was added and the reaction mixture was stirred at room temperature for 1 day. The suspension was filtered to remove triethylamine hydrochloride and the filtrate was concentrated. The resulting residue was

5 chromatographed over 50 g of flash silica using dichloromethane/ methanol 9:1 as the eluant to provide the desired product as a light yellow solid (310 mg, 49%). ^1H NMR (400 MHz, CDCl_3) δ 0.25-0.45 (m, 1H), 0.65-0.85 (m, 2H), 0.95-1.10 (m, 2H), 1.30-1.60 (m, 7H), 1.75-1.85 (m, 2H), 1.85-2.2 (m, 2H), 2.31 (s, 3H), 2.35-2.50 (m, 3H), 2.65-2.80 (m, 2H), 4.60-4.70 (m, 1H), 7.35-7.50 (m, 2H), 7.58 (t, $J = 7.3$, 1H), 7.78 (d, $J = 7.7$ Hz, 1H), 7.81 (br s, 1H), 8.91 (br s, 1H). MS, m/z 458 = $M+1$.


EXAMPLE 82



N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-(1,1-dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-propionamide.

20 The title compound was prepared starting from 3-chloro-benzo[d]isothiazole 1,1-dioxide and 2-amino -N-(4-cyano-1-propyl-piperidin-4-yl)-3-cyclohexylpropionamide bis hydrochloride salt according to the procedure from Example 81, except that the compound was further purified by HPLC using a 20 x 250 mm C_{18} reverse phase column with the method being 20% acetonitrile in water to acetonitrile. MS, m/z 486 = $M+1$.


EXAMPLE 83



5 2-(1,1-dioxo-1H-1λ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid(4-
cyano-1-propylpiperidin-4-yl)-amide.

The title compound was prepared starting from 3-chloro-benzo[d]isothiazole 1,1-dioxide and 2-amino -4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)amide bis
 10 hydrochloride salt according to the procedure from Example 81, except that the
 compound was further purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column
 with the method being 20% acetonitrile in water to acetonitrile. MS, m/z 460 = M+1.


EXAMPLE 84



15 N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-(1,1-dioxo-1H-1λ⁶-
 20 benzo[d]isothiazol-3-ylamino)-propionamide.

The title compound was prepared starting from 3-chloro benzo[d]isothiazole 1,1-dioxide and 2-amino-N-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 81, except that the
5 compound was further purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 40% acetonitrile in water to acetonitrile. MS, m/z 500 = M+1.

EXAMPLE 85



N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H-1λ⁶-
15 benzo[d]isothiazol-3-ylamino)-propionamide.


The title compound was prepared starting from 3-chloro benzo[d]isothiazole 1,1-dioxide and 2-amino-N-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 81, except that the
20 compound was further purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 40% acetonitrile in water to acetonitrile. MS, m/z 512 = M+1.

EXAMPLE 86

N-(4-Cyano-methyl-piperidin-4-yl)-3-cyclohexyl-2-(3-oxo-3H-isindol-1-ylamino)-
25 propionamide.

The title compound was prepared starting from 3-imino-2, 3-dihydro-isoindol-1-one and 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 81, except that refluxing THF was used as the solvent. The compound was further purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 20% acetonitrile in water to acetonitrile. MS, m/z 422.5 = M+1.


EXAMPLE 87



4,4-Dimethyl-2-(3-oxo-3H-isoindol-1-ylamino)-pentanoic acid-(4-cyano-1-propyl-piperidin-4-yl)-amide.

The title compound was prepared from 3-imino-2, 3-dihydro-isoindol-1-one and 2-amino-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)amide bis hydrochloride salt according to the procedure from Example 86. MS, m/z 424.5 = M+1.

EXAMPLE 88



N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(5,6-difluoro-3-oxo-3H-isoindol-1-ylamino) propionamide.

5

(a) 2-Chloro-4,5-difluorobenzoic acid methyl ester.

2-Chloro-4,5-difluorobenzoic acid (1.93 g, 10 mmol) was dissolved in 20 mL of acetone.

10 Cesium carbonate (5.29 g, 15 mmol) was added followed by iodomethane (1.0 mL, 15 mmol). This reaction mixture was heated under reflux for 1 h and then cooled to room temperature. This suspension was then diluted with 40 mL of ethyl ether. The solid was removed by filtration and washed with ethyl ether. The filtrate was evaporated *in vacuo* to give the title compound in quantitative yield as a clear oil.

15

(b) 2-Cyano-4,5-difluorobenzoic acid methyl ester.

The above oil (2.06 g, 10 mmol) was dissolved in 10 mL of *N*-methyl pyrrolidinone.

Copper (I) cyanide (1.79 g, 20 mmol) was added. This mixture was heated at 195 °C

20 under nitrogen for 1 h. After cooling to room temperature, this solution was diluted with 100 mL of water. The resulting solid was collected by filtration. This solid was then suspended in a rapidly stirred solution of potassium cyanide (0.5 g) in 30 mL of water for 1 h. EtOAc (30 mL) was added. The mixture was filtered through diatomaceous earth. The organic phase was separated and the aqueous phase was extracted with EtOAc (20

mL x 2). The combined organic phase was washed with brine and dried over magnesium sulfate. The solvent was removed in vacuo. The residue was crystallized from ethyl ether and petroleum ether to give the title compound as a yellow solid (1.26 g, 64 %).


5 (c) 5,6-Difluoro-2,3-dihydro-3-imino-1*H*-isoindol-1-one.

The above solid (0.493 g, 2.5 mmol) was dissolved in 20 mL of MeOH. This solution was saturated with ammonia at 0 °C and then stirred in a pressure tube at room temperature for 3 days. The solid was collected by filtration and washed with ethyl ether
10 to give the title compound as a yellow solid (0.363 g, 80 %).

The title compound was prepared from 5,6-difluoro-2,3-dihydro-3-imino-1*H*-isoindol-1-one and 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt according to the procedure from Example 86. MS, m/z 458.3 = $M+1$.

15

EXAMPLE 89



20


N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(2-oxo-2H-benzo[e][1,3] oxazin-4-ylamino)-propionamide.

The title compound was prepared starting from 4-chloro-benzo[e][1,3] oxazin-2-one
25 (prepared from benzo [e][1,3] oxazin-2,4-dione and PCl_5 in refluxing toluene) and 2-

amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis
hydrochloride salt according to the procedure from Example 81. MS, m/z 438 = M+1.

5

EXAMPLE 90




10

N-(4-cyano-1-methyl-piperidin-4-yl)-2-(4-cyano-pyrimidin-2-ylamino)-3-cyclohexyl-
propionamide (Method G).

2-Chloro-4-pyrimidinecarbonitrile (0.3 mmol, Daves, G. D. Jr., O'Brien, D. E., Cheng, C.
15 C. J. *Het. Chem*, 1964, 1, 130) and 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-
cyclohexyl-propionamide (0.7 mmol) were dissolved in acetonitrile (10 mL) containing
N,N-diisopropylethylamine (0.6 mmol). The solution was heated to a gentle reflux for 17
h. The volatiles were evaporated and the residue was subjected to chromatography (silica
gel, eluant = EtOAc then MeOH). The methanolic fraction was concentrated to a
20 colorless solid which was rechromatographed (10% MeOH/EtOAc) to afford the title
compound as a colorless solid (52%). The material was recrystallized from
dichloromethane/petroleum ether.

25

EXAMPLE 91




N-(4-cyano-1-methyl-piperidin-4-yl)-2-(4-trifluoromethyl-pyrimidin-2-ylamino)-3-cyclohexyl-propionamide.

- 5 The title compound was prepared from 2-chloro-4-trifluoromethyl pyrimidine and 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide according to the procedure from Example 90. MS, m/z 439.5 = $M+1$.

10

EXAMPLE 92



- 15 N-(4-cyano-1-methyl-piperidine-4-yl)-3-cyclohexyl-2[N-cyano-morpholine-4-carboximidoyl]-amino-propionamide (Method H).

(a) 2-(N-Cyano-iminomethylene-amino)-N-(4-cyano-1-methyl-piperidine-4-yl)-3-cyclohexyl-propionamide.

20


A solution of diphenylcyanocarbonimidate (455 mg, 1.91 mmol), 2-amino-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide bis hydrochloride salt (680 mg, 1.86 mmol) and N,N-diisopropylethylamine (482 mg, 3.73 mmol) in isopropanol (5.0 mL) was stirred overnight at room temperature. The reaction mixture was then filtered to provide the desired carbodiimide as a white powder (140 mg, 22%). This material was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 0.80-1.00 (m, 2H), 1.05-1.20 (m, 1H), 1.20-1.40 (2H), 1.50-1.85 (m, 8H), 2.32 (s, 3H), 2.40-2.50 (m, 2H), 2.55-2.70 (m, 4H), 2.85-2.95 (m, 2H), 4.10-4.20 (m, 1H), 8.77 (br s, 1H). MS, m/z 343 = M+1.

- 10 (b) 2-(N-Cyano-benzimidoyl-amino)-N-(4-cyano-1-methyl-piperidine-4-yl)-3-cyclohexyl-propionamide.

A suspension of 2-(N-Cyano-iminomethylene-amino)-N-(4-cyano-1-methyl-piperidine-4-yl)-3-cyclohexyl-propionamide (120 mg, 0.35 mmol) in tetrahydrofuran (1 mL) was treated with morpholine (4 mL, 45.9 mmol). The reaction mixture was stirred at room temperature for 3 days then concentrated to dryness. The residue was purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 20% acetonitrile in water to 90% acetonitrile in water. MS, m/z 430 = M+1.

20

EXAMPLE 93



N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[[diethyl-carbamoylimino]-morpholin-4-yl-methyl]-amino}-propionamide (Method H).

(a) N,N-Diethyl carbamoyl thiocyanate.

5

A suspension of sodium thiocyanate (3.30 g, 40.7 mmol) in dry acetonitrile (25 mL) at 80°C was treated dropwise with a solution of N,N-diethyl carbamoyl chloride (5.0 g, 36.9 mmol) in dry acetonitrile (15 mL). The reaction mixture was stirred at 80°C for 50 min, cooled to room temperature, then filtered through a fine glass frit. The resulting filtrate
10 was used as a 0.9 M solution of N,N-diethyl carbamoyl thiocyanate in acetonitrile.

(b) N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(3-diethylamino-carbonyl-thioureido)-propionamide

15 A solution of 2-amino-N-(4-cyano-1-propyl-piperidin-4-yl)-3-cyclohexylpropionamide bis hydrochloride salt (560 mg, 1.53 mmol) and triethylamine (500 µL, 3.59 mmol) in acetonitrile (4 mL) was treated with a solution of N,N-diethyl carbamoyl thiocyanate in acetonitrile (3.0 mL, 2.7 mmol). The reaction mixture was stirred overnight at room temperature and concentrated on a rotary evaporator. The resulting residue was
20 chromatographed (ethyl acetate: hexanes 1:1 then ethyl acetate and finally methanol: methylene chloride 1:9 as the eluant) to provide the desired product as a light yellow solid (340 mg, 49%). MS, m/z 451.3 = M+1.

The title compound was prepared by treating a solution of the resulting thiourea (340 mg,
25 0.75 mmol) and triethylamine (230 µL, 1.65 mmol) in dry acetonitrile (4 mL) with mercury (II) chloride (225 mg, 0.83 mmol) and morpholine (200 µL, 2.23 mmol). The reaction mixture was stirred at room temperature for 4 h then filtered through a 0.45 µm filter disc. The resulting filtrate was filtered through a column of silica (5% methanol/methylene chloride as the eluant) and the resulting crude product was further
30 purified by HPLC using a 20 x 250 mm C₁₈ reverse phase column with the method being 20% acetonitrile in water to acetonitrile. MS, m/z 504.6 = M+1.

The following examples were prepared by Method H in a parallel fashion:

EXAMPLE 94

5

{[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-pyrrolidin-1-yl-methyl}-carbamic acid ethyl ester. MS, m/z 461 = M+1.

EXAMPLE 95

10

{[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-piperidin-1-yl-methyl}-carbamic acid ethyl ester. MS, m/z 477 = M+1.

EXAMPLE 96

15

{Azepan-1-yl-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-methylene}-carbamic acid ethyl ester. MS, m/z 490 = M+1.

EXAMPLE 97

20

{Azocan-1-yl-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-methylene}-carbamic acid ethyl ester. MS, m/z 504 = M+1.

EXAMPLE 98

25

1- {[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-ethoxycarbonylimino-methyl}-piperidine-4-carboxylic acid ethyl ester. MS, m/z 548 = M+1.

30

EXAMPLE 99

1-[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-ethoxycarbonylimino-methyl]-piperidine-3-carboxylic acid ethyl ester. MS, m/z 548 = M+1.

5

EXAMPLE 100

10

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(4-pyrrolidin-1-yl-piperidin-1-yl)-methylene]-carbamic acid ethyl ester. MS, m/z 545 = M+1.

15

EXAMPLE 101

{[1,4']Bipiperidinyl-1'-yl-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-methylene}-carbamic acid ethyl ester. MS, m/z 559 = M+1.

20

EXAMPLE 102

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(4-phenyl-piperazin-1-yl)-methylene]-carbamic acid ethyl ester. MS, m/z 553 = M+1.

25

EXAMPLE 103

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(4-ethyl-piperazin-1-yl)-methylene]-carbamic acid ethyl ester. MS, m/z 505 = M+1.

30

EXAMPLE 104

{{(4-Acetyl-piperazin-1-yl)-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-methylene}-carbamic acid ethyl ester. MS, m/z 519 = M+1.

EXAMPLE 105

5

4-[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-ethoxycarbonylimino-methyl]-piperazine-1-carboxylic acid ethyl ester. MS, m/z 549 = M+1.

10

EXAMPLE 106

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(3,3,5-trimethyl-6-aza-bicyclo[3.2.1]oct-6-yl)-methylene]-carbamic acid ethyl ester. MS, m/z 544 = M+1.

15

The following examples may also be made by the methods described above:

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

{[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

{[1-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

{[1-(3-Cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

((1-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butylamino)-morpholin-4-yl-methylene)-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

((1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butylamino)-morpholin-4-yl-methylene)-carbamic acid ethyl ester

((1-[3-Cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butylamino)-morpholin-4-yl-methylene)-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid [3-cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

{{1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

{{1-(3-Cyano-1-phenethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4,4-dimethyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-propionamide

N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-[(methanesulfonylimino-morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-N-(4-cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(methanesulfonylimino-morpholin-4-yl-methyl)-amino]-propionamide

{[1-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-propionamide

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-[(methanesulfonylimino-morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-N-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-3-cyclohexyl-propionamide

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-[(methanesulfonylimino-morpholin-4-yl-methyl)-amino]-propionamide

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-N-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-propionamide

{[1-(3-Cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

((1-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethylamino)-morpholin-4-yl-methylene)-carbamic acid ethyl ester

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-[(ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-N-[3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-
[(methanesulfonylimino-morpholin-4-yl-methyl)-amino]-propionamide

N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-[(methanesulfonylimino-
morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-N-[3-cyano-1-(1-ethyl-propyl)-
pyrrolidin-3-yl]-3-cyclohexyl-propionamide

N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-[(ethylcarbamoylimino-
morpholin-4-yl-methyl)-amino]-propionamide

{[1-[3-Cyano-1-(1H-indol-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-
ethylamino]-morpholin-4-yl-methylene)-carbamic acid ethyl ester

N-[3-Cyano-1-(1H-indol-3-ylmethyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-
[(ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-propionamide

N-[3-Cyano-1-(1H-indol-3-ylmethyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-
[(methanesulfonylimino-morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-N-[3-cyano-1-(1H-indol-3-ylmethyl)-
pyrrolidin-3-yl]-3-cyclohexyl-propionamide

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-[(carbamoylimino-morpholin-4-yl-methyl)-amino]-
3-cyclohexyl-propionamide

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-morpholin-4-
yl-methyl)-amino]-propionamide

{[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethylamino]-morpholin-4-
yl-methylene)-carbamic acid ethyl ester

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-3-cyclohexyl-2-[(methanesulfonylimino-morpholin-
4-yl-methyl)-amino]-propionamide

N-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-[(methanesulfonylimino-morpholin-4-yl-methyl)-amino]-propionamide

{[1-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)carbamoyl]-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

N-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-H-(3-cyano-1-phenethyl-pyrrolidin-3-yl)-3-cyclohexyl-propionamide

{[1-(4-Cyano-1-propyl-piperidin-4-yl)carbamoyl]-3-methyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

{[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-3-methyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

{[1-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

{[1-(3-Cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

([1-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-3-methyl-butylamino]-morpholin-4-yl-methylene)-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

((1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3-methyl-butylamino)-morpholin-4-yl-methylene)-carbamic acid ethyl ester

((1-[3-Cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-3-methyl-butylamino)-morpholin-4-yl-methylene)-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid [3-cyano-1-(1-H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

{[1-(3-Cyano-1-phenethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-morpholin-4-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

{[1-(4-Cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylamino]-phenyl-methylene}-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-phenyl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Methanesulfonylimino-phenyl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[(Carbamoylimino-phenyl-methyl)-amino]-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

{[1-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butylamino]-piperazin-1-yl-methylene}-carbamic acid ethyl ester

2-[(Ethylcarbamoylimino-piperazin-1-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Methanesulfonylimino-piperazin-1-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-[(Carbamoylimino-piperazin-1-yl-methyl)-amino]-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

{{1-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethylamino}-pyridin-4-yl-methylene)-carbamic acid ethyl ester

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-[(ethylcarbamoylimino-pyridin-4-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-pyridin-4-yl-methyl)-amino]-N-[3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-[(methanesulfonylimino-pyridin-4-yl-methyl)-amino]-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(ethylcarbamoylimino-pyrazin-2-yl-methyl)-amino]-propionamide

2-[(Carbamoylimino-pyrazin-2-yl-methyl)-amino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(methanesulfonylimino-pyrazin-2-yl-methyl)-amino]-propionamide

{{1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino}-pyrazin-2-yl-methylene}-carbamic acid ethyl ester

{{1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butylamino]-carbamoylimino-methyl}-carbamic acid benzyl ester

{{1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butylamino]-ethylcarbamoylimino-methyl}-carbamic acid benzyl ester

{{1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butylamino]-methanesulfonylimino-methyl}-carbamic acid benzyl ester

2-(1,1-Dioxo-1H-1λ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid [3-cyano-1-(1H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-propionamide

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-(1,1-dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-propionamide

N-[3-Cyano-1-(1H-indol-3-ylmethyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-(1,1-dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-propionamide

N-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-propionamide

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-propionamide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid [3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid [3-cyano-1-(1H-indol-3-ylmethyl)-pyrrolidin-3-yl]-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid (3-cyano-1-phenethyl-pyrrolidin-3-yl)-amide

2-(1,1-Dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-ylamino)-4-methyl-pentanoic acid (1-benzyl-3-cyano-pyrrolidin-3-yl)-amide

N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-(3-oxo-3H-isoindol-1-ylamino)-propionamide

4-Methyl-2-(3-oxo-3H-isoindol-1-ylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(3,4-dihydro-1H-pyrano[4,3-c]pyridin-5-ylamino)-propionamide

2-(3,4-Dihydro-1H-pyrano[4,3-c]pyridin-5-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

2-(3,4-Dihydro-1H-pyrano[4,3-c]pyridin-5-ylamino)-4-methyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(isoquinolin-1-ylamino)-propionamide

2-(Isoquinolin-1-ylamino)-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-(Isoquinolin-1-ylamino)-4-methyl-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide

2-(Imidazo[1,5-a]pyridin-3-ylamino)-4-methyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

2-(Imidazo[1,5-a]pyridin-3-ylamino)-4,4-dimethyl-pentanoic acid (3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-amide

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(imidazo[1,5-a]pyridin-3-ylamino)-propionamide

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-(8-oxo-8,9-dihydro-7H-purin-6-ylamino)-propionamide

4-Methyl-2-(8-oxo-8,9-dihydro-7H-purin-6-ylamino)-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

4,4-Dimethyl-2-(8-oxo-8,9-dihydro-7H-purin-6-ylamino)-pentanoic acid [3-cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-amide

2-{1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butylamino}-pyrimidine-4-carboxylic acid amide

2-{1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3-methyl-butylamino}-pyrimidine-4-carboxylic acid amide

2-{1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethylamino}-pyrimidine-4-carboxylic acid amide

4-Methyl-2-(2-oxo-1,2-dihydro-quinazolin-4-ylamino)-pentanoic acid (3-cyano-1-cyclopentylmethyl-pyrrolidin-3-yl)-amide

N-(3-Cyano-1-cyclopentylmethyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(2-oxo-1,2-dihydro-quinazolin-4-ylamino)-propionamide

4,4-Dimethyl-2-(2-oxo-1,2-dihydro-quinazolin-4-ylamino)-pentanoic acid (3-cyano-1-cyclopentylmethyl-pyrrolidin-3-yl)-amide

N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-(4-oxo-3,4-dihydro-phthalazin-1-ylamino)-propionamide

4-Methyl-2-(4-oxo-3,4-dihydro-phthalazin-1-ylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

4,4-Dimethyl-2-(4-oxo-3,4-dihydro-phthalazin-1-ylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-(1H-Indazol-3-ylamino)-4-methyl-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

2-(1H-Indazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

N-[4-Cyano-1-(2-hydroxy-ethyl)-piperidin-4-yl]-3-cyclohexyl-2-(1H-indazol-3-ylamino)-propionamide

4-Methyl-2-(2-oxo-2H-benzo[e][1,3]oxazin-4-ylamino)-pentanoic acid (4-cyano-1-methyl-piperidin-4-yl)-amide

4,4-Dimethyl-2-(2-oxo-2H-benzo[e][1,3]oxazin-4-ylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

N-[4-Cyano-1-(2-hydroxy-ethyl)-piperidin-4-yl]-3-cyclohexyl-2-(2-oxo-2H-benzo[e][1,3]oxazin-4-ylamino)-propionamide

2-(6-Hydroxy-1,1-dioxo-1H-1λ⁶-benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methyleneamide

4-Methyl-piperazine-1-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methyleneamide

4,4-Dimethyl-2- {[morpholin-4-yl-(2-morpholin-4-yl-ethylcarbamoylimino)-methyl]-amino}-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2- {[N-(5-methyl-oxazol-2-yl)-morpholine-4-carboximidoyl]-amino}-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2- {[N-(1-methyl-1H-imidazol-2-yl)-morpholine-4-carboximidoyl]-amino}-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2- {[N-(2-methyl-2H-[1,2,4]triazol-3-yl)-morpholine-4-carboximidoyl]-amino}-propionamide

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)-methylene]-carbamic acid ethyl ester

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(2-methoxymethyl-morpholin-4-yl)-methylene]-carbamic acid ethyl ester

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(2,6-dimethyl-morpholin-4-yl)-methylene]-carbamic acid ethyl ester

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-{{N-(4-methoxy-phenyl)-morpholine-4-carboximidoyl}-amino}-propionamide

4-({N-[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl}-morpholine-4-carboximidoyl)-amino)-benzamide

2-({N-[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl}-morpholine-4-carboximidoyl)-amino)-oxazole-5-carboxylic acid amide

2-({N-[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl}-morpholine-4-carboximidoyl)-amino)-oxazole-4-carboxylic acid amide

5-({N-[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl}-morpholine-4-carboximidoyl)-amino)-pyridine-2-carboxylic acid amide

2-({N-[1-(4-Cyano-1-methyl-piperidin-4-yl)carbamoyl]-2-cyclohexyl-ethyl}-morpholine-4-carboximidoyl)-amino)-3H-imidazole-4-carboxylic acid amide

2-[(N-Benzooxazol-2-yl-morpholine-4-carboximidoyl)-amino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-[(N-thiazol-2-yl-morpholine-4-carboximidoyl)-amino]-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-{{N-(5-phenyl-thiazol-2-yl)-morpholine-4-carboximidoyl}-amino}-propionamide

2-{{N-(5-Carbamoylmethyl-oxazol-2-yl)-morpholine-4-carboximidoyl}-amino}-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2- {[N-(2-methyl-oxazol-5-yl)-morpholine-4-carboximidoyl]-amino}-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(5,6-dihydro-8H-imidazo[5,1-c][1,4]oxazin-3-ylamino)-propionamide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(5,6,8,8a-tetrahydro-1H-imidazo[5,1-c][1,4]oxazin-3-ylamino)-propionamide

[[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-(2-methylcarbamoyl-morpholin-4-yl)-methylene]-carbamic acid ethyl ester

{[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-[3-(1-methylcarbamoyl-2-phenyl-ethylcarbamoyl)-morpholin-4-yl]-methylene}-carbamic acid ethyl ester

N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(1H-indol-2-yl-amine)-propionamide

[(1-(4-cyano-1-methyl-piperidin-4-yl-carbamoyl)-2-naphthalen-2-yl-ethylamino]-morpholin-4-yl-methylene)-carbamic acid ethyl ester

[(1-(4-cyano-1-methyl-piperidin-4-yl-carbamoyl)-2-(6-dimethylaminomethyl,naphthalen-2-yl-ethylamino]-morpholin-4-yl-methylene)-carbamic acid ethyl ester

2-(Benzooxazol-2-ylamino)-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

2-(Benzooxazol-2-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-(Benzo[thiazol-2-ylamino)-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

2-(Benzo[thiazol-2-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-(1H-Benzoimidazol-2-ylamino)-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

2-(1H-Benzoimidazol-2-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(6-methanesulfonylamino-2H-indazol-3-ylamino)-propionamide

2-(6-Methanesulfonylamino-2H-indazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-(Benzo[d]isoxazol-3-ylamino)-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

2-(Benzo[d]isoxazol-3-ylamino)-4,4-dimethyl-hexanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-(Benzo[d]isothiazol-3-ylamino)-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

2-(Benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(7-methanesulfonylamino-imidazo[1,5-d]pyridin-3-ylamino)-propionamide

2-(7-Methanesulfonylamino-imidazo[1,5-a]pyridin-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[1-(2-Carbamoyl-ethyl)-1H-imidazol-2-ylamino]-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(3-ureido-pyridin-2-ylamino)-propionamide

4,4-Dimethyl-2-(3-ureido-pyridin-2-ylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

2-[1-(2-Carbamoyl-ethyl)-1H-imidazol-2-ylamino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide

4,4-Dimethyl-2-(4-trifluoromethyl-pyrimidin-2-ylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

{[1-(1-Benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylimino]-morpholin-4-yl-methyl}-methyl-carbamic acid ethyl ester

{[1-(4-Cyano-1-isopropyl-piperidin-4-ylcarbamoyl)-3-methyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid benzyl ester

{[1-(4-Cyano-1-ethyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid cyclopentyl ester

{[1-(4-Cyano-1-phenethyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylamino]-morpholin-4-yl-methylene}-carbamic acid 2-methoxy-ethyl ester

{[1-(4-Cyano-1-cyclohexyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylamino]-phenyl-methylene}-carbamic acid ethyl ester

5

METHODS OF THERAPEUTIC USE

The compounds of the invention are useful in inhibiting the activity of cathepsin S, K, F,
10 L and B. In doing so, these compounds are useful in blocking disease processes mediated by these cysteine proteases.

Compounds of this invention effectively block degradation of the invariant chain to CLIP by cathepsin S, and thus inhibit antigen presentation and antigen-specific immune
15 responses. Control of antigen specific immune responses is an attractive means for treating autoimmune diseases and other undesirable T-cell mediated immune responses. Thus, there is provided methods of treatment using the compounds of this invention for such conditions. These encompass autoimmune diseases and other diseases involving inappropriate antigen specific immune responses including, but not limited to,
20 rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, multiple sclerosis, Guillain-Barre syndrome, psoriasis, Grave's disease, myasthenia gravis, scleroderma, glomerulonephritis, atopic dermatitis, insulin-dependent diabetes mellitus and asthma. The compounds of the invention can also be used to treat other disorders associated with extracellular proteolysis such as Alzheimer's disease and
25 atherosclerosis. The compounds of the invention can also be used to treat other disorders

associated with inappropriate autoimmune responses, T-cell mediated immune responses, or extracellular proteolysis mediated by cathepsin S, unrelated to those listed above or discussed in the Background of the Invention. Therefore, the invention also provides methods of modulating an autoimmune disease comprising administering to a patient in
5 need of such treatment a pharmaceutically effect amount of a compound according to the invention.

Compounds of the invention also inhibit cathepsin K. In doing so, they may block inappropriate degradation of bone collagen and other bone matrix proteases. Thus, there
10 is provided a method for treating diseases where these processes play a role such as osteoporosis. Inhibition of cathepsins F, L, and B are also within the scope of the invention due to similarity of the active sites in cysteine proteases as described above.

For therapeutic use, the compounds of the invention may be administered in any
15 conventional dosage form in any conventional manner. Routes of administration include, but are not limited to, intravenously, intramuscularly, subcutaneously, intrasynovially, by infusion, sublingually, transdermally, orally, topically or by inhalation. The preferred modes of administration are oral and intravenous.

The compounds of this invention may be administered alone or in combination with
20 adjuvants that enhance stability of the inhibitors, facilitate administration of pharmaceutical compositions containing them in certain embodiments, provide increased dissolution or dispersion, increase inhibitory activity, provide adjunct therapy, and the like, including other active ingredients. Advantageously, such combination therapies
25 utilize lower dosages of the conventional therapeutics, thus avoiding possible toxicity and adverse side effects incurred when those agents are used as monotherapies. Compounds of the invention may be physically combined with the conventional therapeutics or other adjuvants into a single pharmaceutical composition. Advantageously, the compounds may then be administered together in a single dosage form. In some embodiments, the
30 pharmaceutical compositions comprising such combinations of compounds contain at least about 15%, but more preferably at least about 20%, of a compound of the invention

(w/w) or a combination thereof. Alternatively, the compounds may be administered separately (either serially or in parallel). Separate dosing allows for greater flexibility in the dosing regime.

- 5 As mentioned above, dosage forms of the compounds of this invention include pharmaceutically acceptable carriers and adjuvants known to those of ordinary skill in the art. These carriers and adjuvants include, for example, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, buffer substances, water, salts or electrolytes and cellulose-based substances. Preferred dosage forms include, tablet, capsule, caplet,
- 10 liquid, solution, suspension, emulsion, lozenges, syrup, reconstitutable powder, granule, suppository and transdermal patch. Methods for preparing such dosage forms are known (see, for example, H.C. Ansel and N.G. Popovich, *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 5th ed., Lea and Febiger (1990)). Dosage levels and requirements are well-recognized in the art and may be selected by those of ordinary skill
- 15 in the art from available methods and techniques suitable for a particular patient. In some embodiments, dosage levels range from about 10-1000 mg/dose for a 70 kg patient. Although one dose per day may be sufficient, up to 5 doses per day may be given. For oral doses, up to 2000 mg/day may be required. As the skilled artisan will appreciate, lower or higher doses may be required depending on particular factors. For instance,
- 20 specific dosage and treatment regimens will depend on factors such as the patient's general health profile, the severity and course of the patient's disorder or disposition thereto, and the judgment of the treating physician.

ASSESSMENT OF BIOLOGICAL PROPERTIES

25

Expression and Purification of recombinant human Cathepsin S

Cloning of human cathepsin S:

- U937 RNA was subjected to reverse transcriptase / polymerase chain reaction with
- 30 primer A (5'cacaatgaaacggctggtttg 3') and primer B (5'ctagattctgggtaagaggg 3') designed to specifically amplify the cathepsin S cDNA. The resulting 900 bp DNA

fragment was subcloned into pGEM-T (Promega) and sequenced to confirm its identity. This construct was used for all subsequent manipulations. This procedure is typical for cloning of known genes and is established in its field.

- 5 Human Pre-Pro-Cat S was removed from pGem-T vector (Promega, 2800 Woods Hollow Rd, Madison, WI 53711) by digestion with restriction enzyme SacII, followed by treatment with T4 DNA polymerase to generate a blunt end, and a second restriction enzyme digest with Sall. It was subcloned into pFastBac1 donor plasmid (GibcoBRL, 8717 Grovemont Cr., Gaithersburg, MD 20884) which had been cut with restriction
10 enzyme BamHI and blunt-ended and then cut with restriction enzyme Sall. The ligation mixture was used to transform DH5a competent cells (GibcoBRL) and plated on LB plates containing 100ug/ml ampicillin. Colonies were grown in overnight cultures of LB media containing 50ug/ml Ampicillin, plasmid DNA isolated and correct insert confirmed by restriction enzyme digestion. Recombinant pFastBac donor plasmid was
15 transformed into DH10Bac competent cells (GibcoBRL). Large white colonies were picked from LB plates containing 50ug/ml kanamycin, 7ug/ml gentamicin, 10ug/ml tetracycline, 100ug/ml Blueo-gal, and 40ug/ml IPTG. DNA was isolated and used to transfect Sf9 insect cells using CellFECTIN reagent (GibcoBRL). Cells and supernatant were harvested after 72 hours. Viral supernatant was passaged twice and presence of Cat
20 S confirmed by PCR of the supernatant.

- Sf9 cells were infected with recombinant baculovirus at a MOI of 5 for 48-72 hrs. Cell pellet was lysed and incubated in buffer at pH 4.5 at 37 for 2 hours to activate Cat S from pro-form to active mature form (Bromme, D & McGrath, M., Protein Science, 1996,
25 5:789-791.) Presence of Cat S was confirmed by SDS-PAGE and Western blot using rabbit anti-human proCat S.

Inhibition of Cathepsin S

- 30 Human recombinant cathepsin S expressed in Baculovirus is used at a final concentration of 10nM in buffer. Buffer is 50mM Na Acetate, pH 6.5, 2.5mMEDTA, 2.5mMTCEP. Enzyme is incubated with either compound or DMSO for 10 min at 37C. Substrate 7-

amino-4-methylcoumarin, CBZ-L-valyl-L-valyl-L-arginineamide (custom synthesis by Molecular Probes) is diluted to 20uM in water (final concentration of 5uM), added to assay and incubated for additional 10 minutes at 37 C. Compound activity is measured by diminished fluorescence compared to DMSO control when read at 360nm excitation and 460nm emission.

Examples listed above were evaluated for inhibition of cathepsin S in the above assay. All had IC₅₀ values of 100 micromolar or below.

Inhibition of Cathepsin K, F, L and B:

Inhibition of these enzymes by particular compounds of the invention may be determined without undue experimentation by using art recognized methods as provided hereinbelow each of which is incorporated herein by reference:

Cathepsin B, and L assays are to be found in the following references:

1. Methods in Enzymology, Vol.244, Proteolytic Enzymes: Serine and Cysteine Peptidases, Alan J. Barrett, ed.

Cathepsin K assay is to be found in the following reference:

2. Bromme, D., Okamoto, K., Wang, B. B., and Biroc, S. (1996) *J. Biol. Chem.* **271**, 2126-2132.

Cathepsin F assays are to be found in the following references:


3. Wang, B., Shi, G.P., Yao, P.M., Li, Z., Chapman, H.A., and Bromme, D. (1998) *J. Biol. Chem.* **273**, 32000-32008.

4. Santamaria, I., Velasco, G., Pendas, A.M., Paz, A., and Lopez-Otin, C (1999) *J. Biol. Chem.* **274**, 13800-13809.

Preferred compounds to be evaluated for inhibition of Cathepsin K, F, L and B in the above assays desirably have IC₅₀ values of 100 micromolar or below.

What is claimed is

1. A compound of formula (I):



wherein:

10 Het is piperidinyl, pyrrolidinyl, azetidiny, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, hexahydropyrimidinyl, hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidin-2-ylamine, dihydro-oxazolyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide or imidazolidinyl-2,4-dione, each
15 being optionally substituted with one or more R_5 ;

Y is O or S;

20 R_1 is C1-5 alkyl, C1-5 alkoxy, aryloxy, C3-7 cycloalkyl, phenyl, benzyl, naphthyl, tetrahydronaphthyl, C1-5alkylsulfonylC1-5alkyl, C3-7cycloalkylsulfonylC1-5alkyl, arylsulfonylC1-5alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl,
25 isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl, quinoxalinyl, or amino; wherein R_1 is optionally substituted by one or more R_a ;

R_a is C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkoxy, C1-5alkanoyl, C1-5alkanoyloxy, aryloxy, benzyloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_a may be further optionally substituted by one or more R_b ;

R_b is C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino;

5 R₂ is hydrogen or C1-3 alkyl;

R₃ is hydrogen, C1-5 alkyl, C2-5alkylene, C3-7 cycloalkyl, arylC1-3alkyl or aryl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is C1-5 alkyl, C3-7 cycloalkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxaliny, C1-5 alkoxy, aryloxy, C1-5
15 alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl,
20 indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny,
or R_c is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
25 independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny,
30 C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5

alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylalkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino;

R₄ is hydrogen or C1-3 alkyl;

R₅ is C1-5 alkyl chain optionally interrupted by one or two O or S, phenyl, naphthyl, arylC1-3alkyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, C1-5 alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, benzyloxycarbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl or quinolinyl, or R₅ is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or disubstituted by alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyridinylcarbonyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl or arylsulfonyl, or R₅ is halogen, hydroxy, oxo,

carboxy, cyano, nitro, amidino or guanidino, R_5 may be further optionally substituted by one or more R_6 ;

R_6 is C1-5 alkyl, C3-6 cycloalkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl, quinoxalinyl, C1-5 alkoxy, aryloxy, aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, furanyl, thienyl, pyrrolyl or pyridinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, benzyloxy, arylC1-3alkoxycarbonyl, amidino or guanidino;

X is O or S and

pharmaceutically acceptable derivatives thereof.

2. The compound according to claim 1

wherein:

Het is piperidinyl, pyrrolidinyl, tetrahydropyranyl or tetrahydrothiopyranyl each ring being substituted with one or more R_5 ;

Y is O;

R_1 is C1-3 alkyl, C1-3 alkoxy, C3-7 cycloalkyl, phenyl, benzyl, naphthyl, tetrahydronaphthyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl, pyridinyl, isoxazolyl, pyrazinyl, indolyl, quinolinyl, benzofuranyl, benzimidazolyl, benzoisoxazolyl or amino; wherein R_1 is optionally substituted by one or more R_a ;

R_a is C1-3 alkyl, phenyl, naphthyl, piperidinyl, indolinyl, morpholinyl, piperazinyl, furanyl, thienyl, benzimidazolyl, C1-3 alkoxy, C1-3 alkanoyl, phenoxy, naphthyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzoyloxy, carbamoyl
5 wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl or pyridinyl, or R_a is C1-5 alkanoylamino, benzoylamino, C1-3alkylsulfonyl, phenylsulfonyl, ureido wherein either nitrogen atom may be independently substituted by alkyl, phenyl, piperidinyl, morpholinyl, furanyl, thienyl or pyridinyl, C1-3
10 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, C1-5 alkylsulfonylamino, phenylsulfonylamino, C1-5 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl or pyridinyl, halogen, hydroxy, oxo, carboxy, nitro or cyano, R_a may be further optionally
15 substituted by one or more R_b ;

R_b is halogen, hydroxy, benzyloxy, oxo or cyano;

R_2 is hydrogen;

R_3 is C1-5 alkyl or C2-5 alkylene, C4-6 cycloalkyl or benzyl wherein R_3 is optionally substituted by one or more R_c ;

R_c is C1-4 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, C1-4 alkoxy, phenoxy, benzoyl, benzyloxy, indolinyl, imidazolyl, C1-3alkylthio, C1-3alkylsulfonyl, halogen, hydroxy, oxo, carboxy, nitro or cyano, R_c may be further optionally substituted by one or more R_d ;

R_d is methyl, phenyl, benzyl, benzyloxy, C1-3alkoxy, halogen, hydroxy, nitro or cyano;

R₄ is hydrogen;

R₅ is C1-4alkyl chain optionally interrupted by one O or S atom, phenyl, phenylC1-2alkyl, furanyl, pyrimidinyl, thienyl, C1-3 alkanoyl, benzoyl, C1-4 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl, phenyl, piperidinyl, morpholinyl, piperazinyl, furanyl, thienyl or pyridinyl, C1-3 alkylthio, phenylthio, C1-5 alkylaminosulfonyl, phenylaminosulfonyl, C1-5alkylamino wherein the nitrogen atom may be independently mono- or disubstituted by naphthylsulfonyl or pyridinylcarbonyl, halogen, hydroxy, carboxy, oxo or cyano, R₅ may be further optionally substituted by one or more R_c;

R_c is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthylmethyl, piperidinyl, morpholinyl, piperazinyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, benzimidazolyl, quinolinyl, isoquinolinyl, C1-4 alkoxy, benzoyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, piperidinyl, morpholinyl, furanyl, thienyl or pyridinyl, halogen, hydroxy, oxo or cyano; and

X is O.

3. The compound according to claim 2

wherein:

R₁ is methyl, ethyl, phenyl, piperidinyl, morpholinyl, piperazinyl, pyridinyl, pyrazinyl, furanyl, thienyl, benzyl, benzofuranyl, cyclohexyl, quinolinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is C1-3 alkyl, phenyl, piperidinyl, thienyl, C1-3 alkoxy, phenoxy, C1-3alkanoyl, C1-3 alkoxycarbonyl, benzyloxy, C1-3 alkanoylamino, thiophenyl, benzimidazolyl, C1-3 alkylthio or chloro, R_a may be further optionally substituted by one or more R_b ;

5

R_b is bromo, chloro, fluoro, iodo, hydroxy, oxo or cyano;

R_3 is methyl, ethyl, n-propyl, n-butyl, isobutyl, propenyl, butenyl, isobutenyl, C3-7 cycloalkyl or benzyl wherein R_3 is optionally substituted by one or more R_c ;

10

R_c is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, methoxy, ethoxy, methylthio, ethylthio, cyclohexyl, phenyl, naphthyl, imidazolyl, indolyl, bromo, chloro, fluoro, iodo, hydroxy, oxo, carboxy, nitro, benzoyl, benzyloxy, N-benzylimidazolyl or cyano, R_c may be further optionally substituted by one or more R_d ;

15

R_d is methyl, methoxy, ethoxy, chloro, fluoro, nitro or hydroxy;

R_5 is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, phenyl, methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, isobutyloxycarbonyl, tert-butoxycarbonyl and pyrimidinyl, R_5 may be further optionally substituted by one or more R_e ;

20

R_e is methyl, ethyl, n-propyl, isopropyl, phenyl, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, tert-butoxycarbonyl, bromo, chloro, fluoro, iodo, hydroxy, oxo or cyano.

25

4. The compound according to claim 3

30

wherein:

Het is piperidinyl or pyrrolidinyl;

R₁ is N-acetylamino-phenyl, chlorophenyl, methoxyphenyl, m-phenoxyphenyl,
5 morpholinyl, pyrazinyl, pyridinyl, furanyl, chlorothienyl, thienyl or thienylmethyl;

R₃ is n-butyl, isobutyl, 2,2-dimethylpropyl, cyclohexylmethyl, p-methoxybenzyl or 2-naphthylmethyl; and
wherein the configuration at the stereocenter defined by R₂ and R₃ when they are
10 different and the carbon they are attached to is defined as L; and

R₅ is methyl, propyl, isopropyl, ethoxycarbonyl, benzyloxycarbonyl, benzyl, phenethyl,
N,N-dimethylaminoacetyl or pyrimidinyl.

15

5. The compound according to claim 4

wherein:

20 Het is piperidin-4-yl or pyrrolidinyl;

R₁ is morpholinyl or N-acetylamino-phenyl;


R₃ is 2,2-dimethylpropyl or cyclohexylmethyl; and

25

R₅ is methyl, propyl, isopropyl, ethoxycarbonyl, benzyloxycarbonyl, benzyl, phenethyl,
N,N-dimethylaminoacetyl or pyrimidinyl.

30

6. A compound of formula (II):



5

wherein:

- Het is azepanyl, piperidinyl, pyrrolidinyl, azetidiny, oxepanyl, tetrahydropyranyl,
 10 tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, azocanyl, oxocanyl, 1,3-diazocanyl,
 1,4-diazocanyl, 1,5-diazocanyl, 1,3-dioxocanyl, 1,4-dioxocanyl, 1,5-dioxocanyl, 1,3-
 oxazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,3-diazepanyl, 1,4-diazepanyl, 1,3-
 dioxepanyl, 1,4-dioxepanyl, 1,3-oxazepanyl, 1,4-oxazepanyl, 1,2-thiazocanyl-1,1-
 dioxide, 1,2,8-thiadiazocanyl-1,1-dioxide, 1,2-thiazepanyl-1,1-dioxide, 1,2,7-
 15 thiadiazepanyl-1,1-dioxide, tetrahydrothiophenyl, hexahydropyrimidinyl,
 hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, pyrazolidinyl, dihydro-
 oxazolyl, dihydrothiazolyl, dihydroimidazolyl, isoxazoliny, oxazolidinyl, 1,2-
 thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide,
 imidazolidinyl-2,4-dione, imidazolidinyl, morpholinyl, dioxanyl, tetrahydropyridinyl,
 20 thiomorpholinyl, thiazolidinyl, dihydropyranyl, dithianyl, decahydro-quinolinyl,
 decahydro-isoquinolinyl, 1,2,3,4-tetrahydro-quinolinyl, indolinyl, octahydro-quinoliziny,
 dihydro-indoliziny, octahydro-indoliziny, octahydro-indolyl, decahydroquinazoliny,
 decahydroquinoxaliny, 1,2,3,4-tetrahydroquinazoliny or 1,2,3,4-tetrahydroquinoxaliny;
- 25 A C6-C10 bridged bicyclo wherein one or more carbon atoms are optionally replaced by
 a heteroatom chosen from N, O and S;

each being optionally substituted with one or more R₅;

Y is C(O), C(S) or S(O)₂;

5

R₁ is a bond, hydrogen, C1-10 alkyl, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, C3-8 cycloalkyloxy, aryl, benzyl, tetrahydronaphthyl, indenyl, indanyl, C1-10alkylsulfonylC1-10alkyl, C3-8cycloalkylsulfonylC1-10alkyl, arylsulfonylC1-10alkyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, 10 indolinyl, pyranyl, tetrahydropyranyl, tetrahydrothiopyranyl, thiopyranyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, tetrazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzisoxazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl, 15 tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, hydroxy or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond, C1-10 alkyl, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, 20 indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10 alkoxy, C1-10alkanoyl, C1- 25 10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, aryloxy carbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, 30 indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

with the proviso that R₁ and R_a simultaneously cannot be a bond;

R_b is a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more carbon atoms are optionally replaced by O, N, S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2 oxo groups, -NH₂, or one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl;

or R_b is C3-6 cycloalkyl, aryl, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, mono-C1-5alkylamino, di-C1-5alkylamino, carboxamide, amidino or guanidino;

5 R₂ is hydrogen or C1-3 alkyl;

R₃ is a bond, hydrogen, C1-10 alkyl, C2-10alkylene, C3-8 cycloalkyl, arylC1-5alkyl or aryl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is C1-10 alkyl, C3-8 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, decahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranyl, tetrahydropyranyl, tetrahydrothiopyranyl,
15 thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, dihydrobenzofuranyl, octohydrobenzofuranyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, tetrahydroquinolinyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10 alkoxy, aryloxy, C1-10 alkanoyl, aroyl, C1-10 alkoxycarbonyl, aryloxycarbonyl, C1-10 alkanoyloxy, aroyloxy,
20 carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
25 or R_c is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom
30 may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl,

oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl,
pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl,
quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
or R_c is C1-10 alkoxy carbonylamino, aryloxy carbonylamino, C1-10
5 alkyl carbamoyloxy, aryl carbamoyloxy, C1-10 alkyl sulfonylamino,
aryl sulfonylamino, C1-10 alkyl aminosulfonyl, aryl aminosulfonyl, amino wherein
the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl,
aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,
indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl,
10 tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,
benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or
quinoxalinyl,
or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c
may be further optionally substituted by one or more R_d;

15 R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-5alkyl, C1-5 alkoxy,
aryloxy, arylC1-5alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy,
cyano, nitro, amidino or guanidino;

20 R₂ and R₃ together with the carbon they are attached optionally form a nonaromatic 5-7
membered cycloalkyl or heterocyclic ring;

R₄ is hydrogen, hydroxy or C1-3 alkyl;

25 R₅ is a bond, hydrogen, carbonyl, C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-
10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, aryl, benzyl,
tetrahydronaphthyl, indenyl, indanyl, C3-7cycloalkylsulfonylC1-5alkyl, arylsulfonylC1-
5alkyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl,
30 thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, thiopyranyl,
tetrahydrothiopyranyl, furanyl, tetrahydrofuranlyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl,

thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridizinyl, tetrazolyl, triazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl, tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the
5 heterocyclyl moiety is selected from those herein described in this paragraph, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, benzyloxy, C1-10alkoxycarbonyl, arylC1-5alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl,
10 thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
or R₅ is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to
15 a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
20 or R₅ is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl,
25 thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R₅ may be further optionally substituted by one or more R₆;

R_e is C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-10 alkoxy, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, thiopyranyl, tetrahydrothiopyranyl, pyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, arylC1-3alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_e is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_e is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl,

tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_c may be further optionally substituted by one or more R_g ;

R_f is C1-5 alkyl, C3-6 cycloalkyl, tolylsulfonyl, C1-5 alkoxy, aryl, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

X is O or S and

pharmaceutically acceptable derivatives thereof.

7. The compound according to claim 6

wherein:

Het is piperidinyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, azetidiny, azepanyl, oxepanyl, tetrahydrofuranyl, oxetanyl, hexahydropyrimidinyl, hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, octahydro-indoliziny, octahydro-quinoliziny, decahydro-quinolinyl, 1,2,3,4-tetrahydro-quinolinyl, dihydro-oxazolyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-

1,1-dioxide, imidazolidinyl, pyrazolidinyl or a bridged bicyclo chosen from aza-bicyclo[3.2.1]octane, aza-bicyclo[2.2.1]heptane, aza-bicyclo[2.2.2]octane, aza-bicyclo[3.2.2]nonane, aza-bicyclo[2.1.1]hexane, aza-bicyclo[3.1.1]heptane, aza-bicyclo[3.3.2]decane and 2-oxa or 2-thia-5-aza-bicyclo[2.2.1]heptane;

each ring being substituted with one or more R_5 ;

Y is C(O) or S(O)₂;

R₁ is a bond, hydrogen, C1-7 alkyl, C1-7 alkoxy, C3-7 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, tetrahydronaphthyl, C1-7alkylsulfonylC1-7alkyl, C3-7cycloalkylsulfonylC1-7alkyl, arylsulfonylC1-7alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, isoxazolyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoisoxazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond C1-7 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-7 alkoxy, C1-7alkanoyl, C1-7alkanoyloxy, aryloxy, benzyloxy, C1-7 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_a is C1-7 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-7 alkyl-carbamoyloxy, aryl-carbamoyloxy, C1-7 alkylsulfonylamino, arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b ;

R_b is C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

R_2 is hydrogen or methyl or ethyl;

R_3 is a bond, hydrogen, C1-5 alkyl, C2-5 alkylene, C3-7 cycloalkyl, aryl-C1-3 alkyl or aryl wherein R_3 is optionally substituted by one or more R_c ;

R_c is C1-5 alkyl, C3-7 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranyl, tetrahydropyranyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5 alkanoyl, aroyl, C1-5 alkoxy-carbonyl, aryloxy-carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl,

thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

5 or R_c is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, 10 oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_c is C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, 15 arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, 20 benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

25 R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-4 alkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, halogen, hydroxy, oxo or cyano;

R₄ is hydrogen or methyl;

R₅ is a bond, hydrogen, carbonyl, C1-8 alkyl, C1-8alkoxyC1-8alkyl, C1-8alkylaminoC1-8alkyl, C1-8alkylthioC1-8alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-8 alkoxy,, aryloxy, C3-7 cycloalkyl, aryl, benzyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, thiopyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranlyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, tetrazolyl, triazolyl, pyrazolyl, indolyl, benzofuranlyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-7alkanoyl, aroyl, C1-7alkanoyloxy, benzyloxy, C1-7 alkoxycarbonyl, arylC1-4alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranlyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranlyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R₅ is C1-7 alkoxycarbonylamino, aryloxycarbonylamino, C1-7 alkylcarbamoyloxy, arylcarbamoyloxy, C1-7 alkylsulfonylamino, arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranlyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-7 alkyl, C1-7alkoxyC1-7alkyl, C1-7alkylaminoC1-7alkyl, C1-
5 7alkylthioC1-7alkyl wherein the sulfur atom may be oxidized to a sulfoxide or
sulfone, C1-7 alkoxy, C3-7 cycloalkyl, aryl, tetrahydronaphthyl, indanyl,
pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thiopyranyl,
tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl,
oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
10 benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl,
quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkanoyl, aroyl, C1-
5alkanoyloxy, aryloxy, benzyloxy, C1-5 alkoxycarbonyl, aryloxy carbonyl,
aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or
di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl,
15 thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl,
pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,
benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or
quinoxalinyl,

or R_e is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom
20 may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may
be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl,
morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl,
imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,
25 benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or
quinoxalinyl,

or R_e is C1-5 alkoxycarbonylamino, aryloxy carbonylamino, C1-5
alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino,
arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein
30 the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl,
aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl,

thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
benzofuranyl, benzothenyl, benzimidazolyl, benzthiazolyl, quinolinyl,
isoquinolinyl, quinazolinyl or quinoxalinyl,
or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or
5 guanidino, R_e may be further optionally substituted by one or more R_f ;

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, C1-3 alkoxy, cyclopropyl,
cyclohexyl, phenyl, naphthyl, phenoxy, benzyloxy, fluoro, chloro, bromo,
hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

10 and
X is O.

15 8. The compound according to claim 7

wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl,
20 tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, octahydro-indoliziny, octahydro-
quinoliziny or aza-bicyclo[3.2.1]octanyl, each ring being optionally substituted with one
or more R_5 ;

R_1 is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl,
25 naphthyl, C1-3alkylsulfonylC1-3alkyl, C3-6cycloalkylsulfonylC1-3alkyl,
arylsulfonylC1-3alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl,
piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, isoxazolyl,
pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl,
benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R_1 is optionally
30 substituted by one or more R_a ;

R_a is a bond, C1-3 alkyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl
5 wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

10 or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

15 or R_a is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

20 or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b ;

R_b is C1-3 alkyl, C3-6 cycloalkyl, aryl, C1-3 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or
25 guanidino;

R_2 is hydrogen or methyl;

R_3 is a bond, hydrogen, C1-5 alkyl, C2-5alkylene, C4-6 cycloalkyl or arylC1-2alkyl
30 wherein R_3 is optionally substituted by one or more R_c ;

R_c is C1-4 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, indolinyl, furanyl, tetrahydrofuranlyl, pyranlyl, tetrahydropyranlyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranlyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-4 alkoxy, phenoxy, naphthyloxy, C1-3 alkanoyl, benzoyl, C1-3 alkoxycarbonyl, phenoxycarbonyl, C1-3 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, or R_c is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or aryl, or R_c is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-3 alkyl, C3-6 cycloalkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, phenylC1-3alkoxy, benzoyl, halogen, hydroxy, oxo or cyano;

R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-6 alkyl, C1-6alkoxyC1-6alkyl, C1-6alkylaminoC1-6alkyl, C1-6alkylthioC1-6alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-6 alkoxy, , phenoxy, naphthyloxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl,

indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl and benzoxazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-4alkanoyloxy, benzyloxy, C1-4 alkoxycarbonyl, arylC1-2alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-4 alkanoylamino, aroylamino, C1-4 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-4 alkoxycarbonylamino, phenoxycarbonylamino, C1-4 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-4 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-4 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R₆;

R₆ is C1-4 alkyl, C1-4 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-4 alkanoyl, aroyl, C1-4alkanoyloxy, phenoxy, naphthyloxy, benzyloxy, C1-4

alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl,
5 benzimidazolyl, or benzthiazolyl,

or R_c is C1-4 alkanoylamino, benzoylamino, C1-4 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl,
10 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_c is C1-4 alkoxycarbonylamino, phenoxycarbonylamino, C1-4 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-4 alkylsulfonylamino, phenylsulfonylamino, C1-4 alkylaminosulfonyl, phenylaminosulfonyl, amino
15 wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

20 or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_c may be further optionally substituted by one or more R_f,

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, methoxy, cyclopropyl, phenyl, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy or
25 carboxamide.

9. The compound according to claim 8

30 wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R₅;

5

R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

10

15

R_a is a bond, C1-3 alkyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

20

25

30

R_b is methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, methoxy, ethoxy, n-propoxy, i-propoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₂ is hydrogen;

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, C1-3 alkoxy, phenoxy, naphthyloxy, C1-2 alkanoyl, benzoyl, C1-2 alkoxy carbonyl, phenoxy carbonyl, C1-2alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or aryl, or R_c is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or aryl, or R_c is C1-2 alkoxy carbonylamino, phenoxy carbonylamino, C1-2 alkyl carbamoyloxy, aryl carbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, or R_c is halogen, hydroxy, oxo, carboxy or cyano, R_c may be further optionally substituted by one or more R_d;

R_d is methyl, cyclopropyl, cyclohexyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, fluoro, chloro, oxo or cyano;

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-3alkanoyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzyloxycarbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-3 alkoxycarbonylamino, phenoxy carbonylamino, C1-3 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C1-3 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl,

thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-3 alkanoyl, aroyl, C1-3alkanoyloxy, phenoxy, benzyloxy, C1-3 alkoxy carbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl

wherein the nitrogen atom may be independently mono or di-substituted by C1-3

alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl,

thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur

atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen

atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl,

piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl,

pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-3 alkoxy carbonylamino, phenoxycarbonylamino, C1-3

alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino,

phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino

wherein the nitrogen atom may be independently mono or di-substituted by C1-3

alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,

oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or

benzthiazolyl,

or R_e is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R_e may be further optionally substituted by one or more R_f ,

and

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide.

10. The compound according to claim 9

wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl or tetrahydropyranyl each ring being substituted with one or more R_5 ;

5 Y is C(O);

R_1 is a bond, methyl, ethyl, i-propyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl
10 or amino; wherein R_1 is optionally substituted by one or more R_a ;

R_a is a bond, methyl, ethyl, cyclopropyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, acetyl, acetoxyl, phenoxy, benzyloxy, methoxycarbonyl, phenoxycarbonyl, benzoyloxy,
15 carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
or R_a is acetylamino, benzoylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,
20 or R_a is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
or R_a is fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or
25 carboxamide, R_a may be further optionally substituted by one or more R_b ;

R_b is methyl, cyclopropyl, phenyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

30 R_3 is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R_3 is optionally substituted by one or more R_c ;

R_c is methyl, ethyl, n-propyl, i-propyl, C5-6 cycloalkyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, thienyl, oxazolyl, thiazolyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or aryl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or aryl,

or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_c is fluoro, chloro or oxo, R_c may be further optionally substituted by one or more R_d ;

R_d is methyl, cyclopropyl, phenyl, methoxy, fluoro, chloro or oxo;

R_5 is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-4alkoxyC1-4alkyl, C1-4alkylaminoC1-4alkyl, C1-4alkylthioC1-4alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-4 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocycloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-2alkanoyl, benzoyl, naphthoyl, C1-2alkanoyloxy, benzyloxy, C1-2 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be

independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C1-2 alkoxy, C3-6 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-2 alkanoyl, aroyl, C1-2 alkanoyloxy, phenoxy, benzyloxy, C1-2 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R_e is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R_e is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-2 alkylsulfonylamino,

phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-2
alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,
oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
5 or R_e is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R_e may be
further optionally substituted by one or more R_f,
and

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro,
chloro, hydroxy, oxo, carboxy or carboxamide.

11. The compound according to claim 10

wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl, azepan-3-yl, azepan-4-
yl or tetrahydropyran-4-yl, each ring being optionally substituted with one or more R₅;

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, cyclopropyl, cyclohexyl, phenoxy, phenyl,
benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,
20 furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is
optionally substituted by one or more R_a;

R_a is methyl, phenyl, thienyl, methoxy, acetyl, acetoxy, phenoxy, benzyloxy,
methoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be
independently mono or di-substituted by methyl or phenyl,
or R_a is acetylamino, methylthio, phenylthio wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
independently substituted by methyl or phenyl,
30 or R_a is methoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy,
methylsulfonylamino, phenylsulfonylamino, amino wherein the nitrogen atom
may be independently mono or di-substituted by methyl or phenyl,

or R_a is fluoro, chloro, hydroxy, oxo, carboxy, cyano or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, propenyl, butenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

5

R_c is methyl, ethyl, n-propyl, i-propyl, cyclohexyl, cyclopentyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxyl, benzoyloxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro, chloro or oxo;

10

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-2alkoxyC1-2alkyl, C1-2alkylaminoC1-2alkyl, C1-2alkylthioC1-2alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-2 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocycloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

20

or R₅ is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

25

or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

30

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is methyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidiny, morpholiny, indolyl, thienyl, pyridiny, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_e is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R_e is methoxycarbonylamino, ethoxycarbonylamino, phenoxy carbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f,

and

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo.

12. The compound according to claim 11

wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl or tetrahydropyran-4-yl, each ring being substituted with one or more R₅;

R₁ is i-propyl, benzyloxy, cyclohexyl, phenyl, 4-(acetylamino)-phenyl, 4-(methanesulfonylamino)-phenyl, 4-methoxyphenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-chlorophenyl, naphthyl, thienylmethyl, piperidinyl, morpholinyl, pyrrolidinyl, piperazinyl, furanyl, thienyl, 5-chlorothienyl,
5 pyridin-4-yl, pyrazinyl, methylamino, ethylamino, dimethylamino or diethylamino;

R₃ is ethyl, n-propyl, propenyl, butenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is methyl, cyclohexyl, cyclopentyl, indanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro or chloro;

R₅ is a bond, carbonyl, methyl, ethyl, n-propyl, n-butyl, t-butyl, i-propyl, i-butyl,
15 cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, piperidinyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy, R₅ may be further optionally substituted by one or more R_c;

20 R_c is methyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, thienyl, 5-methylthienyl, methoxy, phenoxy, benzyloxy, piperidinyl, pyridinyl, indolyl, 1-(tolyl-sulfonyl)-indolyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, phenyl or benzyl,
or R_c is hydroxy, fluoro, chloro, oxo, dimethylamino or trifluoromethyl;

25

13. The compound according to claim 12

30 wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl or azetidin-3-yl, each ring being substituted with one or more R₅;

R₁ is phenyl, 4-(acetylamino)-phenyl, 4-(methanesulfonylamino)-phenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-fluorophenyl, thienylmethyl, morpholinyl, pyrrolidinyl, piperidinyl, piperazinyl, 5-chlorothienyl, pyridin-4-yl or pyrazinyl;

R₃ is n-butyl, i-butyl, 2,2-dimethylpropyl, cyclohexylmethyl, propenyl, i-butenyl, 4-methoxybenzyl, 4-chlorobenzyl, 3,4-dichlorobenzyl, 3-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylbenzyl, 3-methylbenzyl or naphth-2-ylmethyl; wherein the configuration at the stereocenter defined by R₂ and R₃ when they are different and the carbon they are attached to is defined as L;

and

R₅ is a bond, methyl, ethyl, n-propyl, n-butyl, n-pentyl, 2-pentyl, 3-pentyl, phenethyl, phenpropyl, 2,2-dimethylpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, phenyl, benzyl, naphthylmethyl, indanylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, 5-methylthienylmethyl, piperidinyl, piperidinylcarbonyl, pyridinylcarbonyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, methylcarbamoyl, phenylcarbamoyl, benzylcarbamoyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, methylcyclohexyl, methylbenzyl, methoxybenzyl, phenoxybenzyl, benzyloxybenzyl, N-[(4-methylphenyl)-sulfonyl]-indolylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, N,N-dimethylaminoacetyl, trifluoromethylbenzyl, fluoro, oxo or carboxy.

14. The compound according to claim 8

wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R_5 ;

- 5 R_1 is a bond, C1-4 alkyl, C1-4 alkoxy, cyclopropyl, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R_1 is optionally substituted by one or
10 more R_a ;

- R_a is methyl, ethyl, propyl, i-propyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, ethoxy, acetyl, acetoxy, phenoxy, naphthyloxy, benzyloxy,
15 methoxycarbonyl, ethoxycarbonyl, phenoxycarbonyl, naphthyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
or R_a is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be
20 oxidized to a sulfoxide or sulfone, ethylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
25 or R_a is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, naphthylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, naphthylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, naphthylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl,
30 phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

5 R_b is methyl, ethyl, cyclopropyl, cyclohexyl, phenyl, methoxy, ethoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₂ is hydrogen or methyl;

10 R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

15 R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, naphthyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

20 or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,

25 or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

or R_c is chloro, fluoro, hydroxy, oxo, carboxy or cyano;

30 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl,

tetrahydrofuranyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl or tetrahydrothiophenyl;

R₄ is hydrogen;

5

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, naphthyloxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-

10

15 or R₅ is acetylamino, benzoylamino, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, phenylsulfonylamino, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

20

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_c;

25

R_c is methyl ethyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidinyl, morpholinyl, indolyl, thienyl, pyridinyl, methoxy, ethoxy, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

30

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio methylthio wherein the sulfur atom

may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,
or R_e is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino,
5 phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f ;

10 R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo.

15 15. The compound according to claim 14

wherein:

R_1 is a bond, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, benzyloxy, cyclopropyl, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl,
20 morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R_1 is optionally substituted by one or more R_a ;

25 R_a is methyl, cyclopropyl, phenyl, halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R_3 is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl, benzyl or naphthylmethyl wherein R_3 is optionally substituted by one or more
30 R_c ;

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

5 or R_c is acetylamino, benzoylamino, methylthio, methoxycarbonylamino, methylcarbamoyloxy, methylsulfonylamino, methylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl,

or R_c is fluoro or oxo;

10 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, pyrrolidinyl or piperidinyl;

15 R₅ is methyl, ethyl, n-propyl, n-butyl, n-pentyl, 2-pentyl, 3-pentyl, phenethyl, phenpropyl, 2,2-dimethylpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methylbenzyl, 3-methylbenzyl, 4-methylbenzyl, 2,6-dimethylbenzyl, 2,5-dimethylbenzyl, 2,4-dimethylbenzyl, 2,3-dimethylbenzyl, 3,4-dimethylbenzyl, 3,5-dimethylbenzyl, 2,4,6-
20 trimethylbenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-phenoxybenzyl, 3-phenoxybenzyl, 4-phenoxybenzyl, 2-benzyloxybenzyl, 3-benzyloxybenzyl, 4-benzyloxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 2,6-difluorobenzyl, 2,5-difluorobenzyl, 2,4-difluorobenzyl, 2,3-difluorobenzyl, 3,4-difluorobenzyl, 3,5-difluorobenzyl, 2,4,6-trifluorobenzyl, 2-trifluoromethylbenzyl, 3-
25 trifluoromethylbenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, indanylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, 5-methylthienylmethyl, piperidinyl, piperidinylcarbonyl, pyridinylcarbonyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, methylcarbamoyl, phenylcarbamoyl, benzylcarbamoyl, methylsulfonylamino, phenylsulfonylamino,
30 methylamino, dimethylamino, fluoro, oxo or carboxy.

16. The compound according to claim 15

wherein:

5 R_1 is methoxy, benzyloxy, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R_1 is optionally substituted by one or more R_a ;

10

R_a is methyl, phenyl, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R_3 is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl or benzyl wherein R_3 is optionally substituted by one or more R_c ;

15

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, acetylamino, methylthio, methylsulfonylamino or fluoro;

20 R_2 and R_3 together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl or tetrahydrofuranyl;

25 R_5 is methyl, ethyl, n-propyl, n-butyl, phenethyl, phenpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclohexyl, cyclopropylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl 4-fluorobenzyl, 3,5-difluorobenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, phenylcarbamoyl, phenylsulfonylamino or fluoro.

30

17. The compound according to claim 16

wherein:

Het is pyrrolidinyl, piperidinyl or tetrahydropyranyl;

5

R₁ is benzyloxy, phenoxy, naphthyloxy, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyridinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or phenylamino;

10 R₃ is n-propyl, i-butyl, propenyl, i-butenyl or 2,2-dimethylpropyl;

R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, or cycloheptyl;

15 R₅ is methyl, ethyl, n-propyl, phenethyl, t-butyl, i-propyl, i-butyl, cyclohexyl, cyclohexylmethyl, benzyl, 4-fluorobenzyl, naphthylmethyl, acetyl, benzoyl or benzyloxycarbonyl.

20 18. A compound selected from the group consisting of:

Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

25 4-Acetylamino-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

4-Acetylamino-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

30

Morpholine-4-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

35

Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide hydrochloride;

5 Morpholine-4-carboxylic acid {1-[4-cyano-1-(1-methyl-ethyl)-piperidin-4-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-phenethyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

10 Morpholine-4-carboxylic acid {1-[3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

15 Morpholine-4-carboxylic acid [1-(4-cyano-1-isopropyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

20 Morpholine-4-carboxylic acid [1-(1-phenethyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(1-n-propyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

25 Morpholine-4-carboxylic acid [1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

4-Acetylamino-*N*-[1-(1-benzyl-4-cyano-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

30 4-Acetylamino-*N*-[1-(4-cyano-1-isopropyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

35 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

40 4-(Acetylamino-methyl)-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-isonicotinamide;

45 Pyrazine-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

5-Chloro-thiophene-2-carboxylic acid [1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

4-Chloro-*N*-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-3-phenoxy-benzamide;

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl-ethyl]-isonicotinamide;

Pyrazine-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(cyclohexyl-methyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

Pyrazine-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-methyl-ethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

N-[1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide;

N-[1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-methanesulfonylamino-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethyl]-4-fluoro-benzamide;

N-[1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-4-methanesulfonylamino-benzamide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-ethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

5 Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(3-methyl-benzyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

10 Morpholine-4-carboxylic acid {1-[3-cyano-1-(2-methyl-pent-2-enyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid {1-[3-cyano-1-(1*H*-indol-3-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

15 Pyrrolidine-1-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

20 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

25 Morpholine-4-carboxylic acid [1-(3-cyano-1-isopropyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

30 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

35 Morpholine-4-carboxylic acid {1-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-phenethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

40 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopropylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-methyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

45

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-azetidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

5 Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-propyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

10 Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

15 Morpholine-4-carboxylic acid [1-(3-cyano-1-isobutyl-piperidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

20 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

25 Morpholine-4-carboxylic acid {1-[3-cyano-1-(*trans*-4-methyl-cyclohexyl)-pyrrolidin-3-ylcarbamoyl]-3,3-dimethyl-butyl}-amide;

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-naphthalen-2-yl-ethyl]-amide;

30 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-2-(4-chloro-phenyl)-ethyl]-amide;

35 Morpholine-4-carboxylic acid {1-[3-cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethyl}-amide;

Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-but-3-enyl]-amide;

40 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclopentylmethyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexyl-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

45

4-Chloro-*N*-[1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-benzamide;

5 Pyrazine-2-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

4,4-dimethyl-2-(2-thiophen-2-yl-acetylamino)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide;

10 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-cyclohexyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide;

15 Morpholine-4-carboxylic acid [2-(4-chloro-phenyl)-1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-ethyl]-amide;

20 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-(3,4-dichloro-phenyl)-ethyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-2-naphthalen-2-yl-ethyl]-amide;

25 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-amide;

Morpholine-4-carboxylic acid [1-(4-cyano-1,2-dimethyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butyl]-amide

30 and the pharmaceutically acceptable derivatives thereof.

35 19. The compound according to claim 14 wherein the compound is selected from:

[1-(1-Benzyl-4-cyano-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;

40 [1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-cyclohexyl]-carbamic acid *t*-butyl ester;

[1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;

45 [1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-cyclohexyl]-carbamic acid benzyl ester;

Naphthalene-2-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

- 5 Morpholine-4-carboxylic acid [1-(4-cyano-1-propyl-piperidin-4-ylcarbamoyl)-3-methyl-butyl]-amide;

Naphthalene-2-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;

- 10 [1-(1-Benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;

- 15 Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;


[1-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-carbamic acid benzyl ester;

- 20 Morpholine-4-carboxylic acid [1-(3-cyano-1-cyclohexylmethyl-pyrrolidin-3-ylcarbamoyl)-3-methyl-butyl]-amide;


Morpholine-4-carboxylic acid [1-(1-benzyl-3-cyano-pyrrolidin-3-ylcarbamoyl)-3-methyl-but-3-enyl]-amide and

- 25 and the pharmaceutically acceptable derivatives thereof.

- 30 20. A compound of formula (Ia) or (Ib):



(Ia),



(Ib)

wherein:

Het is

azepanyl, piperidinyl, pyrrolidinyl, azetidiny, oxepanyl, tetrahydropyranyl,
 tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, azocanyl, oxocanyl, 1,3-diazocanyl,
 1,4-diazocanyl, 1,5-diazocanyl, 1,3-dioxocanyl, 1,4-dioxocanyl, 1,5-dioxocanyl, 1,3-
 5 oxazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,3-diazepanyl, 1,4-diazepanyl, 1,3-
 dioxepanyl, 1,4-dioxepanyl, 1,3-oxazepanyl, 1,4-oxazepanyl, 1,2-thiazocanyl-1,1-
 dioxide, 1,2,8-thiadiazocanyl-1,1-dioxide, 1,2-thiazepanyl-1,1-dioxide, 1,2,7-
 thiadiazepanyl-1,1-dioxide, tetrahydrothiophenyl, hexahydropyrimidinyl,
 hexahydropyridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, pyrazolidinyl, dihydro-
 10 oxazolyl, dihydrothiazolyl, dihydroimidazolyl, isoxazoliny, oxazolidinyl, 1,2-
 thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide,
 imidazolidinyl-2,4-dione, imidazolidinyl, morpholinyl, dioxanyl, tetrahydropyridinyl,
 thiomorpholinyl, thiazolidinyl, dihydropyranyl, dithianyl, decahydro-quinolinyl,
 decahydro-isoquinolinyl, 1,2,3,4-tetrahydro-quinolinyl, indolinyl, octahydro-quinoliziny,
 15 dihydro-indoliziny, octahydro-indoliziny, octahydro-indolyl, decahydroquinazoliny,
 decahydroquinoxaliny, 1,2,3,4-tetrahydroquinazoliny or 1,2,3,4-tetrahydroquinoxaliny;

A C6-C10 bridged bicyclo wherein one or more carbon atoms are optionally replaced by
 a heteroatom chosen from N, O and S;

each being optionally substituted with one or more R₅;

R₁ is a bond, hydrogen, C1-10 alkyl, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, C3-8
 25 cycloalkyloxy, aryl, benzyl, tetrahydronaphthyl, indenyl, indanyl, C1-10alkylsulfonylC1-
 10alkyl, C3-8cycloalkylsulfonylC1-10alkyl, arylsulfonylC1-10alkyl, heterocyclyl
 selected from azepanyl, azocanyl, pyrrolidinyl, piperidinyl, morpholinyl,
 thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl,
 tetrahydrothiopyranyl, thiopyranyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, oxazolyl,
 30 isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl,
 tetrazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl,

benzisoxazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl, tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocycloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, hydroxy or amino; wherein R₁ is optionally substituted by one or more R_a;

5

R_a is a bond, C1-10 alkyl, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-10 alkoxy, C1-10alkanoyl, C1-10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_a is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,

30

indoliny, furanyl, thienyl, pyrroly, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny,

5 or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b ; with the proviso that R_1 and R_a simultaneously cannot be a bond;

R_b is a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more carbon
10 atoms are optionally replaced by O, N, S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2 oxo groups, -NH₂, or one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indoliny, furanyl, thienyl, pyrroly, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl,
15 pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny;

or R_b is C3-6 cycloalkyl, aryl, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, mono-C1-5alkylamino, di-C1-5alkylamino,
20 carboxamide, amidino or guanidino;

R_2 is hydrogen or C1-3 alkyl;

R_3 is a bond, hydrogen, C1-10 alkyl, C2-10alkylene, C3-8 cycloalkyl, arylC1-5alkyl or
25 aryl wherein R_3 is optionally substituted by one or more R_c ;

R_c is C1-10 alkyl, C3-8 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, decahydronaphthyl,
30 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indoliny, furanyl, tetrahydrofuranyl, pyranyl, tetrahydropyranyl, tetrahydrothiopyranyl,

thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, dihydrobenzofuranyl, octohydrobenzofuranyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, tetrahydroquinoliny, quinoliny, tetrahydroisoquinoliny, isoquinoliny, quinazoliny, quinoxaliny, C1-10 alkoxy, aryloxy, C1-10 alkanoyl, aroyl, C1-10 alkoxycarbonyl, aryloxycarbonyl, C1-10 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, indoliny, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny, or R_c is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, indoliny, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny, or R_c is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, indoliny, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny, or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-5alkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino;

- 5 R_2 and R_3 together with the carbon they are attached optionally form a nonaromatic 5-7 membered cycloalkyl or heterocyclic ring;

each R_4 is independently hydrogen, hydroxy or C1-3 alkyl;

- 10 R_5 is a bond, hydrogen, carbonyl, C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-10 alkoxy, aryloxy, C3-8 cycloalkyl, aryl, benzyl, tetrahydronaphthyl, indenyl, indanyl, C3-7cycloalkylsulfonylC1-5alkyl, arylsulfonylC1-5alkyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl,
- 15 thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, thiopyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranlyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridizinyl, tetrazolyl, triazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, quinazolinyl,
- 20 tetrahydroquinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, benzyloxy, C1-10alkoxycarbonyl, arylC1-5alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl,
- 25 morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,
- 30 or R_5 is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently

substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny, or R₅ is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxaliny, or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R₅ may be further optionally substituted by one or more R₆;

R₆ is C1-10 alkyl, C1-10alkoxyC1-10alkyl, C1-10alkylaminoC1-10alkyl, C1-10alkylthioC1-10alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-10 alkoxy, C3-8 cycloalkyl, aryl, tetrahydronaphthyl, indenyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, thiopyranyl, tetrahydrothiopyranyl, pyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxaliny, C1-10alkanoyl, aroyl, C1-10alkanoyloxy, aryloxy, benzyloxy, C1-10 alkoxycarbonyl, arylC1-3alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl,

benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

or R_e is C1-10 alkanoylamino, aroylamino, C1-10 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom

5 may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-10 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

10 or R_e is C1-10 alkoxycarbonylamino, aryloxycarbonylamino, C1-10 alkylcarbamoxyloxy, arylcarbamoxyloxy, C1-10 alkylsulfonylamino, arylsulfonylamino, C1-10 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-10 alkyl, 15 aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

20 or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_e may be further optionally substituted by one or more R_f,

R_f is C1-5 alkyl, C3-6 cycloalkyl, tolylsulfonyl, C1-5 alkoxy, aryl, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, 25 carboxamide, amidino or guanidino;

R₆ is

hydrogen, hydroxy, nitrile or

30 a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH,

S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2
oxo groups, -NH₂, one or more C₁₋₄ alkyl, pyrrolidinyl, piperidinyl, morpholinyl,
thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl,
oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
5 benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl,
quinazoliny, benzoxazolyl or quinoxaliny;

wherein R₁ and R₆ in the formulas (I) or (II) optionally form a 4 to 8 membered mono-
or 7-12 membered polycyclo heteroring system, each aromatic or nonaromatic, wherein
10 each heteroring is optionally substituted by one or more R₇;

each R₇ and R₈ are independently:

15 C1-5 alkyl chain optionally interrupted by one or two N, O or S(O)_m and optionally
substituted by 1-2 oxo, amino, hydroxy, halogen, C1-4alkyl, pyrrolidinyl, piperidinyl,
morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl,
thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl,
pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl,
20 isoquinolinyl, quinazoliny, benzoxazolyl or quinoxaliny,

aryl, aryloxy, aroyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, C1-5
alkanoyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, benzyloxycarbonyl,
C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio C1-5 alkylsulfonylamino,
25 arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, C3-6 cycloalkyl and
benzyloxy
each of the aforementioned are optionally halogenated,
halogen, hydroxy, oxo, carboxy, nitrile, nitro or NH₂C(O)-;

30 m is 0, 1 or 2;

X is =O, =S or =N-R₆ wherein R₆ is as defined above, and

pharmaceutically acceptable derivatives thereof.

5

21. The compound according to claim 20

wherein:

10 Het is piperidinyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, azetidiny, azepanyl, oxepanyl, tetrahydrofuranyl, oxetanyl, hexahydropyrimidinyl, hexahydropryridazinyl, piperazinyl, 1,4,5,6-tetrahydropyrimidinyl, octahydro-indoliziny, octahydro-quinoliziny, decahydro-quinoliny, 1,2,3,4-tetrahydro-quinoliny, dihydro-oxazolyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-
15 1,1-dioxide, imidazolidinyl, pyrazolidinyl or a bridged bicyclo chosen from aza-bicyclo[3.2.1]octane, aza-bicyclo[2.2.1]heptane, aza-bicyclo[2.2.2]octane, aza-bicyclo[3.2.2]nonane, aza-bicyclo[2.1.1]hexane, aza-bicyclo[3.1.1]heptane, aza-bicyclo[3.3.2]decane and 2-oxa or 2-thia-5-aza-bicyclo[2.2.1]heptane; each ring being substituted with one or more R₅;

20

R₁ is a bond, hydrogen, C1-7 alkyl, C1-7 alkoxy, C3-7 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, tetrahydronaphthyl, C1-7alkylsulfonylC1-7alkyl, C3-7cycloalkylsulfonylC1-7alkyl, arylsulfonylC1-7alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indoliny, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridiny, isoxazolyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinoliny, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoisoxazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

30

R_a is a bond C1-7 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridiny, pyrimidinyl, pyrazinyl,

indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliny, isoquinoliny, quinazoliny, quinoxaliny, C1-7 alkoxy, C1-7alkanoyl, C1-7alkanoyloxy, aryloxy, benzyloxy, C1-7 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or

5 di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,

10 or R_a is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl,

15 imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,

or R_a is C1-7 alkoxycarbonylamino, aryloxycarbonylamino, C1-7 alkylcarbamoyloxy, arylcarbamoyloxy, C1-7 alkylsulfonylamino, arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein

20 the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,

25 or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or

30 guanidino;

R₂ is hydrogen or methyl or ethyl;

R₃ is a bond, hydrogen, C1-5 alkyl, C2-5alkylene, C3-7 cycloalkyl, arylC1-3alkyl or aryl

5 wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-5 alkyl, C3-7 cycloalkyl, aryl, indanyl, indenyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, 10 piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, tetrahydrofuranyl, pyranyl, tetrahydropyranyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5 15 alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, 20 indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_c is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be 25 independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, 30 or R_c is C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino,

arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl, or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-4 alkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, halogen, hydroxy, oxo or cyano;

R₄ is hydrogen or methyl;

R₅ is a bond, hydrogen, carbonyl, C1-8 alkyl, C1-8alkoxyC1-8alkyl, C1-8alkylaminoC1-8alkyl, C1-8alkylthioC1-8alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-8 alkoxy, aryloxy, C3-7 cycloalkyl, aryl, benzyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl, tetrahydropyranyl, thiopyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, tetrazolyl, triazolyl, pyrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-7alkanoyl, aroyl, C1-7alkanoyloxy, benzyloxy, C1-7 alkoxycarbonyl, arylC1-4alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,

benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,

or R₅ is C1-7 alkanoylamino, aroylamino, C1-7 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a

5 sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny,

10 or R₅ is C1-7 alkoxycarbonylamino, aryloxycarbonylamino, C1-7 alkylcarbamoyloxy, arylcarbamoyloxy, C1-7 alkylsulfonylamino, arylsulfonylamino, C1-7 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-7 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, 15 imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny or quinoxaliny, or R₅ is halogen, hydroxy, oxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R_c;

20

R_c is C1-7 alkyl, C1-7alkoxyC1-7alkyl, C1-7alkylaminoC1-7alkyl, C1-7alkylthioC1-7alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-7 alkoxy, C3-7 cycloalkyl, aryl, tetrahydronaphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholiny, thiomorpholiny, piperazinyl, thiopyranyl, 25 tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliny, isoquinoliny, quinazoliny, quinoxaliny, C1-5 alkanoyl, aroyl, C1-5alkanoyloxy, aryloxy, benzyloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or 30 di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholiny,

thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

5 or R_e is C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

10 or R_e is C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl or quinoxalinyl,

20 or R_e is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_e may be further optionally substituted by one or more R_f,

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, C1-3 alkoxy, cyclopropyl, cyclohexyl, phenyl, naphthyl, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R_g is

hydrogen, hydroxy, nitrile or

30 a C1-6 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH,

S(O), S(O)₂ or S and wherein said chain is optionally independently substituted with 1-2
oxo groups, -NH₂, one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl,
thiomorpholinyl, piperazinyl, indolinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl,
oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
5 benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl,
quinazoliny, benzoxazolyl or quinoxaliny;

R₁ and R₆ of the formula (Ia) or formula (Ib) form a monocyclic 5, 6 or 7 membered
aromatic or nonaromatic heterocyclic ring optionally substituted by R₇;

10

or a bicyclic ring having one 5, 6 or 7 membered aromatic or nonaromatic
heterocyclic ring fused to a second 5-7 membered aromatic or nonaromatic heterocyclic
or carbocyclic ring wherein each ring is optionally independently substituted by one or
more R₇;

15

R₇ and R₈ are independently C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy,
benzyloxy each of the aforementioned are optionally halogenated or R_x is halogen,
hydroxy, oxo, carboxy, nitrile, nitro or NH₂C(O)-;

20 m is 0, 1 or 2 and

X is O or S.

25 22. The compound according to claim 21 wherein

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl,
tetrahydrothiopyranyl, tetrahydrofuranyl, oxetanyl, octahydro-indoliziny, octahydro-
30 quinoliziny or aza-bicyclo[3.2.1]octanyl, each ring being optionally substituted with one
or more R₅;

R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, C1-3alkylsulfonylC1-3alkyl, C3-6cycloalkylsulfonylC1-3alkyl, arylsulfonylC1-3alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, isoxazolyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond, C1-3 alkyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl, or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is C1-3 alkyl, C3-6 cycloalkyl, aryl, C1-3 alkoxy, aryloxy, benzyloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino;

5 R₂ is hydrogen or methyl;

R₃ is a bond, hydrogen, C1-5 alkyl, C2-5alkylene, C4-6 cycloalkyl or arylC1-2alkyl wherein R₃ is optionally substituted by one or more R_c;

10 R_c is C1-4 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, indolinyl, furanyl, tetrahydrofuran, pyran, tetrahydropyran, thienyl, pyrrolyl, 15 oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, C1-4 alkoxy, phenoxy, naphthyloxy, C1-3 alkanoyl, benzoyl, C1-3 alkoxy-carbonyl, phenoxycarbonyl, C1-3 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be 20 independently mono or di-substituted by C1-5 alkyl or aryl, or R_c is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or aryl, 25 or R_c is C1-3 alkoxy-carbonylamino, aryloxy-carbonylamino, C1-3 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, 30 or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino or guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is C1-3 alkyl, C3-6 cycloalkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, phenylC1-3alkoxy, benzoyl, halogen, hydroxy, oxo or cyano;

5 R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-6 alkyl, C1-6alkoxyC1-6alkyl, C1-6alkylaminoC1-6alkyl, C1-6alkylthioC1-6alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-6 alkoxy, , phenoxy, naphthyloxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl,
 10 indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, tetrahydrothiopyranyl, furanyl, tetrahydrofuranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl and benzoxazolyl, heterocyclyloxy wherein the heterocyclyl moiety is
 15 selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-4alkanoyloxy, benzyloxy, C1-4 alkoxycarbonyl, arylC1-2alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
 20 or R₅ is C1-4 alkanoylamino, aroylamino, C1-4 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl,
 25 benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-4 alkoxycarbonylamino, phenoxycarbonylamino, C1-4 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-4 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-4 alkyl, aryl, pyrrolidinyl, piperidinyl,
 30 morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R₅ is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R₅ may be further optionally substituted by one or more R_c;

R_c is C1-4 alkyl, C1-4 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl,
5 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,
tetrahydrothiopyranyl, tetrahydropyranyl, tetrahydrofuranyl, thienyl, oxazolyl,
thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl,
benzthiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl,
C1-4 alkanoyl, aroyl, C1-4alkanoyloxy, phenoxy, naphthyloxy, benzyloxy, C1-4
10 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen
atom may be independently mono or di-substituted by C1-3 alkyl, phenyl,
naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,
furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl,
benzimidazolyl, or benzthiazolyl,

15 or R_c is C1-4 alkanoylamino, benzoylamino, C1-4 alkylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen
atom may be independently substituted by C1-3 alkyl, phenyl, naphthyl,
pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl,
20 thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or
benzthiazolyl,

or R_c is C1-4 alkoxycarbonylamino, phenoxycarbonylamino, C1-4
alkylcarbamoyloxy, phenylcarbamoyloxy, C1-4 alkylsulfonylamino,
phenylsulfonylamino, C1-4 alkylaminosulfonyl, phenylaminosulfonyl, amino
25 wherein the nitrogen atom may be independently mono or di-substituted by C1-3
alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl,
piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl,
pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_c is halogen, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_c may be
30 further optionally substituted by one or more R_d;

R_f is methyl, ethyl, t-butyl, tolylsulfonyl, methoxy, cyclopropyl, phenyl, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide.

- 5 R₁ and R₆ of the formula (Ia) or Formula (Ib) optionally form a monocyclic 5 or 6 membered aromatic or nonaromatic heterocyclic ring optionally substituted by R₇;

or a bicyclic ring having one 5, 6 or 7 membered aromatic or nonaromatic heterocyclic ring fused to a second 5-6 membered aromatic or nonaromatic heterocyclic
10 or carbocyclic ring wherein each ring is optionally independently substituted by one or more R₇;

R₇ and R₈ are independently C1-4 alkyl, C5-6 cycloalkyl, C1-4 alkoxy, halogen, hydroxy, oxo, carboxy, nitrile, nitro or NH₂C(O)-;

- 15 and
X is O.

23. The compound according to claim 22 wherein:

- 20 Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R₅;

- 25 R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

- 30 R_a is a bond, C1-3 alkyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, C1-3

alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3
alkoxycarbonyl, aryloxycarbonyl, aroyloxy, carbamoyl wherein the nitrogen atom
may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl,
piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

5 or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom
may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may
be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl,
morpholinyl, thiomorpholinyl or piperazinyl,

10 or R_a is C1-3 alkoxycarbonylamino, aryloxycarbonylamino, C1-3
alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino,
arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein
the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl,
aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

15 or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or
guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclopentyl,
cyclohexyl, phenyl, methoxy, ethoxy, n-propoxy, i-propoxy, phenoxy,
20 benzyloxy, fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano,
nitro or carboxamide;

R₂ is hydrogen;

25 R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl
wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl,
bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl,
30 bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-
tetrahydronaphthyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl,

imidazolyl, pyrimidinyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, C1-3 alkoxy, phenoxy, naphthyloxy, C1-2 alkanoyl, benzoyl, C1-2 alkoxycarbonyl, phenoxycarbonyl, C1-2alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or aryl,
5 or R_c is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or aryl,
or R_c is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2
10 alkylcarbamoyloxy, arylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl,
or R_c is halogen, hydroxy, oxo, carboxy or cyano, R_c may be further optionally
15 substituted by one or more R_d;

R_d is methyl, cyclopropyl, cyclohexyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, fluoro, chloro, oxo or cyano;

20 R_s is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl,
25 benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-3alkanoyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl,
30 morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R₅ is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl,

or R₅ is C1-3 alkoxycarbonylamino, phenoxycarbonylamino, C1-3 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R₅ is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C1-3 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-3 alkanoyl, aroyl, C1-3 alkanoyloxy, phenoxy, benzyloxy, C1-3 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R_e is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_e is C1-3 alkoxycarbonylamino, phenoxycarbonylamino, C1-3 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino,

phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-3
alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl,
oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or
5 benzthiazolyl,

or R_e is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R_e may be further
optionally substituted by one or more R_f;

and

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, bromo,
10 hydroxy, oxo, carboxy or carboxamide;

R₁ and R₆ of the formula (Ia) or Formula (Ib) form a bicyclic ring having one 5 or
6 membered aromatic or nonaromatic heterocyclic ring fused to a second
5-6 membered heteroaryl, heterocycle or phenyl ring;

15 wherein each ring is optionally independently substituted by one or two R₇.

24. The compound according to claim 23 wherein:

20 Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl or tetrahydropyranyl each ring being
substituted with one or more R₅;

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, ethoxy, cyclopropyl, cyclopentyl,
cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl,
25 thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl
or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is a bond, methyl, ethyl, cyclopropyl, phenyl, pyrrolidinyl, piperidinyl,
morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, acetyl,
30 acetoxyl, phenoxy, benzyloxy, methoxycarbonyl, phenoxycarbonyl, benzoyloxy,
carbamoyl wherein the nitrogen atom may be independently mono or di-
substituted by methyl, ethyl or phenyl,

or R_a is acetylamino, benzoylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,
or R_a is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoxyloxy,
5 phenylcarbamoxyloxy, methylsulfonylamino, phenylsulfonylamino,
methyaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
or R_a is fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_a may be further optionally substituted by one or more R_b;

10 R_b is methyl, cyclopropyl, phenyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl
15 wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, i-propyl, C5-6 cycloalkyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-
20 tetrahydronaphthyl, thienyl, oxazolyl, thiazolyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or aryl,

25 or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or aryl,

or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoxyloxy,
30 phenylcarbamoxyloxy, methylsulfonylamino, phenylsulfonylamino,

methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl, or R_c is fluoro, chloro or oxo, R_c may be further optionally substituted by one or more R_d ;

5

R_d is methyl, cyclopropyl, phenyl, methoxy, fluoro, chloro or oxo;

R_5 is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-4alkoxyC1-4alkyl, C1-4alkylaminoC1-4alkyl, C1-4alkylthioC1-4alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-4 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-2alkanoyl, benzoyl, naphthoyl, C1-2alkanoyloxy, benzyloxy, C1-2 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R_5 is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R_5 is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R_5 is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R_5 may be further optionally substituted by one or more R_e ;

R_e is C1-3 alkyl, C1-2 alkoxy, C3-6 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-2 alkanoyl, aroyl, C1-2alkanoyloxy, phenoxy, benzyloxy, C1-2 alkoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl
5 wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
or R_e is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
10 or R_e is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
15 or R_e is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f ,

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide and
25

R_1 and R_6 of the formula (Ia) or Formula (Ib) form a bicyclic ring having one 5-6 membered aromatic or nonaromatic heterocyclic ring fused to a phenyl ring;
30 wherein each ring is optionally independently substituted by one or two R_7 .

25. The compound according to claim 24 wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl, azepan-3-yl, azepan-4-yl or tetrahydropyran-4-yl, each ring being optionally substituted with one or more R₅;

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, cyclopropyl, cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is methyl, phenyl, thienyl, methoxy, acetyl, acetoxy, phenoxy, benzyloxy, methoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

or R_a is acetylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,

or R_a is methoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

or R_a is fluoro, chloro, hydroxy, oxo, carboxy, cyano or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, propenyl, butenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, i-propyl, cyclohexyl, cyclopentyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-

tetrahydronaphthyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro, chloro or oxo;

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-2alkoxyC1-2alkyl, C1-2alkylaminoC1-2alkyl, C1-2alkylthioC1-2alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-2 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl,
5 heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or
10 phenyl,
or R₅ is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,
15 or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further
20 optionally substituted by one or more R_c;

R_c is methyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidinyl, morpholinyl, indolyl, thienyl, pyridinyl, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl,
25 carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
30 independently substituted by methyl, ethyl or phenyl,


or R_e is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f ;

and

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo;

R_1 and R_6 of the formula (Ia) or Formula (Ib) form the bicyclic ring



; wherein W is $-S(O)_n-$, $-O-C(O)-$ or $-N-C(O)-$, n is 0, 1 or 2 and wherein each ring is optionally independently substituted by one or two R_7 .

26. The compound according to claim 25 wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl or tetrahydropyran-4-yl, each ring being substituted with one or more R_5 ;

R_1 is i-propyl, benzyloxy, cyclohexyl, phenyl, 4-(acetylamino)-phenyl, 4-(methanesulfonylamino)-phenyl, 4-methoxyphenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-chlorophenyl, naphthyl, thienylmethyl,

piperidinyl, morpholinyl, pyrrolidinyl, piperazinyl, furanyl, thienyl, 5-chlorothieryl, pyridin-4-yl, pyrazinyl, methylamino, ethylamino, dimethylamino or diethylamino;

R₃ is ethyl, n-propyl, propenyl, butenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is

5 optionally substituted by one or more R_c;

R_c is methyl, cyclohexyl, cyclopentyl, indanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro or chloro;

10

R₅ is a bond, carbonyl, methyl, ethyl, n-propyl, n-butyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, piperidinyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy, R₅ may be

15 further optionally substituted by one or more R_c;

R_e is methyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, thienyl, 5-methylthienyl, methoxy, phenoxy, benzyloxy, piperidinyl, pyridinyl, indolyl, 1-(tolyl-sulfonyl)-indolyl, carbamoyl wherein the nitrogen atom may be

20

independently mono or di-substituted by methyl, phenyl or benzyl,

or R_e is hydroxy, fluoro, chloro, oxo, dimethylamino or trifluoromethyl;

and

25

n is 2.

27. The compound according to claim 20 wherein:

30

R₁ and R₆ remain acyclic;

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more R₅;

- 5 R₁ is a bond, C1-5 alkyl, C1-5 alkoxy, C3-6 cycloalkyl, aryloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

10

R_a is a bond, C1-3 alkyl, cyclopropyl, cyclohexyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, C1-3 alkoxy, C1-3alkanoyl, C1-3alkanoyloxy, aryloxy, benzyloxy, C1-3 alkoxy carbonyl, aryloxy carbonyl, aryloxy, carbamoyl wherein the nitrogen atom
15 may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
or R_a is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
20 independently substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
or R_a is C1-3 alkoxy carbonylamino, aryloxy carbonylamino, C1-3 alkyl carbamoyloxy, aryl carbamoyloxy, C1-3 alkyl sulfonylamino, aryl sulfonylamino, C1-3 alkyl aminosulfonyl, aryl aminosulfonyl, amino wherein
25 the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, aryl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,
or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or guanidino, R_a may be further optionally substituted by one or more R_b;

30

R_b is methyl, ethyl, n-propyl, i-propyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, methoxy, ethoxy, n-propoxy, i-propoxy, phenoxy,

benzyloxy, fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide;

R₂ is hydrogen;

5

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is C1-3 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, indanyl,
10 bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl,
bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-
tetrahydronaphthyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl,
imidazolyl, pyrimidinyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, C1-3
alkoxy, phenoxy, naphthyloxy, C1-2 alkanoyl, benzoyl, C1-2 alkoxy carbonyl,
15 phenoxycarbonyl, C1-2alkanoyloxy, benzyloxy, carbamoyl wherein the nitrogen
atom may be independently mono or di-substituted by C1-3 alkyl or aryl,
or R_c is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur
atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen
20 atom may be independently substituted by C1-3 alkyl or aryl,
or R_c is C1-2 alkoxy carbonylamino, phenoxycarbonylamino, C1-2
alkylcarbamoyloxy, arylcarbamoyloxy, C1-2 alkylsulfonylamino,
phenylsulfonylamino, C1-2alkylaminosulfonyl, phenylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by C1-3
25 alkyl or phenyl,
or R_c is halogen, hydroxy, oxo, carboxy or cyano, R_c may be further optionally
substituted by one or more R_d;

R_d is methyl, cyclopropyl, cyclohexyl, phenyl, benzyl, methoxy, phenoxy,
30 benzyloxy, benzoyl, fluoro, chloro, oxo or cyano;

R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, C3-6 cycloalkyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-3alkanoyl, benzoyl, naphthoyl, C1-3alkanoyloxy, benzyloxy, C1-3 alkoxycarbonyl, benzyloxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R₅ is C1-3 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzofuranyl, benzothienyl, benzimidazolyl or benzthiazolyl, or R₅ is C1-3 alkoxycarbonylamino, phenoxycarbonylamino, C1-3 alkylcarbamoyloxy, phenylcarbamoyloxy, C1-3 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R₅ is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C1-3 alkoxy, C3-7 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl,

benzthiazolyl, benzoxazolyl, C1-3 alkanoyl, aroyl, C1-3alkanoyloxy, phenoxy, benzyloxy, C1-3 alkoxy carbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R_c is C1-3 alkanoylamino, benzoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl, or R_c is C1-3 alkoxy carbonylamino, phenoxy carbonylamino, C1-3 alkyl carbamoyloxy, phenyl carbamoyloxy, C1-3 alkylsulfonylamino, phenylsulfonylamino, C1-3 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,

or R_c is halogen, hydroxy, oxo, carboxy, cyano or carboxamide, R_c may be further optionally substituted by one or more R_f,

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide;

R₆ is

hydroxy, nitrile or

a C1-5 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH, or S(O)₂ and wherein said chain is optionally independently substituted with 1-2 oxo groups, -NH₂, one or more C1-4 alkyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranal, thiopyranal, furanyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl,

benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazoliny, benzoxazolyl or quinoxaliny;
and

5 X is O.

28. The compound according to claim 27 wherein:

10

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

15

R_a is a bond, methyl, ethyl, cyclopropyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl, imidazolyl, methoxy, acetyl, acetox, phenoxy, benzyloxy, methoxycarbonyl, phenoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-

20

substituted by methyl, ethyl or phenyl,

or R_a is acetylamino, benzoylamino, methylthio, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R_a is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino,

25

methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

or R_a is fluoro, chloro, bromo, iodo, hydroxy, oxo, carboxy, cyano, nitro or carboxamide, R_a may be further optionally substituted by one or more R_b;

30

R_b is methyl, cyclopropyl, phenyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R₃ is a bond, C1-3 alkyl, C2-4alkylene, C5-6 cycloalkyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, i-propyl, C5-6 cycloalkyl, indanyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, thienyl, oxazolyl, thiazolyl, indolyl, benzofuranyl, benzothienyl, benzthiazolyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or aryl, or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or aryl, or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl, or R_c is fluoro, chloro or oxo, R_c may be further optionally substituted by one or more R_d;

R_d is methyl, cyclopropyl, phenyl, methoxy, fluoro, chloro or oxo;

R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-4alkoxyC1-4alkyl, C1-4alkylaminoC1-4alkyl, C1-4alkylthioC1-4alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-4 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, benzyl, indanyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl and benzthiazolyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, C1-2alkanoyl, benzoyl, naphthoyl,

C1-2alkanoyloxy, benzyloxy, C1-2 alkoxy carbonyl, benzyloxy carbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
5 or R₅ is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl or benzthiazolyl,
10 or R₅ is C1-2 alkoxy carbonylamino, phenoxy carbonylamino, C1-2 alkyl carbamoyloxy, phenyl carbamoyloxy, C1-2 alkyl sulfonylamino, phenyl sulfonylamino, C1-2 alkyl aminosulfonyl, phenyl aminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
15 or R₅ is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_e;

R_e is C1-3 alkyl, C1-2 alkoxy, C3-6 cycloalkyl, phenyl, naphthyl, indanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl, indolyl,
20 thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, C1-2 alkanoyl, aroyl, C1-2alkanoyloxy, phenoxy, benzyloxy, C1-2 alkoxy carbonyl, phenoxy carbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl,
25 thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,
or R_e is C1-2 alkanoylamino, benzoylamino, C1-2 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-2 alkyl, phenyl, pyrrolidinyl,
30 piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl,

or R_e is C1-2 alkoxycarbonylamino, phenoxycarbonylamino, C1-2 alkylcarbamoxyloxy, phenylcarbamoxyloxy, C1-2 alkylsulfonylamino, phenylsulfonylamino, C1-2 alkylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-2 alkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl or pyrimidinyl, or R_e is fluoro, chloro, bromo, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f;

R_f is methyl, phenyl, tolylsulfonyl, methoxy, phenoxy, benzyloxy, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide and

R_g is

nitrile or

a C1-5 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH, or S(O)₂ and wherein said chain is optionally independently substituted with oxo, -NH₂, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyridinyl, pyrimidinyl or pyrazinyl.

29. The compound according to claim 28 wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl, azepan-3-yl, azepan-4-yl or tetrahydropyran-4-yl, each ring being optionally substituted with one or more R₅;

R₁ is a bond, methyl, ethyl, i-propyl, methoxy, cyclopropyl, cyclohexyl, phenoxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrazinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

R_a is methyl, phenyl, thienyl, methoxy, acetyl, acetoxy, phenoxy, benzyloxy, methoxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
or R_a is acetylamino, methylthio, phenylthio wherein the sulfur atom may be
5 oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,
or R_a is methoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,
10 or R_a is fluoro, chloro, hydroxy, oxo, carboxy, cyano or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, propenyl, butenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

R_c is methyl, ethyl, n-propyl, i-propyl, cyclohexyl, cyclopentyl, indanyl,
15 bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[4.1.0]heptanyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be
20 oxidized to a sulfoxide or sulfone, fluoro, chloro or oxo;

and

wherein the configuration at the stereocenter defined by R₂ and R₃ when they are different and the carbon they are attached to is defined as L; and

25 R₅ is a bond, hydrogen, carbonyl, C1-4 alkyl, C1-2alkoxyC1-2alkyl, C1-2alkylaminoC1-2alkyl, C1-2alkylthioC1-2alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-2 alkoxy, phenoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected
30 from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy,

methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further optionally substituted by one or more R_c;

R_c is methyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidiny, morpholinyl, indolyl, thienyl, pyridinyl, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,

or R_c is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,

or R_c is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_c may be further optionally substituted by one or more R_f;

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or oxo;

R₆ is

nitrile or

- 5 a C1-5 saturated or unsaturated branched or unbranched carbon chain optionally partially or fully halogenated wherein one or more C atoms are optionally replaced by O, NH, or S(O)₂ and wherein said chain is optionally independently substituted with oxo, -NH₂, morpholinyl or piperazinyl.

10

30. The compound according to claim 29 wherein:

Het is piperidin-4-yl, piperidin-3-yl, pyrrolidin-3-yl, azetidin-3-yl or tetrahydropyran-4-yl, each ring being substituted with one or more R₅;

15

R₁ is i-propyl, benzyloxy, cyclohexyl, phenyl, 4-(acetylamino)-phenyl, 4-(methanesulfonylamino)-phenyl, 4-methoxyphenyl, 3-phenoxyphenyl, 4-chlorophenyl, 4-fluorophenyl, 2-fluorophenyl, 2-fluoro-4-chlorophenyl, naphthyl, thienylmethyl, piperidinyl, morpholinyl, pyrrolidinyl, piperazinyl, furanyl, thienyl, 5-chlorothienyl,
20 pyridin-4-yl, pyrazinyl, methylamino, ethylamino, dimethylamino or diethylamino;

R₃ is ethyl, n-propyl, propenyl, butenyl, i-butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

25

R_c is methyl, cyclohexyl, cyclopentyl, indanyl, 1,2,3,4-tetrahydronaphthyl, methoxy, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, fluoro or chloro;

30

R₅ is a bond, carbonyl, methyl, ethyl, n-propyl, n-butyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, benzyl, piperidinyl, tetrahydropyran-yl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, methylsulfonylamino,

phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy, R₅ may be further optionally substituted by one or more R_e;

R_e is methyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, thienyl, 5-methylthienyl, methoxy, phenoxy, benzyloxy, piperidinyl, pyridinyl, indolyl, 1-(tolyl-sulfonyl)-indolyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, phenyl or benzyl, or R_e is hydroxy, fluoro, chloro, oxo, dimethylamino or trifluoromethyl;

and

R₆ is acetyl, C1-3alkylaminocarbonyl or C1-3alkoxycarbonyl.

31. The compound according to claim 30 wherein:

Het is piperidin-4-yl or pyrrolidin-3-yl;

R₁ is morpholin-4-yl, p-fluorophenyl or p-methoxyphenyl;

R₅ is methyl, propyl, n-pentyl or cyclohexyl
and

R₆ is acetyl, ethylaminocarbonyl or ethoxycarbonyl.

32. The compound according to claim 20 wherein:

Het is piperidinyl, pyrrolidinyl, azetidiny, azepanyl, oxepanyl, tetrahydropyranyl, oxetanyl or tetrahydrothiopyranyl each ring being optionally substituted with one or more

R₅;

R₁ is a bond, C1-4 alkyl, C1-4 alkoxy, cyclopropyl, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl,

benzthiazolyl, benzoxazolyl or amino; wherein R_1 is optionally substituted by one or more R_a ;

R_a is methyl, ethyl, propyl, i-propyl, cyclopropyl, cyclohexyl, phenyl,
5 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, thienyl,
imidazolyl, methoxy, ethoxy, acetyl, acetoxyl, phenoxy, naphthyloxy, benzyloxy,
methoxycarbonyl, ethoxycarbonyl, phenoxycarbonyl, naphthyloxycarbonyl,
benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or
di-substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl,
10 morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, ethylthio wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
15 independently substituted by methyl, ethyl, phenyl, naphthyl, pyrrolidinyl,
piperidinyl, morpholinyl, thiomorpholinyl or piperazinyl,

or R_a is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino,
C1-2 alkylcarbamoyloxy, phenylcarbamoyloxy, naphthylcarbamoyloxy, C1-2
alkylsulfonylamino, phenylsulfonylamino, naphthylsulfonylamino, C1-2
20 alkylaminosulfonyl, phenylaminosulfonyl, naphthylaminosulfonyl, amino wherein
the nitrogen atom may be independently mono or di-substituted by methyl, ethyl,
phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl or
piperazinyl,

or R_a is halogen, hydroxy, oxo, carboxy, cyano, nitro, carboxamide, amidino or
25 guanidino, R_a may be further optionally substituted by one or more R_b ;

R_b is methyl, ethyl, cyclopropyl, cyclohexyl, phenyl, methoxy, ethoxy,
phenoxy, benzyloxy, fluoro, chloro, bromo, hydroxy, oxo, carboxy, cyano,
nitro or carboxamide;

30 R_2 is hydrogen or methyl;

R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl, cyclohexyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more R_c;

5

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, naphthyl, bicyclo[3.1.0]hexanyl, bicyclo[1.1.1]pentanyl, cubanyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, methoxy, ethoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, phenoxycarbonyl, acetoxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

10

or R_c is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl or phenyl,

15

or R_c is methoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl or phenyl,

20

or R_c is chloro, fluoro, hydroxy, oxo, carboxy or cyano;

R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl or tetrahydrothiophenyl;

25

R₄ is hydrogen;

R₅ is a bond, hydrogen, carbonyl, C1-5 alkyl, C1-5alkoxyC1-5alkyl, C1-5alkylaminoC1-5alkyl, C1-5alkylthioC1-5alkyl wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-5 alkoxy, phenoxy, naphthyloxy, cyclopropyl, cyclopentyl, cyclohexyl,

30

phenyl, benzyl, heterocyclyl selected from pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, pyridinyl, and pyrimidinyl, heterocyclyloxy wherein the heterocyclyl moiety is selected from those herein described in this paragraph, acetyl, benzoyl, acetyloxy, benzyloxy, methoxycarbonyl, ethoxycarbonyl, benzyloxycarbonyl,

- 5 benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
or R₅ is acetylamino, benzoylamino, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by methyl, ethyl or phenyl,
10 or R₅ is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, phenylsulfonylamino, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
or R₅ is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R₅ may be further
15 optionally substituted by one or more R_e;

R_e is methyl ethyl, methoxy, ethoxy, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, indanyl, piperidinyl, morpholinyl, indolyl, thienyl, pyridinyl, methoxy, ethoxy, acetyl, benzoyl, acetyloxy, phenoxy, benzyloxy,

- 20 methoxycarbonyl, ethoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl or phenyl,
or R_e is acetylamino, benzoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom
25 may be independently substituted by methyl, ethyl or phenyl,
or R_e is methoxycarbonylamino, ethoxycarbonylamino, phenoxycarbonylamino, methylcarbamoyloxy, phenylcarbamoyloxy, methylsulfonylamino, phenylsulfonylamino, methylaminosulfonyl, phenylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, ethyl
30 or phenyl,

or R_e is fluoro, chloro, hydroxy, oxo, carboxy or carboxamide, R_e may be further optionally substituted by one or more R_f;

R_f is methyl, phenyl, tolylsulfonyl, phenoxy, benzyloxy, fluoro, chloro or
5 oxo.

33. The compound according to claim 32 wherein

R₁ is a bond, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, benzyloxy, cyclopropyl,
10 cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl,
morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl,
imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, quinolinyl,
benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino;
wherein R₁ is optionally substituted by one or more R_a;

15 R_a is methyl, cyclopropyl, phenyl, halogen, hydroxy, oxo, carboxy, cyano, nitro or
carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-
20 butenyl, benzyl or naphthylmethyl wherein R₃ is optionally substituted by one or more
R_c;

R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl,
thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl,
25 carbamoyl wherein the nitrogen atom may be independently mono or di-
substituted by methyl or phenyl,

or R_c is acetylamino, benzoylamino, methylthio, methoxycarbonylamino,
methylcarbamoyloxy, methylsulfonylamino, methylaminosulfonyl, amino
wherein the nitrogen atom may be independently mono or di-substituted by
30 methyl,

or R_c is fluoro or oxo;

R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, pyrrolidinyl or piperidinyl;

5

R₅ is methyl, ethyl, n-propyl, n-butyl, n-pentyl, 2-pentyl, 3-pentyl, phenethyl, phenpropyl, 2,2-dimethylpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methylbenzyl, 3-methylbenzyl, 4-methylbenzyl, 2,6-dimethylbenzyl, 2,5-dimethylbenzyl, 2,4-dimethylbenzyl, 2,3-dimethylbenzyl, 3,4-dimethylbenzyl, 3,5-dimethylbenzyl, 2,4,6-trimethylbenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-phenoxybenzyl, 3-phenoxybenzyl, 4-phenoxybenzyl, 2-benzyloxybenzyl, 3-benzyloxybenzyl, 4-benzyloxybenzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-fluorobenzyl, 2,6-difluorobenzyl, 2,5-difluorobenzyl, 2,4-difluorobenzyl, 2,3-difluorobenzyl, 3,4-difluorobenzyl, 3,5-difluorobenzyl, 2,4,6-trifluorobenzyl, 2-trifluoromethylbenzyl, 3-trifluoromethylbenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, indanylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, 5-methylthienylmethyl, piperidinyl, piperidinylcarbonyl, pyridinylcarbonyl, tetrahydropyranyl, pyrimidinyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, methylcarbamoyl, phenylcarbamoyl, benzylcarbamoyl, methylsulfonylamino, phenylsulfonylamino, methylamino, dimethylamino, fluoro, oxo or carboxy.

25

34. The compound according to claim 33 wherein:

R₁ is methoxy, benzyloxy, cyclohexyl, phenoxy, naphthyloxy, phenyl, benzyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, furanyl, thienyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, quinolinyl, benzofuranyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or amino; wherein R₁ is optionally substituted by one or more R_a;

30

R_a is methyl, phenyl, fluoro, chloro, hydroxy, oxo, carboxy or carboxamide;

R₃ is a bond, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, n-pentyl, propenyl, i-butenyl or benzyl wherein R₃ is optionally substituted by one or more R_c;

5 R_c is methyl, ethyl, cyclohexyl, cyclopentyl, phenyl, furanyl, tetrahydropyranyl, thienyl, oxazolyl, thiazolyl, methoxy, phenoxy, acetyl, benzoyl, methoxycarbonyl, acetylamino, methylthio, methylsulfonylamino or fluoro;

10 R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, cycloheptyl, tetrahydropyranyl, tetrahydrothiopyranyl or tetrahydrofuranlyl;

15 R₅ is methyl, ethyl, n-propyl, n-butyl, phenethyl, phenpropyl, t-butyl, i-propyl, i-butyl, cyclopropyl, cyclohexyl, cyclopropylmethyl, cyclohexylmethyl, phenyl, benzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl 4-fluorobenzyl, 3,5-difluorobenzyl, 4-trifluoromethylbenzyl, naphthylmethyl, pyridinylmethyl, indolylmethyl, thienylmethyl, acetyl, benzoyl, ethoxycarbonyl, benzyloxycarbonyl, t-butoxycarbonyl, phenylcarbamoyl, phenylsulfonylamino or fluoro.

20 35. The compound according to claim 34 wherein:

Het is pyrrolidinyl, piperidinyl or tetrahydropyranyl;

25 R₁ is benzyloxy, phenoxy, naphthyloxy, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyridinyl, indolyl, quinolinyl, benzofuranlyl, benzthienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl or phenylamino;

30 R₃ is n-propyl, i-butyl, propenyl, i-butenyl or 2,2-dimethylpropyl;

R₂ and R₃ together with the carbon they are attached optionally form a ring selected from cyclopentyl, cyclohexyl, or cycloheptyl;

R₅ is methyl, ethyl, n-propyl, phenethyl, t-butyl, i-propyl, i-butyl, cyclohexyl, cyclohexylmethyl, benzyl, 4-fluorobenzyl, naphthylmethyl, acetyl, benzoyl or benzyloxycarbonyl.

36. A compound according to claim 20 selected from:

10 {{1-(4-Cyano-1-propyl-piperidin-4-ylcarbamoyl)-3,3-dimethyl-butylimino]-morpholin-4-yl-methyl}-carbamic acid ethyl ester;

N-(4-Cyano-methyl-piperidin-4-yl)-3-cyclohexyl-2-(3-oxo-3H-isoindol-1-ylamino)-propionamide;

15

4,4-Dimethyl-2-(3-oxo-3H-isoindol-1-ylamino)-pentanoic acid-(4-cyano-1-propyl-piperidin-4-yl)-amide;

20 N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-(2-oxo-2H-benzo[e][1,3] oxazin-4-ylamino)-propionamide;

{{1-(4-cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-piperidin-1-yl-methyl}-carbamic acid ethyl ester;

25

' 2-[(Acetylimino-phenyl-methyl)-amino]-N-(4-cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-propionamide;

30 '{{1-(4-Cyano-1-methyl-piperidin-4-ylcarbamoyl)-2-cyclohexyl-ethylamino]-morpholin-4-yl-methylen}-carbamic acid ethyl ester;

' 2-[(Acetylimino-phenyl-methyl)-amino]-N-[3-cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-propionamide;

35

'({1-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-ylcarbamoyl]-2-cyclohexyl-ethylamino}-morpholin-4-yl-methylene)-carbamic acid ethyl ester;

'N-(4-Cyano-1-methyl-piperidin-4-yl)-3-cyclohexyl-2-

40 [(ethylcarbamoylimino-phenyl-methyl)-amino]-propionamide;

'N-[4-Cyano-1-methyl-piperidin-4-yl]-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

5 'N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

10

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

N-(4-Cyano-1-propyl-piperidin-4-yl)-3-cyclohexyl-2-(1,1-dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-propionamide;

15

'2-(1,1-Dioxo-1H- λ^6 -benzo[d]isothiazol-3-ylamino)-4,4-dimethyl-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide

20 and the pharmaceutically acceptable derivatives thereof.

25 37. A pharmaceutical composition comprising a pharmaceutically effective amount of a compound according to claims 1, 6, 14, 20 or 32.

38. A method of modulating an autoimmune disease, said method comprising administering to a patient in need of such treatment a pharmaceutically effective amount of a compound according to claim 1, 6 or 20.

30

39. The method according to claim 38 wherein the autoimmune disease is rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, multiple sclerosis, Guillain-Barre syndrome, psoriasis, Grave's disease, myasthenia gravis, scleroderma, glomerulonephritis, atopic dermatitis or insulin-dependent diabetes mellitus.

35

40. A method of treating Alzheimer's disease comprising administering to a patient in need of such treatment a pharmaceutically effective amount of a compound according to claim 1, 6 or 20.


5

41. A method of treating atherosclerosis comprising administering to a patient in need of such treatment a pharmaceutically effective amount of a compound according to claim 1, 6 or 20.

10 42. A method of treating osteoporosis comprising administering to a patient in need of such treatment a pharmaceutically effective amount of a compound according to claims 14 or 32.

43. A method of treating asthma comprising administering to a patient in need of such
15 treatment a pharmaceutically effective amount of a compound according to claim 1, 6 or 20.

44. A method of making a compound of the formula (I)




20

wherein R_1 , R_2 , R_3 , R_4 , Het and R_5 are as defined in claim 1 and Y and X are both O;

said method comprising:

a) reacting an amino acid ester (IV), wherein R' is a protecting group, under suitable


5 reaction conditions with a $R_1C(O)L$ wherein L is a leaving group:



10 b) removing the protecting group R' from the compound produced in step a) to produce the corresponding carboxylic acid;


c) reacting the product of step b) under coupling conditions with an amino nitrile bearing "Het- R_5 " shown below to produce a compound of the formula (I):

15



20

45. A method of making a compound of the formula (I)




25 wherein R_1 , R_2 , R_3 , R_4 , Het and R_5 are as defined in claim 1 and Y and X are both O;

said method comprising:


a) reacting under coupling conditions a protected amino acid with a protecting group R' with an nitrile compound possessing "Het-R₅":

5




b) removing the protecting group R' from the compound produced in step a);

10 c) reacting the compound produced from step b) with an activated acid R₁C(Y)L wherein L is a leaving group, to produce a nitrile of the formula (I);




15 46. A method of making a compound of the formula (I)




wherein R₁, R₂, R₃, R₄, Het and R₅ are as defined in claim 1 and Y and X are both O;

said method comprising:


a) reacting ammonia with a protected amino acid bearing "Het-R₅" wherein R" is a protecting group, under suitable coupling conditions:




b) removing the protecting group R" from the product of step a) to produce compound III



c) reacting an amino ester of the formula (IV) with R₁C(O)L wherein L is a leaving group, in the presence of a suitable base, to produce compound (V);




d) hydrolysis of the product of step c) to produce carboxylic acid (VI);



e) reacting under suitable coupling conditions, the product (III) of step b) with the product (VI) of step d), followed by dehydration to produce the nitrile compound of the formula (I).


5 47. A method of making a compound of the formula (I)



wherein R₁, R₂, R₃, R₄, Het and R₅ are as defined in claim 1 and Y and X are both O;

10 said method comprising:

a) reacting under coupling conditions, an amino acid amide compound bearing "Het-R₅" with a protected amino acid bearing groups R₂, R₃:



b) removing the protecting group R' from the compound produced in step a);

c) reacting under suitable conditions the product of step b) with an $R_1C(O)L$ compound wherein L is a leaving group, followed by dehydration to produce a nitrile compound of the formula(I).

INTERNATIONAL SEARCH REPORT

Inte al Application No

PCT/US 00/23584

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 C07D413/12 C07D417/12 C07D205/04 C07D403/06
 A61K31/4535 A61P9/10 A61P25/28 A61P19/10 A61P11/06
 C07D211/66 C07D207/16 C07D309/14 C07D335/02 C07D401/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 776 718 A (KLAUS JEFFREY LEE ET AL) 7 July 1998 (1998-07-07) cited in the application abstract figures 1-8 -----	1,6,20, 37,38, 40-43

☐ Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

31 January 2001

Date of mailing of the international search report

12/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Diederer, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/23584

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5776718 A	07-07-1998	AU 713492 B	02-12-1999
		AU 5367496 A	16-10-1996
		CA 2216151 A	03-10-1996
		CN 1184472 A	10-06-1998
		CZ 9702981 A	18-03-1998
		EP 0817778 A	14-01-1998
		JP 11503417 T	26-03-1999
		NO 974403 A	17-11-1997
		PL 322409 A	19-01-1998
		WO 9630353 A	03-10-1996
		ZA 9602336 A	31-07-1996
